

Hypertensive Nephrosclerosis

Agnes Fogo, MD

The AJKD Atlas of Renal Pathology presents a compilation of figures on a specific pathologic entity. You may download the figures to create your own personal, non-commercial library of images or to create slides for teaching purposes.




Fig 1. Hypertensive nephrosclerosis is characterized by medial thickening and intimal fibrosis of medium-sized and larger vessels, arteriolar thickening, and hyalinosis, frequently associated with global glomerulosclerosis with obsolescence of glomeruli, as shown here. There is proportional tubulointerstitial fibrosis. (Periodic acid-Schiff stain, original magnification $\times 100$).




Fig 2. Hyalinosis is evident in the arteriole and interlobular artery in hypertensive nephrosclerosis, along with mild medial thickening caused by both hypertrophy and hyperplasia of vascular smooth muscle cells. The hyalin material is the result of insudation of plasma proteins, and is pink and glassy-smooth in appearance (ie, hyalin). (Periodic acid-Schiff stain, original magnification $\times 200$).

From the Department of Pathology, Vanderbilt University Medical Center, Nashville, TN.

Medical Photographer: Brent Weedman; with assistance from Kim Solez, MD, of the National Kidney Foundation's cyberNephrology™ Team.

Address author queries to Agnes Fogo, MD, Department of Pathology, Vanderbilt University Medical Center, MCN C-3310, Nashville, TN 37232. E-mail: Agnes.Fogo@vanderbilt.edu

Am J Kidney Dis. 33(6):E1-E2.

© 1999 by the National Kidney Foundation, Inc. Published by Elsevier Inc. All rights reserved.

0272-6386/\$36.00

[http://dx.doi.org/10.1053/S0272-6386\(13\)90017-2](http://dx.doi.org/10.1053/S0272-6386(13)90017-2)




Fig 3. Intimal fibrosis as well as medial thickening is evident in this artery in a patient with hypertensive nephrosclerosis. (Periodic acid-Schiff stain, original magnification $\times 200$).