

Malignant Hypertension

Agnes Fogo, MD

The AJKD Atlas of Renal Pathology presents a compilation of figures on a specific pathologic entity. You may download the figures to create your own personal, non-commercial library of images or to create slides for teaching purposes.




Fig 1. In malignant hypertension, also called accelerated hypertension, there are often underlying changes as seen in hypertensive nephrosclerosis, with superimposed fibrinoid necrosis of the vessel wall. In this vessel, there is also fibrin within the vessel wall, as a consequence of the necrosis. Glomeruli show only ischemic changes. (Jones' silver stain, original magnification $\times 100$).




Fig 2. Fibrinoid necrosis of the vessel wall with pink, amorphous fibrinoid material is present, with activated endothelial cells and inflammatory cells in this artery in a patient with malignant hypertension. There is also underlying medial thickening of the vessel wall, indicative of preexisting hypertensive injury. (Jones' silver stain, original magnification $\times 200$).

From the Department of Pathology, Vanderbilt University Medical Center, Nashville, TN.

Medical Photographer: Brent Weedman; with assistance from Kim Soletz, MD, of the National Kidney Foundation's cyberNephrology™ Team.

Address author queries to Agnes Fogo, MD, Department of Pathology, Vanderbilt University Medical Center, MCN C-3310, Nashville, TN 37232. E-mail: Agnes.Fogo@vanderbilt.edu

Am J Kidney Dis. 34(1):E1-E2.

© 1999 by the National Kidney Foundation, Inc. Published by Elsevier Inc. All rights reserved.

0272-6386/\$36.00

[http://dx.doi.org/10.1053/S0272-6386\(13\)90018-4](http://dx.doi.org/10.1053/S0272-6386(13)90018-4)




Fig 3. The glomeruli in malignant or accelerated hypertension show only ischemic change, seen here as extensive corrugation of the glomerular basement membrane with expanded lamina rara interna. (Transmission electron microscopy, original magnification $\times 8,000$).