

Sarcoidosis

Agnes Fogo, MD

The AJKD Atlas of Renal Pathology presents a compilation of figures on a specific pathologic entity. You may download the figures to create your own personal, non-commercial library of images or to create slides for teaching purposes.




Fig 1. Renal sarcoidosis is manifest by multiple sharply delineated non-necrotizing granulomas with frequent epithelioid-type multinucleated giant cells with accompanying interstitial fibrosis and lymphoplasmacytic interstitial infiltrate, as illustrated here. (Jones' Silver Stain, original magnification $\times 100$).




Fig 2. The well delineated non-necrotizing granulomas typical of sarcoidosis are illustrated here, with adjacent tubulointerstitial fibrosis. Giant cells are evident. (Jones Silver Stain, original magnification $\times 200$).

From the Department of Pathology, Vanderbilt University Medical Center, Nashville, TN.

Medical Photographer: Brent Weedman.


Address author queries to Agnes Fogo, MD, Department of Pathology, Vanderbilt University Medical Center, MCN C-3310, Nashville, TN 37232. E-mail: Agnes.Fogo@vanderbilt.edu

Am J Kidney Dis. 37(2):E12-E13.

© 2001 by the National Kidney Foundation, Inc. Published by Elsevier Inc. All rights reserved.


0272-6386/\$36.00

[http://dx.doi.org/10.1053/S0272-6386\(13\)90037-8](http://dx.doi.org/10.1053/S0272-6386(13)90037-8)




Copyright © 2001 by the National Kidney Foundation

Fig 3. The large giant cells comprising the non-necrotizing granulomas typical of renal sarcoidosis are evident in this biopsy, with surrounding lymphoplasmacytic interstitial infiltrate and early tubulointerstitial fibrosis. (Jones Silver Stain, original magnification $\times 200$).



Copyright © 2001 by the National Kidney Foundation

Fig 4. A well-delineated granuloma is seen surrounding an intact glomerulus in this case of renal sarcoidosis. There are numerous giant cells and surrounding lymphocytes and plasma cells with early interstitial fibrosis. (Hematoxylin and Eosin, original magnification $\times 200$).



Copyright © 2001 by the National Kidney Foundation

Fig 5. The multinucleated giant cell typical of sarcoidosis is shown under high magnification in this biopsy. (Periodic Acid Schiff, original magnification $\times 400$).