

Indinavir Nephropathy

Agnes Fogo, MD

The AJKD Atlas of Renal Pathology presents a compilation of figures on a specific pathologic entity. You may download the figures to create your own personal, non-commercial library of images or to create slides for teaching purposes.




Fig 1. Indinavir may cause a tubulointerstitial nephritis. In this case, there is chronic tubulointerstitial nephritis with interstitial fibrosis and marked lymphoplasmacytic infiltrate. (Masson's trichrome, original magnification $\times 100$).




Fig 2. Indinavir nephropathy is caused by intratubular precipitation of indinavir crystals with histiocytic reaction and sloughed degenerated tubular epithelial cells, as shown here. (Masson's trichrome, original magnification $\times 400$).

From the Department of Pathology, Vanderbilt University Medical Center, Nashville, TN.

Medical Photographer: Brent Weedman.

Address author queries to Agnes Fogo, MD, Department of Pathology, Vanderbilt University Medical Center, MCN C-3310, Nashville, TN 37232. E-mail: Agnes.Fogo@vanderbilt.edu

Am J Kidney Dis. 36(4):E22.

© 2000 by the National Kidney Foundation, Inc. Published by Elsevier Inc. All rights reserved.

0272-6386/\$36.00

[http://dx.doi.org/10.1053/S0272-6386\(13\)90033-0](http://dx.doi.org/10.1053/S0272-6386(13)90033-0)