

Acute Pyelonephritis

Agnes Fogo, MD

The AJKD Atlas of Renal Pathology presents a compilation of figures on a specific pathologic entity. You may download the figures to create your own personal, non-commercial library of images or to create slides for teaching purposes.




Fig 1. Acute pyelonephritis is diagnosed by intratubular aggregations of polymorphonuclear neutrophils (PMNs). There may be surrounding interstitial inflammation with a mixture of PMNs, lymphocytes, and plasma cells, but the predominant inflammation is within the tubules. (Jones' Silver Stain, original magnification $\times 200$).




Fig 2. More extensive intratubular polymorphonuclear neutrophils (PMNs) are present in this case of acute pyelonephritis. The PMNs are easily recognizable by their multilobed irregular nuclei. There is associated tubular degeneration and necrosis, but only minimal interstitial changes. (Jones' Silver Stain, original magnification $\times 200$).

From the Department of Pathology, Vanderbilt University Medical Center, Nashville, TN.

Medical Photographer: Brent Weedman.

Address author queries to Agnes Fogo, MD, Department of Pathology, Vanderbilt University Medical Center, MCN C-3310, Nashville, TN 37232. E-mail: Agnes.Fogo@vanderbilt.edu

Am J Kidney Dis. 35(2):E4.

© 2000 by the National Kidney Foundation, Inc. Published by Elsevier Inc. All rights reserved.

0272-6386/\$36.00

[http://dx.doi.org/10.1053/S0272-6386\(13\)90025-1](http://dx.doi.org/10.1053/S0272-6386(13)90025-1)