

Acute Tubular Necrosis

Agnes Fogo, MD

The AJKD Atlas of Renal Pathology presents a compilation of figures on a specific pathologic entity. You may download the figures to create your own personal, non-commercial library of images or to create slides for teaching purposes.




Fig 1. Acute tubular necrosis may be manifest only by regenerating flattened tubular epithelium without frank necrosis of individual tubular cells, as shown in some tubules to the right of the glomerulus, by blebbing and individual cell degeneration (top left), or by frank necrosis (middle). There is minimal accompanying interstitial infiltrate (periodic acid Schiff stain, $\times 100$).




Fig 2. There may also be degeneration and frank necrosis of individual cells or tubular segments in acute tubular necrosis, or flattened, regenerating type epithelium with degenerated cells in the lumen (middle left) (hematoxylin and eosin, $\times 200$).

From the Department of Pathology, Vanderbilt University Medical Center, Nashville, TN.

Medical Photographer: Brent Weedman.

Address author queries to Agnes Fogo, MD, Department of Pathology, Vanderbilt University Medical Center, MCN C-3310, Nashville, TN 37232. E-mail: Agnes.Fogo@vanderbilt.edu

Am J Kidney Dis. 41(1):E25-E26.

© 2003 by the National Kidney Foundation, Inc. Published by Elsevier Inc. All rights reserved.

0272-6386/\$36.00

[http://dx.doi.org/10.1053/S0272-6386\(13\)90049-4](http://dx.doi.org/10.1053/S0272-6386(13)90049-4)




Fig 3. Acute tubular necrosis is manifest by vacuolated cells and sloughed, necrotic cells in tubular lumina, with some tubules lined by flattened epithelium and some showing frank necrosis (periodic acid Schiff stain, $\times 400$).