

# Finite Observer Consensus as a Reconstruction Principle: Normal Forms, the Standard Model Lie Type, and a Route to the Einstein Field Equation

Bernhard Mueller<sup>1,\*</sup> Alexander Osika Mario Ponder Kai Xue  
Ben Cassie Peter Nguyen Maarten Antoine Visser Kale Arnav Anirudha  
David Matscheko Jonathan Hill William T. Glynn

August 1, 2026

**Paper release: r2011**

## Author affiliation:

<sup>1</sup> Pragma Research Inc., Washington State, United States

\*Corresponding author: [bernhard@floatingpragma.ai](mailto:bernhard@floatingpragma.ai).

## Abstract

We ask whether consensus among finite self-reading observer patches can constrain physical structure. Five layers are distinct: a self-referential closure hypothesis, three finite axioms, named repair and response laws, exact finite consequences, and physical realization maps. The patches carry state, records, overlap readback, and an oriented twelve-port boundary. We prove conditions for a canonical observable normal form, stable under refinement and independent of repair schedule. Complete reversible port response with endogenous overlap transport forces the compact Lie-algebra type  $\mathfrak{u}(1) \oplus \mathfrak{su}(2) \oplus \mathfrak{su}(3)$  without an ambient gauge group. A matter premise gives an anomaly-free fifteen-state exterior module with a  $\mathbb{Z}_6$  kernel. The spherical support carries Lorentz kinematics. Given one physical refinement tower with the displayed stress, entropy, continuum, and scale properties, null tomography and the Bianchi identity yield the Einstein field equation up to a cosmological constant. An icosahedral theorem fixes the first allowed anisotropic harmonic at angular rank six. On the registered primitive twelve-port propagation branch, the quartic coefficient fixes both sixth-order coefficients and the rotated rank-six template. This frozen prospective prediction differs from the zero-coefficient minimal locally Lorentz-invariant Standard Model plus General Relativity baseline. Its physical sector, coherent frame, exclusivity, continuum tower, and laboratory attachments enter as premises.

**Keywords:** foundations of physics; Observer Patch Holography; finite reconstruction; quantum records; Einstein field equation; gauge structure.

## 1 Introduction and claim boundary

Quantum probability, Lorentzian kinematics, gravitational dynamics, and gauge structure enter standard theories through distinct mathematical inputs [15, 16, 17, 38, 47]. Observer Patch Holography (OPH) asks which structures follow from a specified finite patch system whose components retain records, compare overlap readbacks, and repair disagreement. The source object is narrower than generic observer consistency: it includes a declared twelve-port carrier, complete reversible

response, endogenous overlap transport, explicit repair hypotheses, a finite algebra-state representation, and the interpretation maps stated with each physical conclusion.

The logical structure has five layers. First, the closure hypothesis says that the physical universe is a fixed point of its own internal description and repair; it is a structural premise beside the axioms. Second, A1 fixes the finite carrier and complete reversible-response tangent, A2 fixes agreement and endogenous overlap transport, and A3 selects the least-informative state left by those constraints. Third, named repair, response, and propagation laws act on the carrier. The axioms do not silently supply termination, confluence, or a physical kinetic operator. Fourth, finite theorems and certified computations state what follows from each displayed premise set. Fifth, physical realization maps identify finite records, currents, clocks, stress, and scales with laboratory or spacetime objects. The Lorentz statement is a group-theoretic consequence of the spherical support. The Einstein statement is tensor completion from a realized null balance, Ward conservation, the Bianchi identity, and an independent scale.

OPH is developed through a linked paper and artifact stack. This paper gives its synthesis and claim boundary; companion work supplies the finite formalization, exact-computation certificates, consensus construction, screen architecture, and simulation evidence [1, 5, 6, 7, 8, 11, 10, 2]. Together these sources specify a proposed observer-patch architecture and its conditional physical consequences. Each conclusion retains its own premises and verification artifacts, which readers can inspect directly.

Table 1 gives the seven status descriptions used throughout.

Table 1: Claim-status vocabulary used in the manuscript.

|                                             |                                                                                                                                                                                                                                               |
|---------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Finite theorem</b>                       | A deductive result proved from displayed premises on a stated finite domain. An exhaustive computation counts as part of the proof only with a specified domain and a soundness argument.                                                     |
| <b>Certified finite computation</b>         | An executable witness whose inputs, controls, and output are fixed by a reproducible certificate; execution alone does not promote its interpretation to a theorem.                                                                           |
| <b>Reconstruction implication</b>           | An exact implication whose antecedents include displayed geometric or physical premises.                                                                                                                                                      |
| <b>Open realization map</b>                 | A required map from a finite object to a physical event, current, field, clock, scale, or continuum structure for which no construction is asserted.                                                                                          |
| <b>Diagnostic</b>                           | A certified computation compared against measured values that it consumes; never counted as a prediction.                                                                                                                                     |
| <b>Registered conditional test</b>          | A diagnostic with a custody-bound target definition, kill band, precision floor, and decision rule fixed before comparison.                                                                                                                   |
| <b>Frozen prospective branch prediction</b> | A source calculation and decision rule fixed before any eligible comparison, conditional on a named physical branch. Failure rejects that branch; its scope extends to the whole framework only if the branch is proved forced and exclusive. |

Throughout, “source” means computed from the declared finite source specification, including every explicitly named source law, with no measured physical input. “Source-only” marks a quantity whose entire dependency cone is of that kind. A physical realization map is not source-only merely because the finite branch calculation is target-clean.

The distinction is mathematical: a finite implication and a physical realization of its antecedents are different claims. Machine-readable receipts record that distinction and reject a physical verdict when a required map is absent.

*Boundary 1.1.* This paper claims no completed derivation of particle masses, no source-only determination of the fine-structure constant, no completed cosmology, and no claim that formal verification establishes physical truth. Quantitative surfaces that consume measured values are labeled diagnostics. The primitive twelve-port propagation branch is a frozen prospective conditional prediction. Section 12 states its decision rule and the physical premises that set the scope of any verdict.

**Definition 1.2** (Physical realization map). A physical realization map assigns finite public records, event data, currents, modular generators, and scales to physical records, spacetime events, laboratory currents, clocks, and units. It must preserve the algebraic operations, overlap restrictions, causal order, normalizations, and refinement relations consumed by the conclusion in question. A resemblance of spectra, dimensions, or symmetry names is insufficient.

## 1.1 Principal contributions

The observer-indexed normal-form framework separates existence, observable determination, schedule independence, stability, and computational cost. It gives a canonical partial normalizer, a sharp two-output stability bound, an accumulated refinement bound, and complexity barriers for succinct boundary problems. On the declared transactional repair branch, semantic dependency closure and revalidation yield the local diamond and hence a schedule-independent public record.

Complete reversible response gives a faithful compact twelve-dimensional unitary current algebra. Endogenous overlap transport makes every proper carrier action inner on that same algebra. Its one-dimensional fixed space and compact-simple classification force  $\mathfrak{u}(1) \oplus \mathfrak{su}(2) \oplus \mathfrak{su}(3)$  without an ambient gauge-group premise. A declared four-band matrix map is an exact conditional witness for this type. Under a separate trace-balanced matter contract, the exterior module gives a faithful image with  $\mathbb{Z}_6$  cover kernel and an anomaly-free fifteen-state representation. The source contains no selection of the matrix current, physical matter action, global quotient, or laboratory current; the physical reading consumes these as premises.

The completed finite record surface carries exact projection-event, conditioning, expectation, and Clauser–Horne–Shimony–Holt identities. The geometry branch distinguishes observer-velocity kinematics from event geometry. Its Einstein result is a reconstruction implication whose modular, stress, entropy, area, Ward, continuum, and scale premises are displayed in one contract.

The quantitative section proves the positive-chamber Koide identity and the capacity-transfer sign, states the conditional tau test and charged-lepton diagnostic, and derives the primitive twelve-port propagation prediction. The latter fixes the two sixth-order coefficients and the rotated angular-rank-six template from the quartic coefficient. Its custody record excludes every data product used by the related exploratory cosmic-microwave-background search.

Operational quantum reconstructions [15, 16, 17], relational and modular accounts of physical facts and time [27, 28], thermodynamic routes to gravitational dynamics [38, 39], and compact-group reconstruction [46, 47] provide the principal comparison points. Born and Lüders rules, Tsirelson’s inequality, conformal–Lorentz isomorphisms, Tannaka reconstruction, and the familiar one-generation exterior representation are used as established ingredients. The originality asserted here is limited to the observer-fiber normalizer with sharp refinement control, the transaction contract yielding the local diamond, the A1–A2 forcing theorem with its fixed-space exclusion, the conditional four-band witness and exhaustive exterior selection, the nine-direction null design with exact determinant and decoder constants, and the frozen coefficient-linked primitive-port prediction.

## 1.2 The three axioms

The model uses the following three mathematical postulates.

**A1 (oriented twelve-port observer screen).** In plain language, the model assigns every carrier a finite observer screen with local state, readback, records, repair moves, checkpoints, typed overlaps, and an oriented twelve-port boundary. Concisely, for each regulator  $r$  there is a typed object

$$\mathfrak{N}_r = (\mathcal{P}_r, \mathcal{A}_r, \mathcal{R}_r, \mathcal{I}_r, \mathcal{U}_r, \mathcal{C}_r, N_r, S_r, b_r)$$

whose carrier boundary  $K_{r,i} = (P_{r,i}, E_{r,i}, F_{r,i}, o_{r,i})$  has  $|P| = 12$ ,  $|E| = 30$ ,  $|F| = 20$ , and the oriented icosahedral incidence relations. The bridge  $b_r : N_r \rightarrow S_r$  carries a designated oriented two-cycle to the fundamental class of the spherical support, and the typed maps commute with refinement. For each complete carrier, put  $V_{r,i} = \mathbb{R}^{P_{r,i}}$ . A1 supplies a finite-dimensional unitary space  $H_{r,i}$  and an injective real-linear response derivative

$$D_{r,i} : V_{r,i} \longrightarrow \mathfrak{u}(H_{r,i})$$

whose image  $\mathfrak{g}_{r,i} = D_{r,i}(V_{r,i})$  is closed under commutators. The primitive port probes span  $V_{r,i}$ , their ordered compositions generate every accepted infinitesimal reversible response, and no public response direction is omitted. The pairing

$$\langle v, w \rangle_D = -\text{Tr}(D_{r,i}(v)D_{r,i}(w))$$

is positive definite. The response data and their completeness commute with refinement, and

$$G_{D,r,i}^0 = \langle \exp(tD_{r,i}(v)) : t \in \mathbb{R}, v \in V_{r,i} \rangle^0$$

is the connected response group. A1 constrains the finite carrier, federation, operational interfaces, spherical-support tower, and complete public reversible-response tangent. It does not imply semantic agreement, repair termination, confluence, a particular matrix response, an inverse-port law, a compact Lie type, a global group, or a laboratory identification [4].

**A2 (observer agreement).** In plain language, observers assign the same operational meaning to accepted data on their shared boundary. Every proper recharting of a complete carrier is implemented by their own reversible overlap response. Formally, the interpretation functor

$$\mathcal{J}_r : \text{Data}_r \longrightarrow \text{Meaning}_r$$

is natural under every visible restriction, recharting, seam translation, higher-overlap map, federation map, and refinement map. On an overlap  $O$ ,

$$\mathcal{J}_O(\text{res}_{P \rightarrow O} d_P) = \mathcal{J}_O(\tau_{Q \rightarrow P} \text{res}_{Q \rightarrow O} d_Q).$$

The accepted reversible overlap transports form a groupoid  $\mathcal{O}_r$ . For a complete carrier chart  $o$ , let  $\text{Hol}_r(o)$  be its closed overlap paths. A2 requires the port projection

$$\Pi_{r,i} : \text{Hol}_r(o) \longrightarrow \text{Aut}^+(K_{r,i})$$

to be surjective. For every proper carrier automorphism  $a \in \text{Aut}^+(K_{r,i})$ , A2 supplies a closed path  $\gamma_a$  and a projective unitary implementer  $[U_a]$  satisfying

$$\Pi_{r,i}(\gamma_a) = a, \quad \text{Ad}_{[U_a]} D_{r,i}(v) = D_{r,i}(a \cdot v).$$

The implementer is endogenous to the same response:

$$[U_a] = [g_a c_a], \quad g_a \in G_{D,r,i}^0, \quad c_a \in C_{U(H_{r,i})}(\mathfrak{g}_{r,i}).$$

The centralizer factor acts trivially on  $\mathfrak{g}_{r,i}$ . A2 constrains operational meaning on accepted shared data and makes proper carrier rechartings inner on the complete port-response algebra. It does not imply global state extension, termination, confluence, unique normal forms, durable records, a global compact group, or a laboratory current [4].

**A3 (maximum randomness under agreement constraints).** In plain language, the selected state is least informative relative to the declared reference after every observer-visible constraint has been imposed. At finite regulator  $r$ , let  $\mathcal{K}_r$  be the nonempty convex set of compatible local state families and let

$$\mathcal{D}_r(\rho \parallel \tau_r) = \sum_{P \in \mathcal{G}_r} w_{r,P} D(\rho_{r,P} \parallel \tau_{r,P}), \quad w_{r,P} > 0.$$

The observer cover is state-determining on  $\mathcal{K}_r$ , and

$$\rho_r = \arg \min_{\rho \in \mathcal{K}_r} \mathcal{D}_r(\rho \parallel \tau_r).$$

A3 constrains state selection inside one A1-fixed feasible space. It does not select a field list, repair law, response map, particle multiplicity, or continuum limit [4].

None of the axioms contains a gauge group, a particle list, an event-manifold dimension, or a recovery law. A1 does fix a two-dimensional spherical support and hence the Lorentz kinematic type used below. The results below state which structures follow from these axioms, which require added premises, and which realization maps are consumed as premises.

### 1.3 The self-referential closure hypothesis

One structural hypothesis sits beside the axioms. The universe is modeled as a timeless, closed, self-referential mathematical structure: the simulating description and the simulated system are one object, and the physical configuration is a fixed point  $T(\mathfrak{U}) = \mathfrak{U}$  of a universe-level closure operator. There is no global clock anywhere in the model. Repair order is a partial order on commits, and time is emergent per observer, read along each observer's own chain of completed records.

On the reading adopted here, a structure closed under its own description contains subsystems that read, record, compare, and reconstruct it. Within their emergent subjective time, observers assemble a consistent account of the structure and recover its architecture. The closure condition requires the rebuilt description and the inhabited structure to return the same invariant quantities. It supplies no finite repair law, current derivative, physical kinetic operator, or existence and uniqueness proof for the cosmic fixed point.

Two finite results give the hypothesis exact content on declared branches. The packet-quotient closure theorem proves, on the declared finite consensus branch, that the closure map is an affine idempotent whose fixed points are exactly the packets supported on consensus normal forms; its habitat-level extension is typed open [7]. The universe-level record-closure equation  $N = \log M_0(\mathfrak{U}_N)$  is posed exactly, and the complete declared direct source class is proved non-identifying, so any successful completion requires an additional source law whose two readings agree [9]. The A2 endogeneity clause is the hypothesis's local expression within the axioms: every proper recharting is an operation implemented from inside the response.

The hypothesis also fixes how the declared carrier is to be read. Under closure there is no outside vantage from which an architecture could be tuned; the twelve-port carrier is part of the solution that observers inside the structure reverse engineer, and the axioms are the recovered description of that solution. This reading motivates the declaration and is typed as a hypothesis: the theorems of this paper consume the declared carrier as a premise and stand without the reading, and no uniqueness theorem over carriers is claimed. The quantitative closure candidates the hypothesis generates, the screen-grain equation and the capacity coordinate, are developed with full typing in Section 9.1. Existence, uniqueness, and stability of the physical fixed point are separate determinacy tests, and the finite theorems prove closure only on their declared finite branches.

## 1.4 How the constraints work together

Physical structure is selected at the intersections of constraints. The carrier incidence fixes a finite response space, its symmetry action, and its invariant subspaces. By itself, that geometry admits many equivariant response laws. A1 restricts the menu to faithful, complete, compact, commutator-closed reversible responses. A2 requires every proper recharting visible through overlap agreement to act internally on that same response. For the twelve-port carrier, the resulting one-dimensional fixed space excludes the sole centreless compact alternative and fixes the local Standard Model gauge algebra. Geometry, reversible dynamics, and agreement are all used in this conclusion.

Repair dynamics occupy a distinct part of the architecture. A named repair law produces a schedule-independent public record only when its termination and local-diamond premises hold. The simulator can test whether a proposed law realizes those premises and whether its records support the response and holonomy interfaces consumed by later branches. The gauge theorem does not silently import such a law: it consumes the complete response clause of A1 and the endogenous transport clause of A2. Source derivation of those two objects is a separate realization problem.

The same rule applies to measures: carrier counting and algebraic traces supply finite candidate measures, agreement and repair determine whether they descend to the public quotient, and a physical realization map must then preserve them as spacetime or laboratory measures. This dependency discipline separates forced structure from formal resemblance.

## 1.5 The theorem chain

The results form branches with different premises:

|                                                |   |                                               |
|------------------------------------------------|---|-----------------------------------------------|
| finite carrier and convergent repair           | → | consensus normal form,                        |
| algebra-state reading of the completed record  | → | finite quantum record identities,             |
| A1 response and A2 transport                   | → | Standard Model local Lie type,                |
| displayed matter representation                | → | conditional $\mathbb{Z}_6$ matter image,      |
| oriented spherical support                     | → | Lorentz observer-velocity kinematics,         |
| realized null balance, Ward/Bianchi, and scale | ⇒ | $G_{ab} + \Lambda g_{ab} = 8\pi G_N T_{ab}$ . |

The first four rows are finite results with the displayed premises. The fifth consumes the declared spherical support. The sixth is a reconstruction implication. A common modular–stress–entropy tower is the proposed route to its null-balance antecedent; its physical realization is an open realization map. Sections 2 through 9 develop these branches. Section 10 records the physical interpretation maps. Section 11 describes the verification stack. Section 12 states the falsification architecture and the empirical decision rule. Section 13 compares the program with neighboring

approaches. The conclusion states the program’s full ambition, a derivation of the constants and structural properties of nature from consistency requirements alone.

## 2 The primitive finite architecture

**Definition 2.1** (Observer patch). An observer patch is a finite object with local state, an observable boundary map, durable records, readback, and a set of authorized repair moves. A patch sees a fragment of the world: its own state and the boundary data of its authorized overlaps.

**Definition 2.2** (Overlap and repair). For patches  $x_i, x_j$  with an authorized overlap  $e = (i, j)$ , both induced boundary records must agree for the pair to hold public data. A repair move changes local state while preserving protected readout. A configuration is a normal form when no authorized repair applies.

The carrier realization used throughout is the twelve-port icosahedral architecture of A1: each carrier’s boundary packet is combinatorially the oriented icosahedron boundary  $(P, E, F, o)$  with  $|P| = 12$ ,  $|E| = 30$ ,  $|F| = 20$ ; carriers federate through typed seam algebras with coherent triple-overlap cocycles into a nerve carrying a degree-one bridge to the oriented spherical support. This is the declared source architecture. We do not claim a uniqueness theorem selecting it among all possible carriers; where results depend on it, the dependence is displayed.

Two exact features of this architecture enter the later constructions. First, the proper rotation group of the icosahedral packet is the alternating group  $A_5$ , with sixty rotations acting on the twelve ports; the real port-coefficient space has the character decomposition

$$P_{12} \cong_{A_5} \mathbf{1} \oplus \mathbf{3} \oplus \mathbf{3}' \oplus \mathbf{5}, \quad (1)$$

an exact multiplicity-free decomposition obtained by standard finite-group character theory [33] and checked by certified projectors. Second, record keeping is integer valued: a write appends  $+1$ , a retraction appends  $-1$ , and readback sums atomic events per port; a conservative repair transfers one unit across a seam whenever the oriented mismatch has magnitude at least two. With  $V(N) = \sum_i N_i^2$ , each such repair strictly decreases  $V$  by  $2(d - 1)$  for mismatch  $d \geq 2$ , so repair terminates by a finite theorem. Divisibility of total load by twelve is a necessary consensus condition and is not sufficient by itself. For the certificate’s source-generated full-pile packet, an explicit eighteen-move schedule reaches consensus.

### 2.1 Hardware and protocol: one substrate, typed dynamics, several readouts

The finite substrate and the operations acting on it are distinct parts of the model. Every physical conclusion requires a specified operation and its own realization map.

The *substrate and reversible-response tangent* are the content of A1: a federation of finite carriers, each with the oriented twelve-port icosahedral boundary packet, joined through typed seam algebras with coherent triple overlaps into a nerve that bridges to the oriented spherical support, together with the complete reversible response space of each carrier. The companion microphysics paper specifies this substrate at the level of patch hardware, records, and synchronization interfaces [8]. This layer fixes what exists at each finite regulator: ports, seams, records, and response directions. It does not select a repair schedule, convergent repair law, or physical propagation operator.

The *agreement and state-selection protocol* combines A2 and A3 with named branch dynamics. Patches write and retract integer record events, read back port sums, compare induced records

on shared boundaries, and interpret accepted data through the natural functor of A2. A named repair law moves oriented mismatches under quadratic descent; its termination and local-diamond premises are proved separately. A3 selects the residual state by constrained information projection after the agreement constraints are fixed. The companion consensus paper develops this layer as an asynchronous agreement protocol with explicit safety and liveness assumptions [7]. The protocol fixes what happens: which configurations count as repaired, which records become public, and which transports implement recharting.

The claim structure of the paper is that distinct readouts of this one protocol-on-hardware system yield the distinct pillars of observed physics. Table 2 states the correspondence and names the section that proves each step.

Table 2: The reconstruction pipeline: typed operation, finite substrate, resulting structure, and its location in this paper.

| Operation or branch premise                                                   | Finite substrate                          | Resulting structure                                                                                                | Where |
|-------------------------------------------------------------------------------|-------------------------------------------|--------------------------------------------------------------------------------------------------------------------|-------|
| Repair to a fixed point under quadratic descent and the transactional diamond | carrier records and seams                 | schedule-independent public record; measurement as completed consensus                                             | §3    |
| Algebra-state reading of the completed record surface                         | protected record projectors               | Born weights, Lüders conditioning, Tsirelson bound                                                                 | §4    |
| Overlap transport around closed paths                                         | seam groupoid and triple-overlap cocycles | central defects, edge sectors, edge-center entropy                                                                 | §5    |
| Record-germ separation along refinement                                       | federation nerve and spherical bridge     | Lorentz kinematics; conditional Lorentzian event manifold                                                          | §6    |
| Modular flow, entropy stationarity, null tomography on one tower              | the same record tower                     | conditional Einstein field equation                                                                                | §7    |
| Endogenous recharting of the complete port response                           | twelve-port incidence and response space  | forced gauge Lie type $\mathfrak{u}(1) \oplus \mathfrak{su}(2) \oplus \mathfrak{su}(3)$ ; conditional matter image | §8    |
| Capacity accounting and closure readback                                      | screen capacity and grain                 | exact quantitative closures and physical tests                                                                     | §9    |
| Equal-weight primitive-port propagation with coherent physical frame          | the twelve-port direction orbit           | linked quartic and sixth-order coefficients with one rotated angular-rank-six template                             | §9.6  |

Two features of this pipeline carry the unification claim. The branches consume one substrate with typed operations: the same completed records that define measurement feed the event geometry, and the same overlap transport that defines agreement feeds the gauge forcing. Every arrow in the table is typed as a finite theorem, a certified computation, or an open realization map. The universe claim of the program is the conjunction: a physical carrier that realizes the hardware and runs the protocol produces public records whose readouts include quantum probability, Lorentzian causal structure, Einstein dynamics, and the Standard Model gauge and matter skeleton, with the listed open realization maps as its premises.



### 3 Observable normal forms and finite consensus

A completed finite configuration  $x$  is a fixed point of the named repair-and-agreement map:

$$\mathcal{R}_{\text{cons}}(x) = x, \quad (2)$$

where  $\mathcal{R}_{\text{cons}}$  is the composite finite dynamics. This map is distinct from the universe-level closure operator  $T$  of Section 1.3. Fixed-point existence, observable determination, and schedule independence are separate questions. The following framework isolates them [5].

#### 3.1 Exact observable fibers

**Definition 3.1** (Observable quotient system). An observable quotient system is a tuple

$$\mathfrak{S} = (Q, C, \mathcal{B}, B),$$

where  $Q$  is a set of configurations,  $C \subseteq Q$  is the consistent subset,  $\mathcal{B}$  is a set of protected records, and  $B : Q \rightarrow \mathcal{B}$  is the record map. For  $b \in \mathcal{B}$ , write  $C_b = C \cap B^{-1}(b)$ .

The fiber  $C_b$  is unrealizable, reconstructing, or ambiguous according as its cardinality is zero, one, or greater than one. This trichotomy avoids assigning a state to a record that has either no consistent extension or several observationally indistinguishable extensions.

**Theorem 3.2** (Canonical observable normalizer). *The following statements are equivalent:*

- (i)  $B|_C$  is injective;
- (ii) there is a unique map  $N : Q \rightarrow C \sqcup \{\perp\}$  that preserves  $B$  on its  $C$ -valued outputs, fixes every point of  $C$ , is constant on each  $B$ -fiber, and returns  $\perp$  exactly on fibers with no consistent extension;
- (iii)  $B|_C : C \rightarrow B(C)$  is a bijection.

If  $B(Q) \subseteq B(C)$ , the map is a total idempotent retraction  $Q \rightarrow C$ .

*Proof.* Injectivity makes every nonempty  $C_b$  a singleton, which defines  $N$  and forces all its properties fiber by fiber. Conversely, if  $c, c' \in C$  have  $B(c) = B(c')$ , the fixed-point and fiber clauses give  $c = N(c) = N(c') = c'$ . The third statement is the restriction of the first to the image  $B(C)$ .  $\square$

This normalizer is defined by observable fibers, independent of a chosen repair schedule. A repair relation implements it only when the relation preserves  $B$ , reaches the consistent set, and satisfies an appropriate liveness condition. Confluence from one source does not by itself prove determination across distinct sources with the same protected record.

#### 3.2 Approximate outputs and refinement

Let  $(Q, d_Q)$  be finite, let  $(\mathcal{B}, d_B)$  be metric, and let  $C = \Phi^{-1}(0)$  be nonempty for a nonnegative consistency residual  $\Phi$ . For  $t, r \geq 0$ , define

$$\eta_\Phi(t) = \max\{\text{dist}_Q(x, C) : \Phi(x) \leq t\}, \quad (3)$$

$$\omega_B(r) = \max\{d_Q(c, c') : c, c' \in C, d_B(Bc, Bc') \leq r\}. \quad (4)$$

The first modulus turns residual error into distance from consistency. The second measures how well protected observations distinguish consistent outputs.

**Theorem 3.3** (Sharp two-output stability bound). *Suppose  $B$  is  $L_B$ -Lipschitz. If*

$$\delta_x, \delta_y, \varepsilon \geq 0, \quad \Phi(x) \leq \delta_x, \quad \Phi(y) \leq \delta_y, \quad d_B(Bx, By) \leq \varepsilon,$$

*then*

$$d_Q(x, y) \leq \eta_\Phi(\delta_x) + \eta_\Phi(\delta_y) + \omega_B(\varepsilon + L_B[\eta_\Phi(\delta_x) + \eta_\Phi(\delta_y)]). \quad (5)$$

*At zero residual the  $\omega_B$  term is exact, and the coefficient one on each of the two residual-distance terms is optimal in the class of finite metric systems.*

*Proof.* Choose nearest consistent points  $c_x, c_y$ . The two residual moduli bound the outer legs  $x \rightarrow c_x$  and  $c_y \rightarrow y$ . Lipschitz continuity bounds  $d_B(Bc_x, Bc_y)$  by the argument of  $\omega_B$  in (5); the inverse-observation modulus bounds the middle leg. The triangle inequality gives the result. Exactness at zero residual follows from the definition of  $\omega_B$ . To test the first residual coefficient, take the two-point metric  $Q = \{c, x\}$  with  $C = \{c\}$ ,  $d_Q(c, x) = s$ , constant  $B$ ,  $y = c$ ,  $\Phi(c) = 0$ , and  $\Phi(x) = 1$  at  $(\delta_x, \delta_y, \varepsilon) = (1, 0, 0)$ . The bound reduces to  $s \leq \eta_\Phi(1) = s$ , so that coefficient cannot be smaller than one. Exchanging  $x$  and  $y$  proves the same statement for the second coefficient.  $\square$

Let  $(Q_n, d_n)$  be nonempty finite metric spaces. For  $m \geq j \geq n$ , let  $\rho_{m,n} : Q_m \rightarrow Q_n$  be restriction maps satisfying  $\rho_{n,n} = \text{id}_{Q_n}$  and  $\rho_{m,n} = \rho_{j,n} \rho_{m,j}$ . Let  $N_n : Q_n \rightarrow Q_n$  be total level normalizers. Set

$$a_j = \max_{q \in Q_{j+1}} d_j(\rho_{j+1,j} N_{j+1} q, N_j \rho_{j+1,j} q), \quad K_{j,n} = \text{Lip}(\rho_{j,n}).$$

**Theorem 3.4** (Accumulated refinement bound). *For  $m > n$  and  $q \in Q_m$ ,*

$$d_n(\rho_{m,n} N_m q, N_n \rho_{m,n} q) \leq \sum_{j=n}^{m-1} K_{j,n} a_j.$$

*In particular, if the restrictions are nonexpansive and  $\sum_{j=0}^{\infty} a_j < \infty$ , then*

$$\sup_{m > n} \max_{q \in Q_m} d_n(\rho_{m,n} N_m q, N_n \rho_{m,n} q) \leq \sum_{j=n}^{\infty} a_j \longrightarrow 0 \quad (n \rightarrow \infty).$$

*This is asymptotic naturality of the normalizers.*

*Proof.* Insert the intermediate points  $p_j = \rho_{j,n} N_j \rho_{m,j} q$ . Consecutive points differ by at most  $K_{j,n} a_j$ . Summing the telescoping chain from  $p_m$  to  $p_n$  gives the bound.  $\square$

### 3.3 Computational boundary

The normalizer can be canonical without being cheap. For a Boolean circuit  $F : \{0, 1\}^n \rightarrow \{0, 1\}$ , take  $C = F^{-1}(1)$  and let  $B_S$  project onto a declared coordinate set  $S$ .

**Theorem 3.5** (Complexity of succinct boundary reconstruction). *For succinct Boolean boundary systems:*

(i) *deciding whether a protected record has a consistent extension is NP-complete;*

- (ii) deciding whether  $B_S|_C$  is injective is coNP-complete, even when all but one variable are observed;
- (iii) deciding whether every protected collar value has some consistent interior extension, equivalently whether an unrestricted total strong repair exists, is  $\Pi_2^P$ -complete.

*Proof.* For (i), a consistent extension is a polynomial witness and satisfiability is the case  $S = \emptyset$ . For (ii), noninjectivity is witnessed by two distinct satisfying assignments with the same projection; unsatisfiability reduces to injectivity after adding one hidden tie-breaking bit and one always-consistent baseline assignment. For (iii), a repair exists exactly when the corresponding circuit relation satisfies  $\forall d \exists w R(w, d) = 1$ , the canonical one-alternation quantified Boolean formula problem.  $\square$

### 3.4 Repair realization on the twelve-port carrier

For a prepared batch of support-closed transactions, join  $\tau$  and  $\sigma$  in the conflict graph when

$$W_\tau \cap (R_\sigma \cup W_\sigma) \neq \emptyset \quad \text{or} \quad W_\sigma \cap (R_\tau \cup W_\tau) \neq \emptyset.$$

The conflict components below are obtained by iterating the following finite closure: compute semantic support to a fixed point inside every aggregate, rebuild the conflict graph on the expanded read and write sets, merge every connected component, and repeat until the partition and all aggregate supports are unchanged.

**Proposition 3.6** (Transactional local diamond). *Let an accepted repair be an aggregate transaction with read set  $R_\tau$ , write set  $W_\tau$ , a read snapshot, and a snapshot-determined payload. Assume:*

- (i) *every read set contains the semantic dependency closure of its write set, and every affected acceptance functional is revalidated at commit;*
- (ii) *the terminal closure above exists, and each of its connected conflict components has one coherent canonical aggregate payload;*
- (iii) *a prepared component whose snapshot remains valid survives commits of independent components; and*
- (iv) *accepted commits preserve the protected record and strictly decrease the integer descent functional.*

*Then every one-step quotient peak*

$$t \longleftarrow s \longrightarrow u$$

*has a quotient state  $v$  with  $t \longrightarrow v \longleftarrow u$ .*

*Proof.* Two different first steps cannot originate in one conflict component because that component has a unique aggregate payload. They therefore originate in distinct components of the terminal conflict closure. Rebuilding the graph after every aggregate support expansion makes their final read and write sets satisfy

$$W_\tau \cap (R_\sigma \cup W_\sigma) = \emptyset, \quad W_\sigma \cap (R_\tau \cup W_\tau) = \emptyset.$$

Their writes are disjoint and neither commit changes the other's read snapshot. Semantic dependency closure ensures that neither commit changes an acceptance functional consumed by the other.

Revalidation and the surviving component rule therefore make both second commits legal. Since the payloads are snapshot-determined and the writes are disjoint,

$$\text{Apply}_\sigma \text{Apply}_\tau(s) = \text{Apply}_\tau \text{Apply}_\sigma(s) = v.$$

Protected-record preservation makes this equality valid on the physical quotient.  $\square$

**Theorem 3.7** (Consensus normal form). *Suppose the accepted repair relation satisfies Proposition 3.6, the exact quadratic descent of Section 2, and repair completeness, meaning that its normal forms are exactly the consistent states. Every maximal asynchronous repair schedule from one initial state then terminates at the same consistent quotient normal form. The induced global repair map is idempotent, fixes consistent states, and preserves the protected record.*

*Proof.* Quadratic descent makes the accepted relation terminating. Proposition 3.6 makes it locally confluent, so Newman’s lemma [26] makes it confluent. A terminating confluent relation has a unique normal form below each source. Repair completeness puts that normal form in the consistent set. Applying the normalizer a second time does nothing, and protected-record preservation holds along every accepted edge.  $\square$

*Boundary 3.8.* Atomic commits, disjoint write sets, or termination alone do not imply the local diamond. Countermodels omitting semantic dependency closure, canonical aggregation, or the surviving component rule are retained with the finite receipt. The reference engine exhaustively checks its states, peaks, payload hashes, protected records, and descent comparisons. The theorem is finite and asserts no continuum limit.

The declared class also has gauge compatibility (repair commutes with local relabeling), idempotence of the completed dynamics, and refinement compatibility where proved. The public object is the protected quotient record, not an arbitrary intermediate configuration.

The architecture sits close to distributed agreement in the tradition of Lamport, Shostak, and Pease [24]: patches play the role of protocol nodes, overlap repair of a quorum vote, and the completed fixed point of the decided state. A component paper of the stack proves safety and liveness for a Byzantine-tolerant reading of repair under explicit quorum and partial-synchrony assumptions [7]; the impossibility boundary of Fischer, Lynch, and Paterson [25] is the reason those assumptions are named.

## 4 Finite record algebra and quantum identities

Once a compare, write, and verify slice has completed, its accessible events are represented in a finite-dimensional  $\ast$ -algebra with a normalized state. This algebra-state representation is an explicit input at this stage. The consensus theorem identifies which record is public; it does not derive the classification of finite-dimensional  $C^\ast$ -algebras or the trace pairing.

**Theorem 4.1** (Finite public-record identities). *Let  $P_E$  be the projector of a completed public event  $E$  and  $\rho$  the normalized state of the completed record surface. The event weight and nonzero-weight update*

$$\Pr(E) = \text{Tr}(\rho P_E), \quad \rho|E = \frac{P_E \rho P_E}{\text{Tr}(\rho P_E)} \quad (6)$$

*define a normalized probability measure on every orthogonal event partition and a normalized positive state on the post-event corner. For self-adjoint dichotomic observables  $A_0, A_1$  and  $B_0, B_1$  in commuting record subalgebras, with  $A_i^2 = B_j^2 = I$ , define*

$$S_{\text{CHSH}} = \text{Tr}[\rho\{A_0(B_0 + B_1) + A_1(B_0 - B_1)\}].$$

Then

$$|S_{\text{CHSH}}| \leq 2\sqrt{2}. \quad (7)$$

*Proof.* Positivity of  $\rho$  and  $0 \leq P_E \leq I$  give  $0 \leq \text{Tr}(\rho P_E) \leq 1$ . For a complete orthogonal partition  $(P_k)$ , linearity and  $\sum_k P_k = I$  give  $\sum_k \text{Tr}(\rho P_k) = 1$ . If the event weight is nonzero,  $P_E \rho P_E / \text{Tr}(\rho P_E)$  is positive, normalized, and supported on  $P_E \mathcal{A} P_E$ .

For  $\mathcal{B} = A_0(B_0 + B_1) + A_1(B_0 - B_1)$ , the commuting-subalgebra calculation gives

$$\mathcal{B}^2 = 4I - [A_0, A_1][B_0, B_1].$$

The two commutator norms are at most two, so  $\|\mathcal{B}^2\| \leq 8$  and  $\|\mathcal{B}\| \leq 2\sqrt{2}$ . Taking its expectation in  $\rho$  proves the stated bound.  $\square$

The first identity is the Born rule on this finite surface [18]; the second is Lüders conditioning [19]; the third is the Tsirelson bound [21] for the Clauser–Horne–Shimony–Holt combination [20] at fixed cutoff. Compatibility structure for commuting events follows the standard finite operational pattern within the same library.

**Theorem 4.2** (Two-level partition expectation). *Let  $(P_i)_{i=1}^k$  be pairwise orthogonal projectors with  $\sum_i P_i = I$ . Define*

$$\mathcal{P}(X) = \sum_i P_i X P_i, \quad \mathcal{A}(X) = \sum_i \frac{\text{Tr}(X P_i)}{\text{Tr}(P_i)} P_i,$$

*with a zero projector contributing zero. Then  $\mathcal{P}$  and  $\mathcal{A}$  are positive, unital, trace-preserving idempotent linear maps. The exact range of  $\mathcal{P}$  is the commutant of the partition; the exact range of  $\mathcal{A}$  is the commutative span of the  $P_i$ ; and*

$$\mathcal{A}\mathcal{P} = \mathcal{A} = \mathcal{P}\mathcal{A}.$$

*The average preserves every partition-event weight. For a partition member of nonzero weight, averaging commutes with Lüders conditioning and both orders give the normalized projector.*

*Proof.* Orthogonality gives  $\mathcal{P}^2 = \mathcal{P}$  and  $\mathcal{A}^2 = \mathcal{A}$ . Each map is a sum of positive corner maps, is unital, and preserves the trace. An operator is fixed by  $\mathcal{P}$  exactly when its off-diagonal partition blocks vanish, which is equivalent to commuting with every  $P_i$ . An operator is fixed by  $\mathcal{A}$  exactly when it is a scalar multiple of  $P_i$  on each nonzero block. These range descriptions imply  $\mathcal{A}\mathcal{P} = \mathcal{A} = \mathcal{P}\mathcal{A}$ . Finally,  $\text{Tr}(\mathcal{A}(X)P_i) = \text{Tr}(X P_i)$ ; applying this identity to  $P_i X P_i$  and normalizing gives the conditioning statement.  $\square$

These are standard finite-matrix identities related to trace-preserving conditional expectations [22, 23]. The accompanying formal development contributes a checked arbitrary-partition interface, exact range and uniqueness statements, and adapters from projection events to the state-level Clauser–Horne–Shimony–Holt bound [6]. No priority claim is made for Born probability, Lüders conditioning, pinching, or Tsirelson’s inequality.

*Boundary 4.3.* This is a finite operational record interface. It does not reconstruct an arbitrary quantum state space or an interacting continuum quantum field theory. Its role is narrower: the completed consensus record carries the finite probability, conditioning, expectation, and correlation operations consumed by later branches.

## 5 Overlap defects and edge-center entropy

Overlap agreement has a second layer beyond equality of scalar records. Transport maps can compose only up to a central multiplier. The resulting defect is a finite gluing invariant; removing triangle defects and obtaining endpoint-only transport are distinct operations.

**Theorem 5.1** (Central defect strictification and residual holonomy). *Let  $N_\Sigma$  be the nerve of a finite overlap cover. Suppose  $U_{ji} = U_{ij}^{-1}$  and*

$$U_{ij}U_{jk} = z_{ijk}U_{ik}, \quad z_{ijk} \in Z_\Sigma,$$

where the identified coefficient group  $Z_\Sigma$  is abelian and overlap transport acts trivially on it. Then:

(i)  $z$  is a Čech 2-cocycle,

$$z_{jkl}z_{ikl}^{-1}z_{ijl}z_{ijk}^{-1} = 1;$$

(ii) its class  $[z] \in \check{H}^2(N_\Sigma, Z_\Sigma)$  is invariant under changes of local frame;

(iii) a central edge rephasing removes every triangle multiplier exactly when  $[z] = 0$ ;

(iv) after such a strictification, endpoint-only transport on a charge block  $\mathcal{C}_\alpha$  holds exactly when the represented holonomy of every closed loop is the identity on  $\mathcal{C}_\alpha$ .

*Proof.* Associativity compares  $(U_{ij}U_{jk})U_{kl}$  with  $U_{ij}(U_{jk}U_{kl})$  and gives the cocycle identity. A local frame change conjugates the multiplier, hence leaves a central element unchanged. An edge rephasing by a 1-cochain changes  $z$  by its Čech coboundary, proving the third statement. Once the edge maps form a strict 1-cocycle, the discrepancy between two paths with common endpoints is the holonomy of the closed loop obtained by composing one path with the inverse of the other.  $\square$

The theorem gives a precise finite meaning to gluing curvature. A vanishing triangle class does not imply trivial transport around noncontractible loops. If the center varies as a local coefficient system, the cocycle equation must use the transported, twisted Čech differential; that case lies outside the untwisted statement above.

### 5.1 Finite edge-center decomposition

Let a regulated collar be cut into left and right halves with common interface  $\Sigma$ . Suppose finite-dimensional Hilbert spaces  $\tilde{H}_L, \tilde{H}_R$  carry diagonal unitary actions of a finite compact group  $G_\Sigma$ . Write

$$\tilde{H}_L = \bigoplus_{\alpha} V_{\alpha} \otimes H_{L,\alpha}, \quad \tilde{H}_R = \bigoplus_{\beta} V_{\beta}^* \otimes H_{R,\beta}.$$

Here the  $V_{\alpha}$  are pairwise inequivalent irreducible unitary representations of  $G_\Sigma$  and  $d_{\alpha} = \dim V_{\alpha}$ .

**Theorem 5.2** (Invariant collar decomposition and one-sided edge entropy). *The invariant collar space and its sector-preserving algebra have the exact forms*

$$H_C = (\tilde{H}_L \otimes \tilde{H}_R)^{G_\Sigma} \cong \bigoplus_{\alpha} H_{L,\alpha} \otimes H_{R,\alpha},$$

$$\mathcal{A}_C = \bigoplus_{\alpha} \mathcal{B}(H_{L,\alpha}) \otimes \mathcal{B}(H_{R,\alpha}), \quad Z(\mathcal{A}_C) = \bigoplus_{\alpha} \mathbb{C} P_{\alpha}^C.$$

Let  $\Omega_\alpha \in V_\alpha \otimes V_\alpha^*$  be the normalized invariant vector corresponding to  $d_\alpha^{-1/2} I_{V_\alpha}$ . A sector-preserving collar state has the form

$$\rho_C = \bigoplus_\alpha p_\alpha \rho_\alpha, \quad \rho_\alpha \in \mathcal{B}(H_{L,\alpha} \otimes H_{R,\alpha}),$$

where  $p_\alpha \geq 0$ ,  $\sum_\alpha p_\alpha = 1$ , and each block with  $p_\alpha > 0$  satisfies  $\rho_\alpha \geq 0$  and  $\text{Tr} \rho_\alpha = 1$ . Embed it in  $\tilde{H}_L \otimes \tilde{H}_R$  through  $\Omega_\alpha$  and trace out  $\tilde{H}_R$ . If  $\rho_{L,\alpha} = \text{Tr}_{H_{R,\alpha}} \rho_\alpha$ , the resulting left state is

$$\rho_L = \bigoplus_\alpha p_\alpha \frac{I_{V_\alpha}}{d_\alpha} \otimes \rho_{L,\alpha},$$

and its entropy splits as

$$S(\rho_L) = H(p) + \sum_\alpha p_\alpha S(\rho_{L,\alpha}) + \sum_\alpha p_\alpha \log d_\alpha.$$

The edge term is the expectation in  $\rho_L$  of the sector observable

$$Z_L = \sum_\alpha (\log d_\alpha) P_\alpha^L,$$

where  $P_\alpha^L$  projects onto  $V_\alpha \otimes H_{L,\alpha} \subset \tilde{H}_L$ .

*Proof.* The tensor product of the two representation decompositions contains  $V_\alpha \otimes V_\beta^*$ . Schur's lemma gives a one-dimensional invariant space when  $\alpha = \beta$  and zero otherwise, yielding the invariant direct sum and the center of its sector-preserving algebra. The partial trace of  $|\Omega_\alpha\rangle\langle\Omega_\alpha|$  over  $V_\alpha^*$  is  $I_{V_\alpha}/d_\alpha$ . The displayed expression for  $\rho_L$  follows. Entropy of a block-diagonal state is the Shannon entropy of the block weights plus the mean block entropy. Tensor-product additivity contributes  $\log d_\alpha$  from each maximally mixed representation factor, and  $\text{Tr}(\rho_L Z_L) = \sum_\alpha p_\alpha \log d_\alpha$ .  $\square$

The theorem distinguishes the gauge-invariant collar state from its one-sided reduced state; the representation-dimension term belongs to the latter. Reading  $Z_L$  as an area observable requires a physical realization map. Exact quantum Markovity alone also does not identify an arbitrary state-dependent Markov decomposition with these preselected edge-center factors; alignment of the state with the collar decomposition is an additional premise. Related edge-mode and algebra-center decompositions occur in lattice gauge theory [43, 44].

## 6 From observer velocities to events and spacetime

### 6.1 Exact kinematics of the spherical support

The oriented conformal spherical support carries

$$\text{Conf}^+(S^2) \cong \text{PSL}(2, \mathbb{C}) \cong \text{SO}^+(3, 1), \quad H^3 \cong \frac{\text{SO}^+(3, 1)}{\text{SO}(3)}, \quad \dim H^3 = 3. \quad (8)$$

These are classical isomorphisms of conformal geometry, consumed as exact mathematics [32]. They identify the three-dimensional hyperbolic homogeneous space of future unit timelike directions. Interpreting those directions as physical observer velocities requires realized events, clocks, and local frames.

*Boundary 6.1.* Observer-velocity geometry and populated event geometry are distinct objects, and the program keeps them distinct. A populated four-dimensional event base requires record separation, local charts, an open-image condition, affine transition data, a quadratic cone, and causal reachability; these are required constructions, not corollaries of (8). The velocity space  $H^3$  and event localization are typed as separate constructions, and no inference passes from one to the other without a certificate.

**Theorem 6.2** (Sufficient gluing criterion for a record-germ event manifold). *Let  $D$  be the algebraic direct limit of a cofinal sequence of finite record-germ stages. Suppose the refinement maps make the common-refinement distance  $d : D \times D \rightarrow [0, \infty)$  a well-defined pseudometric, independent of the chosen stage representatives. Quotient  $D$  by zero distance and take its metric completion; call the result  $X$ . Then  $X$  is a separable, second-countable metric space. Suppose the same cofinal sequence supplies:*

- (E1) *population of every point of  $X$  by realized record germs;*
- (E2) *a covering family of bi-Lipschitz homeomorphisms  $\phi_i : U_i \rightarrow V_i$  from open subsets  $U_i \subset X$  onto open subsets  $V_i \subset \mathbb{R}^4$ ;*
- (E3)  *$C^{1,1}$  affine overlap maps satisfying the triple-overlap cocycle;*
- (E4)  *$C^{1,1}$  tetrads and inverse tetrads on those charts;*
- (E5) *compatibility on overlaps: the chart metrics induced by the tetrads are related by pullback under the transition maps;*
- (E6) *a nondegenerate quadratic event form of inertia  $(1,3)$ , with compatible orientation and time orientation on every overlap; and*
- (E7) *local semantic causal reachability agreeing with the selected cone.*

*Then the event base is a time-oriented Lorentzian four-manifold of regularity  $C^{1,1}$ . The future-unit timelike vectors, equivalently observer velocities modulo spatial rotations, form an  $H^3$  fiber over each event. Stable causality and global hyperbolicity are additional conditions.*

*Proof.* The algebraic direct limit of countably many finite stages is countable and dense in its metric completion, so  $X$  is separable. Every separable metric space is second-countable and Hausdorff. The covering bi-Lipschitz charts and their open images give the local Euclidean topology, while the affine cocycle gives a consistent  $C^{1,1}$  atlas. The tetrads define chartwise Lorentz metrics, and the pullback clause makes them one global  $C^{1,1}$  metric. Inertia, orientation, time orientation, and cone-compatible reachability supply the asserted local causal structure. The unit future timelike vectors of each Lorentzian tangent space form  $\text{SO}^+(3,1)/\text{SO}(3) \cong H^3$ .  $\square$

## 6.2 A measured finite Lorentzian signature

The simulation branch measures the event form. Repair dynamics on federated carriers emits semantic events with ancestry; a declared chart combines one ancestry-depth coordinate with three spectral coordinates; a quadratic event form is fitted on a declared training half of the event pairs and evaluated on the held-out half. The signature is therefore held out from the update rule and is classified as a certified finite computation. The cone margin is the minimum held-out signed score, with sign  $+1$  for ancestry-comparable pairs and  $-1$  for incomparable pairs. A negative margin means that at least one held-out pair lies on the wrong side of the fitted quadratic cone.



On the reported support-adjusted comparison path with (carrier count, observer count, support width) equal to (16,384, 128, 96), (65,536, 256, 96), and (262,144, 512, 384), the held-out event form has inertia (1, 3) at every rung, with cone margins

$$-5.6, \quad -3.2, \quad -1.4 \quad (9)$$

on rows that differ in support width and cross-edge density, so the shrinking magnitudes do not form a convergence sequence. At 262,144 carriers the same-size support-width-96 control has 312 cross-observer edges and inertia (2, 2), while the support-width-384 row has 1,062 edges and inertia (1, 3). In this finite comparison, changing the support and cross-read structure changes the fitted inertia. The comparison does not isolate a unique mechanism or establish a continuum limit [2].

*Boundary 6.3.* The chart allocates four coordinates by declaration; the instrument tests signature and full rank on that chart and does not derive the number of spacetime dimensions. The negative cone margins, one Euclidean local fit in the finite source domain below, closed neighborhoods in place of open charts, and the absence of a certified refinement limit prevent promotion of any finite object in this section to a continuum Lorentzian manifold. These conditions are recorded machine-readably in the receipts; the continuum attachment is an open realization map.

### 6.3 A finite source satisfying part of the local-domain contract

One deterministic capture at 16,384 carriers supplies: a finite causal complex of 2,304 events with exact acyclicity and a strict time function; a certified four-column chart whose global held-out form has inertia (1, 3); six closed observer-visibility neighborhoods with exact induced affine transitions, cocycles, orientation, and time orientation; an observer-visible seam complex of 8,662 carriers and 11,816 seams whose 38 triangles are all frustrated (no global sign assignment satisfies every seam) under the declared orientation-reversing transport, with lift-ambiguity rank 3,117 over  $\mathbb{F}_2$ ; typed scalar, chiral, and gauge sections whose sign-twisted local derivative passes exact integer adjoint, kinetic, covariance, gluing, refinement, and boundary checks; and a signed-graph rank theorem giving a zero twisted kernel, hence an exactly positive dimensionless spectral gap for the declared local operator. The capture is a certified finite computation; its rank and positivity statements are finite theorems. One provenance graph binds the construction, an isolated rerun reproduces the canonical receipt byte for byte, and the verifier rejects any receipt that promotes this finite domain to a continuum spacetime, a physical clock, or a physical mass scale [2].

## 7 Einstein reconstruction

**Theorem 7.1** (Null tomography and metric ambiguity). *Let  $V$  be four-dimensional with Lorentz form  $\eta$ . Suppose a symmetric form  $X$  satisfies  $X(k, k) = 0$  for every null vector  $k$ . Then  $X = \phi\eta$  for a scalar  $\phi$ . Null-null values therefore determine a symmetric tensor modulo the metric line. In an  $\eta$ -orthonormal basis with  $\eta = \text{diag}(-1, 1, 1, 1)$ , set  $s = \sqrt{3}/3$  and take the nine null vectors*

$$\begin{aligned} (1, \pm 1, 0, 0), \quad (1, 0, \pm 1, 0), \quad (1, 0, 0, \pm 1), \\ (1, s, s, s), \quad (1, s, s, -s), \quad (1, s, -s, s). \end{aligned}$$

*On the  $\eta$ -trace-free representative, use coordinates*

$$x = (X_{00}, X_{01}, X_{02}, X_{03}, X_{11}, X_{12}, X_{13}, X_{22}, X_{23}), \quad X_{33} = X_{00} - X_{11} - X_{22}.$$

*The corresponding nine-charge design reconstructs the quotient*

$$\text{Sym}^2(V^*)/\mathbb{R}\eta.$$

Its determinant is  $8192/27$ . With the supremum norms on coordinates and charges, its exact decoder obeys

$$\|x\|_\infty \leq (2 + \sqrt{3})\|q\|_\infty.$$

*Proof.* Choose coordinates with  $\eta = \text{diag}(-1, 1, 1, 1)$  and write every future null ray as  $(1, n)$  with  $|n| = 1$ . The identity

$$X_{00} + 2X_{0i}n_i + X_{ij}n_in_j = 0$$

holds on the unit sphere. Its odd part gives  $X_{0i} = 0$ . The degree-two spherical-harmonic part gives  $X_{ij} = \phi\delta_{ij}$ , and the constant part gives  $X_{00} = -\phi$ . Thus  $X = \phi\eta$ . The quotient therefore has dimension  $10 - 1 = 9$ . Substitution of the nine displayed algebraic directions into the stated trace-free coordinate basis gives determinant  $8192/27$ . The explicit decoder has maximum absolute row sum  $2 + \sqrt{3}$ , which gives the supremum-norm bound. These two constants are exact analytic calculations. The accompanying Lean development proves that the nine directions are null and formalizes the design map, an explicit left-inverse decoder, injectivity, and the metric-line ambiguity [1, 10].  $\square$

The four-dimensional small-ball coefficients used in entanglement-equilibrium arguments [39] can be isolated from their physical interpretation.

With  $G_N > 0$  and  $\ell > 0$ , scalars  $\delta S_{\text{bulk}}, \delta A, t, f$  satisfying

$$\delta S_{\text{bulk}} = \frac{8\pi^2\ell^4}{15}t, \quad \delta A = -\frac{4\pi\ell^4}{15}f, \quad \delta S_{\text{bulk}} + \frac{\delta A}{4G_N} = 0 \quad (10)$$

obey  $f = 8\pi G_N t$  exactly: substitution cancels the common nonzero prefactor. The identity isolates the numerical coefficients from their physical reading and is checked by machine.

The tensor completion begins at an explicitly realized null-balance relation. This is the point at which the physical bridge enters the mathematical implication.

**Theorem 7.2** (Tensor completion of a realized null balance). *Let  $(M, g)$  be a connected, time-oriented,  $C^3$  Lorentzian four-manifold with Einstein tensor  $G_{ab}^{(g)}$ . Let  $T_{ab}$  be a symmetric  $C^1$  tensor satisfying the independent Ward premise  $\nabla^a T_{ab} = 0$ . Suppose a common realized tower supplies a constant  $\kappa > 0$  such that, at every point and for every null vector  $k$ ,*

$$G_{ab}^{(g)}k^ak^b = \kappa T_{ab}k^ak^b, \quad (11)$$

and suppose  $G_N > 0$  with  $\kappa = 8\pi G_N$ . Then there is a constant  $\Lambda$  such that

$$G_{ab}^{(g)} + \Lambda g_{ab} = 8\pi G_N T_{ab} \quad (12)$$

on  $M$ . If a reference event  $p_0$  and scalar  $\Lambda_0$  satisfy

$$(G_{ab}^{(g)} - 8\pi G_N T_{ab})|_{p_0} = -\Lambda_0 g_{ab}|_{p_0},$$

then  $\Lambda = \Lambda_0$ .

*Proof.* At each point, Theorem 7.1 applied to  $G^{(g)} - \kappa T$  gives a scalar  $\phi$  with  $G_{ab}^{(g)} - \kappa T_{ab} = \phi g_{ab}$ . The contracted Bianchi identity  $\nabla^a G_{ab}^{(g)} = 0$ , the Ward premise, and metric compatibility give  $\nabla_b \phi = 0$ . Connectedness makes  $\phi$  constant. Set  $\Lambda = -\phi$  and use  $\kappa = 8\pi G_N$ . Evaluation at  $p_0$  gives  $\Lambda = \Lambda_0$  under the final calibration premise.  $\square$

For OPH to supply the null-balance premise (11), one common refinement tower must realize all of the following:

- (G1) the event manifold of Theorem 6.2, with sufficient regularity for its Levi–Civita connection and contracted Bianchi identity;
- (G2) geometrically normalized cap modular flow and half-sided modular inclusions whose directional charges define one symmetric tensor  $T_{ab}$ ;
- (G3) the Ward identity for that tensor, independently of its reconstruction from directional charges;
- (G4) the edge-center split of Theorem 5.2, the physical normalization  $\delta\langle Z_L \rangle = \delta A/(4G_N)$ , the modular first law, and generalized-entropy stationarity;
- (G5) one family with  $\ell_r > 0$ ,  $\ell_r \rightarrow 0$ , all named remainders  $o(\ell_r^4)$ , the continuum diamond-kernel and fixed-volume area formulas used in the small-ball identity (10), and enough local observer directions to establish (11); and
- (G6) universal coupling of the stress and entropy branches, a vacuum reference, and a physical scale calibration establishing  $\kappa = 8\pi G_N$ .

The exact finite layer contains null tomography, the edge-entropy split, and the coefficient identity. The joint physical tower, its continuum formulas, Ward identity, area normalization, clock and stress identifications, and scale readings are the open realization maps of the composition. The machine-checked composition proves the tensor step from an explicit null-balance premise and separately checks the small-ball arithmetic; it does not construct the bridge between them. The finite signature measurement in Section 6.2 tests one geometric ingredient and does not replace these premises.

## 8 Two gauge reconstructions and a finite matter image

### 8.1 Sector reconstruction

The overlap and edge-center branch gives a structural gauge result before a particular compact group is identified.

**Theorem 8.1** (Bosonic sector reconstruction). *Suppose a cofinal refinement tail carries zero-obstruction, trivial-holonomy bosonic edge sectors and compatible fully faithful pullback functors. Suppose their closure  $\text{Sect}_\infty$  is an essentially small, additive, idempotent-complete, semisimple rigid symmetric  $C^*$ -tensor category with simple unit and finite-dimensional Hom spaces. Suppose it has a faithful, unitary, strong symmetric monoidal,  $*$ -preserving fiber functor*

$$\mathcal{F} : \text{Sect}_\infty \longrightarrow \text{Hilb}_{\text{fd}}.$$

*Then*

$$G_{\text{Tan}} = \text{Aut}_{\otimes}^{u,*}(\mathcal{F})$$

*is compact in the topology of pointwise operator convergence and  $\text{Sect}_\infty \simeq \text{Rep}(G_{\text{Tan}})$  as symmetric  $C^*$ -tensor categories. The reconstructed group is unique up to isomorphism for the specified category and fiber functor.*

*Proof.* Here  $\text{Aut}_{\otimes}^{u,*}(\mathcal{F})$  denotes the unitary monoidal natural  $*$ -automorphisms of  $\mathcal{F}$ . For every object  $X$ , such an automorphism has a component in the compact group  $U(\mathcal{F}(X))$ . Naturality, tensor compatibility, symmetry, and  $*$ -compatibility are closed equations, so this automorphism group is a closed subgroup of the product  $\prod_X U(\mathcal{F}(X))$  over a small skeleton and is compact in the subspace topology. The Doplicher–Roberts/Tannaka reconstruction theorem then gives the equivalence [46, 47]. A second compact group compatible with the same fiber functor is identified with the same automorphism group.  $\square$

Theorem 8.1 classifies the group encoded by the realized sector data. A trivial sector category reconstructs a trivial group. It does not select the Standard Model group without a suitable sector witness, and it does not identify its group with the finite response group below.

## 8.2 The twelve-port response algebra

Let  $A$  be the icosahedral adjacency operator and  $J$  the antipodal involution on the twelve ports. Exact incidence gives

$$10J = A^3 - 4A^2 - 5A + 10I. \quad (13)$$

Put  $\varphi = (1 + \sqrt{5})/2$ . Choose from each antipodal axis one of the certificate’s exact coordinate vectors  $u_i \in \mathbb{Q}(\sqrt{5})^3$ , each with  $\|u_i\|^2 = 2 + \varphi$ , and put  $U = [u_1 \cdots u_6]$ . For a real port field  $f$ , write

$$b_i = \frac{f_i + f_{Ji}}{2}, \quad d_i = \frac{f_i - f_{Ji}}{2}, \quad c = \frac{1}{6} \sum_i b_i, \quad b^0 = b - c\mathbf{1}.$$

The even coordinates split as  $\mathbf{1} \oplus \mathbf{5}$ . The odd coordinates split as the rank-three frame channel detected by  $U$  and its Galois-conjugate companion detected by  $\sigma(U)$ , where  $\sigma(\sqrt{5}) = -\sqrt{5}$ , giving the source-module decomposition

$$P_{12} \cong_{A_5} \mathbf{1} \oplus \mathbf{5} \oplus \mathbf{3} \oplus \mathbf{3}'.$$

**Theorem 8.2** (A1–A2 gauge Lie-type forcing). *Let  $D : P_{12} \rightarrow \mathfrak{u}(H)$  be the complete reversible response derivative supplied by A1, and put  $\mathfrak{g} = D(P_{12})$ . Suppose the endogenous transport clause of A2 covers every proper carrier action  $a \in A_5$  and gives projective implementers satisfying*

$$\text{Ad}_{[U_a]} D(v) = D(a \cdot v), \quad [U_a] = [g_a c_a],$$

where  $g_a \in G_D^0$  and  $c_a$  centralizes  $\mathfrak{g}$  pointwise. Then

$$\mathfrak{g} \cong \mathfrak{u}(1) \oplus \mathfrak{su}(3) \oplus \mathfrak{su}(2).$$

*The center is the unique trivial line. Up to exchanging  $\mathbf{3}$  and  $\mathbf{3}'$  by the outer automorphism of  $A_5$ , the  $\mathfrak{su}(2)$  ideal carries  $\mathbf{3}$  and the  $\mathfrak{su}(3)$  ideal carries  $\mathbf{3}' \oplus \mathbf{5}$ .*

*Proof.* Injectivity of  $D$  gives  $\dim \mathfrak{g} = 12$ . Its commutator-closed image is a finite-dimensional Lie subalgebra of  $\mathfrak{u}(H)$ , and the positive trace pairing makes it compact. For every  $a \in A_5$ , the centralizer factor  $c_a$  acts trivially on  $\mathfrak{g}$ , so  $\text{Ad}_{[U_a]}|_{\mathfrak{g}} = \text{Ad}_{g_a}|_{\mathfrak{g}}$  is inner. The chosen projective lifts need not form a homomorphic section. Covariance through the injective map  $D$  nevertheless gives the homomorphism

$$a \longmapsto (D(v) \mapsto D(a \cdot v)) \quad \text{from } A_5 \text{ to } \text{Int}(\mathfrak{g}).$$

The transitive proper-carrier action fixes only the uniform port line, hence  $\dim \mathfrak{g}^{A_5} = 1$ .

Compactness gives the reductive decomposition  $\mathfrak{g} = \mathfrak{z} \oplus [\mathfrak{g}, \mathfrak{g}]$ . Inner automorphisms fix the center pointwise, so  $\dim \mathfrak{z} \leq 1$ .

If  $\mathfrak{z} = 0$ , the low-dimensional classification of compact simple Lie algebras forces a twelve-dimensional compact semisimple algebra to be  $\mathfrak{su}(2)^4$  [45]. Inner automorphisms preserve each simple ideal. On each three-dimensional ideal, the restriction of the  $A_5$  action is either trivial or faithful because  $A_5$  is simple. Its fixed-space dimension is therefore three or zero: a faithful  $A_5$  subgroup of  $SO(3)$  cannot fix an axis, since a finite rotation group fixing an axis is cyclic. The fixed-space dimension of  $\mathfrak{g}$  would be a multiple of three, contradicting the one-dimensional fixed line. Hence  $\dim \mathfrak{z} = 1$ .

The semisimple part has dimension eleven. The same low-dimensional classification leaves only  $11 = 8 + 3$ , realized by  $\mathfrak{su}(3) \oplus \mathfrak{su}(2)$ . Adding the center proves the Lie-algebra claim. The center must be the unique fixed line. The three-dimensional simple ideal cannot carry the trivial action, since that would add three fixed directions; hence it carries  $\mathbf{3}$  or  $\mathbf{3}'$ , and the eight-dimensional ideal carries the complementary triplet together with  $\mathbf{5}$ .  $\square$

*Boundary 8.3.* No ambient compact group or candidate Lie algebra occurs among the premises of Theorem 8.2. A1 supplies the faithful compact commutator-closed response algebra, and A2 supplies its inner proper-carrier action. The theorem fixes the abstract local Lie-algebra type. The source receipts derive the carrier module, the inverse-port response, and its relative band signs. The source contains no reconstruction of  $D$  from ordered reversible histories or same-current holonomy; the identification consumes both as premises. The physical matrix current, couplings, matter action, global quotient, and laboratory attachment are the open realization maps of this branch. The port count twelve equals the dimension of the forced algebra, so A1 fixes the dimension by declaration; the content of the theorem is the exclusion, through compactness and the one-dimensional fixed space, of the only other twelve-dimensional compact type  $\mathfrak{su}(2)^4$ , and the forcing of the unique center. No uniqueness claim over carriers is made, and under the closure hypothesis of Section 1.3 the carrier is read as part of the reconstructed solution. This interpretation does not enter the theorem, whose declared premises stand independently.

Let

$$\Phi_0(b^0) = \sum_{i=1}^6 b_i^0 u_i u_i^\top, \quad \hat{x} y = x \times y.$$

The conditional matrix witness uses the space

$$H = \mathbb{C}_E^3 \oplus \mathbb{C}_W^3$$

and four nonzero rational coefficients  $\lambda_1, \lambda_5, \lambda_3, \lambda_{3'}$ , one on each source band. These are explicit branch premises. The two three-dimensional response blocks match the two rank-three odd channels; they are not asserted to be a derived physical Hilbert space. Write  $v_p$  for the signed coordinate vector at port  $p$ . For  $g \in A_5$ , let  $R_g \in SO(3)$  be the exact rotation satisfying  $R_g v_p = v_{g(p)}$  on the oriented port frame, and define

$$\Pi(g) = \text{diag}(R_g, \sigma(R_g)).$$

The target  $A_5$  action on  $\mathfrak{u}(H)$  is conjugation by  $\Pi(g)$ .

**Theorem 8.4** (Conditional four-band finite port-response algebra). *On the declared twelve-port carrier and supplied response representation, define*

$$\begin{aligned} K(f) &= \text{diag}(\lambda_3 \widehat{U} d + i\{\lambda_1 c I_3 + \lambda_5 \Phi_0(b^0)\}, \\ &\quad \lambda_{3'} \widehat{\sigma(U)} d) \in \mathfrak{u}(H). \end{aligned} \tag{14}$$

Then  $K$  is injective, its image is commutator-closed, and

$$K(g \cdot f) = \Pi(g)K(f)\Pi(g)^*.$$

The induced  $A_5$  action on  $\text{im } K$  is inner. Let  $K^{-1} : \text{im } K \rightarrow P_{12}$  denote the inverse of  $K$  onto its image. With the bracket pulled back from that image,

$$[x, y]_K = K^{-1}([K(x), K(y)]_{\text{mat}}),$$

one has

$$(P_{12}, [\cdot, \cdot]_K) \cong \mathfrak{u}(3) \oplus \mathfrak{so}(3) \cong \mathfrak{u}(1) \oplus \mathfrak{su}(3) \oplus \mathfrak{su}(2). \quad (15)$$

Its center is the uniform one-dimensional port line, its derived algebra has dimension eleven, and the five-dimensional  $A_5$  band is noncentral. The map has a positive-definite invariant Hilbert–Schmidt pullback.

*Proof.* Equation (13) is obtained by evaluating both sides on the four distinct adjacency eigenspaces  $\mathbf{1}, \mathbf{3}, \mathbf{3}', \mathbf{5}$ . The source protocol solves the common farthest-shell filter on each port and gives the signed response  $-J$ .

The constant coordinate maps to  $i\mathbb{R}I_3$ . The six rank-one axis matrices  $u_i u_i^\top$  form a basis of the symmetric  $3 \times 3$  matrices and satisfy

$$\sum_i u_i u_i^\top = (5 + \sqrt{5})I_3.$$

Their sum-zero coefficients therefore map isomorphically to the traceless symmetric matrices under  $\Phi_0$ . The first two even terms span  $i\text{Sym}_3(\mathbb{R})$ . The map  $U$  is an isomorphism on the  $\mathbf{3}$  odd band and annihilates the  $\mathbf{3}'$  band;  $\sigma(U)$  has the complementary property. Their hat maps span one copy of  $\mathfrak{so}(3)$  in each response block. Because every  $\lambda_\bullet$  is nonzero, these four images are independent and  $K$  has rank  $1 + 5 + 3 + 3 = 12$ .

In the first block,  $i\text{Sym}_3(\mathbb{R}) \oplus \mathfrak{so}(3) = \mathfrak{u}(3)$  as a real Lie algebra, while the second block is  $\mathfrak{so}(3)$ . Thus the image is exactly  $\mathfrak{u}(3) \oplus \mathfrak{so}(3)$  and is closed under commutators. Pullback of the matrix commutator satisfies bilinearity, antisymmetry, and Jacobi and gives (15). The derived algebra is  $\mathfrak{su}(3) \oplus \mathfrak{so}(3)$ , of dimension  $8 + 3 = 11$ ; the remaining line is the center of  $\mathfrak{u}(3)$ . Direct evaluation identifies it with the uniform port vector and exhibits a nonzero commutator involving the five-dimensional band. Equivariance and positive definiteness of the pairing  $-\text{Re tr}(K(f)K(f'))$  are exact calculations in  $\mathbb{Q}(\sqrt{5})$ , verified over the full sixty-element domain by the exact-arithmetic certificate with its manifest, receipt, and negative controls [1]; by the exhaustive-computation rule of Table 1 these calculations are part of the proof. The same certificate verifies, for all sixty elements, that  $\Pi(g)$  is the exponential of an element of  $\text{im } K$ ; hence its conjugation action is inner. The finite refinement and persistence diagrams recorded in the certificate are checked there as separate commuting-square computations.  $\square$

*Boundary 8.5.* The  $A_5$ -module decomposition alone does not determine a Lie bracket: the same module also carries the zero bracket, whose trivial response-generated group fails the A2 endogeneity clause. The independent inputs to Theorem 8.4 are the carrier incidence and orientation, the supplied response representation, and four nonzero band coefficients. The block architecture and coefficients are declared. No measured particle, mass, or coupling value enters the exact verification. The theorem constructs the pulled-back bracket for this response map; it does not select  $K$  from source histories or prove a uniqueness claim over all  $A_5$ -equivariant brackets or finite carriers. The abstract Lie type is forced independently by Theorem 8.2.

### 8.3 Global form and anomaly-free exterior module

**Theorem 8.6** (Finite matter image and exhaustive exterior-menu selection). *Let*

$$V = \mathbb{C}_{-1/3}^3 \oplus \mathbb{C}_{1/2}^2$$

*be an additional, explicitly supplied trace-balanced matter representation of the Lie algebra forced in Theorem 8.2. This  $3 \oplus 2$  matter space is distinct from the supplied response space  $H = \mathbb{C}_E^3 \oplus \mathbb{C}_W^3$  and is not derived from it. Let*

$$\tilde{G} = \text{SU}(3) \times \text{SU}(2) \times \text{U}(1)$$

*act on  $V$  by*

$$(g_3, g_2, z) \cdot (v_3, v_2) = (z^{-2}g_3v_3, z^3g_2v_2).$$

*The differential of this action gives the displayed hypercharges. Its exterior module has the exact branching*

$$\Lambda^2 V \oplus \Lambda^4 V = Q \oplus u^c \oplus e^c \oplus d^c \oplus L \quad (16)$$

*with*

| <i>component</i> | $\text{SU}(3) \times \text{SU}(2)$ <i>type</i> | <i>Y</i> |
|------------------|------------------------------------------------|----------|
| $Q$              | $(3, 2)$                                       | $1/6$    |
| $u^c$            | $(\bar{3}, 1)$                                 | $-2/3$   |
| $e^c$            | $(1, 1)$                                       | $1$      |
| $d^c$            | $(\bar{3}, 1)$                                 | $1/3$    |
| $L$              | $(1, 2)$                                       | $-1/2$   |

*This is the familiar one-generation  $\mathbf{10} \oplus \bar{\mathbf{5}}$  branching associated with  $\text{SU}(5)$  unification [48]. It has total complex dimension fifteen. All perturbative gauge and mixed gravitational anomalies vanish, and the number of weak doublets, counted with color multiplicity, is four. The common kernel of the action of  $\tilde{G}$  on  $V$ , and hence on  $\Lambda^2 V \oplus \Lambda^4 V$ , is  $\mathbb{Z}_6$ . Hence the maximal faithful matter image is*

$$S(U(3) \times U(2)) \cong \frac{\text{SU}(3) \times \text{SU}(2) \times \text{U}(1)}{\mathbb{Z}_6}. \quad (17)$$

*Define*

$$G_{\text{packet}} := S(U(3) \times U(2)),$$

*the maximal faithful image of the displayed source-model module. This quotient is familiar as a possible global form of the Standard Model gauge group [50]. The scan menu consists of the ten nontrivial irreducible summands of  $\Lambda^\bullet V$  after the invariant vacuum  $\Lambda^0 V$  and invariant top line  $\Lambda^5 V$  have been excluded by declaration. Among all  $2^{10} = 1,024$  subsets of this menu, imposing nonemptiness, chirality, and vanishing of the mixed gravitational– $\text{U}(1)$ ,  $\text{SU}(3)^2\text{U}(1)$ ,  $\text{SU}(2)^2\text{U}(1)$ , and  $\text{U}(1)^3$  traces leaves exactly two rank-fifteen masks. They are exchanged by charge conjugation, and the exterior-parity grading is computed as an output.*

*Proof.* The exterior-sum identity

$$\Lambda^k(A \oplus B) = \bigoplus_{p+q=k} \Lambda^p A \otimes \Lambda^q B$$

gives the displayed five components and charges. The anomaly-cancellation conditions are standard [49]. With the conventional quadratic indices  $T(3) = T(2) = 1/2$ , the pure color anomaly is

$$\text{SU}(3)^3 : \quad 2 - 1 - 1 = 0,$$

where the two color fundamentals occur in the weak doublet and the two antifundamentals are  $u^c$  and  $d^c$ . The local  $SU(2)^3$  anomaly vanishes identically because the doublet is pseudoreal. The mixed anomalies are

$$SU(3)^2U(1) : \quad \frac{1}{6} - \frac{1}{3} + \frac{1}{6} = 0, \quad SU(2)^2U(1) : \quad \frac{1}{4} - \frac{1}{4} = 0.$$

The abelian cubic and mixed gravitational sums are

$$6\left(\frac{1}{6}\right)^3 + 3\left(-\frac{2}{3}\right)^3 + 1 + 3\left(\frac{1}{3}\right)^3 + 2\left(-\frac{1}{2}\right)^3 = 0,$$

$$6\left(\frac{1}{6}\right) + 3\left(-\frac{2}{3}\right) + 1 + 3\left(\frac{1}{3}\right) + 2\left(-\frac{1}{2}\right) = 0.$$

There are three color copies of the  $Q$  doublet and one  $L$  doublet, so the Witten parity is even [51].

For the displayed cover action, the element

$$\left(e^{2\pi i/3}I_3, -I_2, e^{i\pi/3}\right)$$

acts trivially on every row of the table and generates a cyclic group of order six. Direct center-action enumeration proves that no larger common kernel exists; the independent integer presentation has Smith invariants  $(1, 1, 1, 1, 1, 6)$ . The cover action has determinant one on  $V$ , so  $\Lambda^4 V \cong V^*$  equivariantly. Its kernel on the displayed exterior module therefore equals its kernel on  $V$ . This gives (17). Finally, the finite scan evaluates nonemptiness, chirality, and the four displayed scan traces on all 1,024 masks in the declared ten-summand menu. Its only survivors are the two exterior-parity masks; the explicit conjugation permutation exchanges them. Parity is recorded after selection and is not a selection predicate.  $\square$

*Boundary 8.7.* The finite response group  $G_{\text{packet}}$  and the sector-reconstructed group  $G_{\text{Tan}}$  arise from different data. Their equality is a separate commuting-square premise. Laboratory current attachment, physical matter identification, exclusion of extra light sectors, scalar attachment, family multiplicity, Yukawa couplings, mixing, masses, and a continuum quantum-field realization are the open realization maps of the matter branch. The theorem classifies a finite source-model representation and its maximal faithful image. The exclusion of  $\Lambda^0 V$ ,  $\Lambda^5 V$ , direct sums, vectorlike additions, and neutral singlets is not derived by this scan.

## 8.4 Massless carriers, conditionally

Three conditional masslessness statements occur on distinct premise sets. They are consistency consequences of their declared branches and supply no independent discriminator from the Standard Model or General Relativity.

*The photon.* A finite one-loop calculation imports the Standard Model electroweak action and its symmetry-restored vacuum relation. The calculation certifies a transverse photon self-energy and a null direction of the neutral mass matrix, hence no hard photon mass term on that fixture. This validates the calculation harness. An OPH derivation additionally requires a source-generated physical action, laboratory-current attachment, and photon-pole map.

*The gluon.* The color factor in (15) lies in the unbroken subalgebra of the conditional global form. Under the separately declared scalar completion, no scalar direction carries color and the finite action contains no hard color-vector mass term. This statement does not construct the physical gluon spectrum. Confinement and hadron physics require strong-coupling dynamics outside the finite calculation.



*The graviton.* The Einstein branch of Theorem 7.2 carries a massless spin-two carrier exactly as far as the composition carries the field equation: linearizing the composed relation about a vacuum yields the standard massless graviton kinematics, as a reconstruction implication conditional on the same premise vector and inheriting its premises. No independent graviton claim is made beyond the composition.

The photon and gluon statements concern finite actions with declared inputs. The graviton statement inherits every premise of the Einstein composition.

## 9 Exact quantitative closures and physical tests

The finite architecture supports a small set of quantitative surfaces sharp enough to be tested numerically. Each surface below is stated with its type from Table 1: the theorems are finite theorems, the numerical enclosures are certified finite computations, and every empirical surface is typed as a diagnostic, a registered conditional test, or a frozen prospective branch prediction. The closure hypothesis of Section 1.3 contributes its quantitative candidates here, in Section 9.1, under the same typing.

### 9.1 Closure candidates: the screen grain and the capacity coordinate

In the displayed proposals,  $A_T(\cdot)$  is the inverse electromagnetic coupling the declared Thomson-limit source map returns at a trial screen grain,  $\alpha_U(\cdot)$  the coupling the same declared map returns at its unification point,  $\mathfrak{U}_N$  the regulated universe-level observer system at capacity budget  $N$ , and  $M_0$  the correctable-code capacity of its terminal public record; the grain scalar of these equations is distinct from the port set and the event projectors used elsewhere in this paper. Certified roots of both displayed maps are compared against the 2022 CODATA inverse coupling [56]. The symbol  $\varphi = (1 + \sqrt{5})/2$  is the golden ratio of Section 8.

**The closure principle.** The axioms describe the observer screen. The global closure principle states that the simulating and the simulated description are one system. Every quantity that has both a construction-side reading and a readback-side reading must take the same value once a typed bridge proves that the two readings denote one invariant. The equality is forced by self-identity. Constructing that bridge and the return map remains part of the physics. Existence, uniqueness, and stability are separate determinacy tests on the resulting equation. The name given to such an equation carries no mathematical weight: after the typed identification, unequal readings would describe two systems rather than the single self-referential universe.

The local screen-grain proposal

$$P = \varphi + \sqrt{\pi}/A_T(P),$$

with  $A_T(P)$  the inverse coupling returned by the declared source map, has one interval-certified fixed point on its physical domain; its laboratory attachment awaits a same-scheme hadronic spectral transport. The displayed map's certified fixed point sits at  $P = 1.6309720959$  and returns  $A_T = 136.9948352$ ; the certified gauge-width variant's fixed point sits at  $P = 1.6309682414$  and returns 137.0356601. Through the same outer equation the measured 2022 CODATA coupling  $\alpha^{-1} = 137.035999177(21)$  reads back the grain value  $P = 1.6309682094$ . The comparisons stand at  $3.0 \times 10^{-4}$  relative for the displayed map and at  $2.5 \times 10^{-6}$  for the gauge-width variant, as diagnostics of the declared source maps.

The direct global proposal is

$$N = \log M_0(\mathfrak{U}_N).$$

The complete declared finite source class does not select a unique cosmic value, so any successful completion needs an additional source law. A separate common-load branch starts from

$$N_0 = \pi \exp\left(\frac{6\pi}{P \alpha_U(P)}\right).$$

On the finite collar branch, the declared total reserve expectation  $P/4$  and six-class equidistribution give presence probability  $P/24$  for each declared class. If one class is physically selected as the blocked event, its scalar-weighted presence receipt is discharged, and that normalized collar-survival factor is proved to act on the global capacity, the corresponding candidate is

$$N_{\text{pres}} = N_0 \left(1 - \frac{P}{24}\right), \quad \ln \frac{N_{\text{pres}}}{\pi} = \frac{6\pi}{P \alpha_U(P)} + \ln \left(1 - \frac{P}{24}\right).$$

The exponential alternative  $N_{\text{Pois}} = N_0 e^{-P/24}$  requires a separate mean-count or continuum carrier. Neither global attachment is derived. The inherited screen/electroweak bridge premises, physical common-load bridge, physical  $\mathbb{Z}_6$  seam action, one-class reserve selection, scalar-weighted receipt, reserve-to-capacity map, and horizon-record identification are open. On the source-forward numerical branch,

$$N_0 = 3.5321 \times 10^{122},$$

so the finite-presence candidate is  $3.2921 \times 10^{122}$  and the exponential candidate is  $3.3001 \times 10^{122}$ . The full weighted Planck base- $\Lambda$ CDM chain gives the comparison coordinate  $3.3129 \times 10^{122}$ . The respective residuals are  $-0.63$  percent and  $-0.39$  percent. Both comparisons were exposed before the branch choice and carry no predictive weight.

## 9.2 The positive-chamber Koide identity

The icosahedral carrier's face structure supplies order-three cyclic response operators. The following exact identity governs every Hermitian three-cycle response [13].

**Theorem 9.1** (Positive-chamber Koide identity). *Let  $R^3 = I$  generate a cyclic action and let*

$$C = aI + bR + \bar{b}R^2, \quad a > 0, \quad b = \rho e^{i\delta} \in \mathbb{C},$$

*be the associated Hermitian circulant, with eigenvalues*

$$\lambda_k = a + 2\rho \cos\left(\delta + \frac{2\pi k}{3}\right), \quad k = 0, 1, 2.$$

*Suppose all three eigenvalues are nonnegative and, for one scale  $s > 0$ ,  $\sqrt{m_k} = \sqrt{s} \lambda_k$ . Then*

$$Q := \frac{\sum_k m_k}{(\sum_k \sqrt{m_k})^2} = \frac{1}{3} + \frac{2}{3} \left(\frac{\rho}{a}\right)^2, \quad \text{so} \quad Q = \frac{2}{3} \iff \frac{\rho}{a} = \frac{1}{\sqrt{2}}. \quad (18)$$

*Proof.* The three cosines at angles separated by  $2\pi/3$  sum to zero, and their squares sum to  $3/2$ . Hence  $\sum_k \lambda_k = 3a$  and  $\sum_k \lambda_k^2 = 3a^2 + 6\rho^2$ . With  $m_k = s\lambda_k^2$  and  $\sqrt{m_k} = \sqrt{s} \lambda_k$ , the ratio is

$$Q = \frac{s(3a^2 + 6\rho^2)}{s(3a)^2} = \frac{1}{3} + \frac{2}{3} \left(\frac{\rho}{a}\right)^2.$$

The equivalence follows by solving  $Q = 2/3$  for  $\rho/a$  on  $a > 0$ ,  $\rho \geq 0$ . □

The phase  $\delta$  cancels from  $Q$  and jointly controls the two mass ratios, so the identity constrains one combination of the spectrum exactly while leaving two ratios free. Equal rank-two event blocks in the finite tracial Gelfand–Naimark–Segal packet supply the balance  $\rho/a = 1/\sqrt{2}$  exactly under the declared tracial premises; the Lean development checks the circulant identity and the equivalence [13].

Under the balance and mass-ordering premises, the measured electron and muon masses [57] determine the tau mass through one quadratic, and the comparison is governed by a registered conditional test whose target definition, precision floor of 0.045 MeV, and decision rule were fixed before comparison with any post-registration measurement [3]. The outward-rounded enclosure is  $[1776.968991, 1776.969063]$  MeV, a 72-eV window whose center sits  $0.43\sigma$  from the measured  $1776.93 \pm 0.09$  MeV [57]. The premise ancestry is declared: the balance premise was first abstracted from the measured triple, so this is a target-informed conditional postdiction whose evidential weight lies in the kill direction. A world-average or dedicated-measurement central value more than three standard uncertainties from 1776.969027 MeV refutes the balanced-circulant premise, and with it the equal-rank-two-block tracial reading of the face-circulant sector, the only exact family-sector mechanism this framework offers. The physical family attachment and phase selection are the open realization maps of this surface, so it is logically independent of the diagnostic below.

### 9.3 The charged-lepton interval diagnostic

A finite eight-path carrier combined with an empirical electromagnetic transport packet yields outward-rounded enclosures for the three charged leptons with logarithmic half-width 1.732%, one-sided multiplicative widths  $-1.72\%$  and  $+1.75\%$ , under the declared payload-coherent anchor-gap premise; the premise-free independent-gap enclosures have half-width 6.554%. The measured triple lies inside all three coherent enclosures [12]. Measured mass ratios and transport anchors enter this branch, so the result is a diagnostic of internal closure: it gives a sharp surface that a source-emitted transport bridge must hit, without claiming a source-only mass prediction. A physical charged determinant line, an absolute clock, and a source-emitted transport bridge are the maps an endpoint prediction requires.

### 9.4 The electroweak chart and the certified pole stack

On the strict source-audit branch, the source chain emits the electroweak chart coordinates 80.330 and 91.119 GeV for the  $W$  and  $Z$  carriers; no measured mass enters that emission [12]. A certified strict one-loop contour stack closes the chart-to-pole step on its declared chart. A directed complex-interval receipt proves zero exclusion of the transverse inverse propagator on the declared principal-sheet boxes, and a companion receipt proves exactly one second-sheet pole of winding number one on the declared channel-by-channel continuation, with interval-Newton pole and residue enclosures on one fixed rational boundary partition replayed at three working precisions; fail-closed checkers recompute both verdicts from the payloads [1]. The chart coordinates sit at relative separations of  $4.9 \times 10^{-4}$  and  $7.6 \times 10^{-4}$  from the measured Breit–Wigner coordinates 80.3692(133) and 91.1880(20) GeV [57]. That comparison is a diagnostic under Table 1: the one-loop transport consumes imported Standard Model self-energy inputs, and an OPH-native source-to-pole packet is a premise of any stronger reading.

### 9.5 Finite capacity and the de Sitter transfer sign

The screen’s finite capacity supports an exact entropy accounting with one conditional gravitational reading [14].

**Proposition 9.2** (Capacity transfer sign). *For finite sector dimensions  $d_i$  with total  $M = \sum_i d_i$  and sector probabilities  $p_i$ , the generalized entropy obeys the exact identity*

$$S_{\text{gen}}(p, d) = - \sum_i p_i \log p_i + \sum_i p_i \log d_i = \log M - D\left(p \parallel \frac{d}{M}\right),$$

where  $D$  is relative entropy; its exact maximum is  $S_{\text{gen}}^{\text{max}} = \log M$  at  $p_i = d_i/M$ . If an observer receives a fraction  $f \in (0, 1)$  of a fixed total capacity and the sectors deplete uniformly, every admissible transfer changes the maximum by

$$\Delta S_{\text{gen}}^{\text{max}} = \log(1 - f) < 0,$$

and the positive-real interpolation in  $f$  is strictly decreasing and concave, so the zero-transfer point is a one-sided boundary maximum.

*Proof.* Expanding  $D(p \parallel d/M) = \sum_i p_i \log p_i - \sum_i p_i \log d_i + \log M$  gives the identity, and nonnegativity of relative entropy with equality at  $p = d/M$  gives the maximum. Uniform depletion replaces  $M$  by  $(1 - f)M$ , so the maximum changes by  $\log(1 - f)$ . Monotonicity and concavity of  $f \mapsto \log(1 - f)$  are elementary.  $\square$

Identifying capacity with horizon area, the transfer coordinate with observer mass, and the associated finite port operator with the gravitational shock supplies a conditional de Sitter time-advance mechanism with a definite sign: under those dictionaries an observer absorbing a capacity fraction  $f$  strictly lowers the maximal generalized horizon entropy by  $\log(1 - f)$ , and the one-sided boundary maximum carries the definite sign of the associated time shift. The finite identities make none of those physical identifications; the horizon and observer dictionaries are the open maps, and the finite calculation supplies no positive cyclic static-patch trace.

## 9.6 The unique rank-six invariant of the primitive carrier

In this subsection, “spin six” means spherical-harmonic angular rank  $j = 6$ . It does not denote a particle with spin six. The mathematical engine is the invariant count for the sixty-element proper icosahedral rotation group. The dimension of the invariant subspace in angular rank  $L$  is

$$m_L = \frac{1}{60} \left[ (2L + 1) + 15 U_{2L}(0) + 20 U_{2L}\left(\frac{1}{2}\right) + 12 U_{2L}\left(\frac{\varphi}{2}\right) + 12 U_{2L}\left(\frac{\varphi-1}{2}\right) \right],$$

where  $U$  denotes the Chebyshev polynomials of the second kind. The exact table gives  $m_1 = \dots = m_5 = 0$  and  $m_6 = 1$ . Thus no nonconstant invariant occurs below angular rank six, and the rank-six invariant line is one-dimensional, in agreement with the classical icosahedral-harmonic classification [34].

**Theorem 9.3** (Icosahedral angular-rank universality). *Let the carrier kinetics be any operator  $\lambda_a(k) = a^{-2} \sum_d w(d) [1 - \cos(a k \cdot d)]$  with a finite direction multiset and weights invariant under the proper icosahedral rotation group. Then the dispersion is exactly isotropic through angular rank five at every order in  $a$ ; every directional term below angular rank ten is one multiple of the unique normalized invariant  $\mathcal{I}_6$ , whose stationary structure is the exact 62-direction census (12 and 20 extrema of opposite index, 30 saddles); and whenever the weighted sixth-moment  $\mathcal{I}_6$  coefficient is nonzero, a condition certified for the equal-weight stencil and for each fundamental orbit, the first directional artifact appears at order  $a^4 k^6$  and one binary refinement step suppresses it by exactly  $1/16$  at that order. A vanishing sixth moment removes the displayed  $a^4 k^6$  residue. Any surviving*

*directional term occurs at higher order. For a single-radius member whose weighted sixth-moment coefficient vanishes, the rank-six component vanishes at every order, so any surviving anisotropy has a higher even invariant rank. Rank six is the least rank any member of the class can populate.*

*Proof.* The order- $a^{2m-2}k^{2m}$  angular content is an invariant polynomial of degree  $2m$ , with harmonic components at even spins  $L \leq 2m$ . The invariant table forces zero at  $L = 2, 4, 8$  and a one-dimensional space at  $L = 6$ , so every anisotropic term below angular rank ten is proportional to  $\mathcal{I}_6$ , and the possible artifact ranks are exactly the even invariant levels  $6, 10, 12, 16, \dots$ . The scaling in  $a$  is fixed by the expansion order, and the census is the exact critical-point theorem of the fingerprint certificate. The invariant table, the census, the constructive orbit verification, the nonvanishing sixth-moment coefficients, and an exact positive-weight tuned member whose spin-six content cancels are certified in exact arithmetic [1].  $\square$

The universality theorem concerns a class of finite operators. A physical prediction requires a narrower branch whose operator and transfer are fixed. Let  $u_1, \dots, u_{12}$  be the unit vectors through the twelve carrier ports. The *primitive twelve-port propagation branch* has the following premises. Its intrinsic spatial kinetic symbol belongs to the real, reciprocal, finite-range cosine class of Theorem 9.3, with scale-independent coefficients. The complete primitive orbit is its sole hop support, and no independent isotropic or directional kinetic term contributes through order  $k^6$ . Proper-carrier covariance acts on this orbit; the carrier scale  $a$  is finite and positive; the continuum quadratic term is normalized to  $k^2$ ; one carrier rest frame is transported coherently over the experiment; and the tested scalar or polarization-independent sector realizes the operator. A photon realization requires equal action on both transverse polarizations. An empirical fit must isolate the carrier contribution from source, medium, gravitational, and instrumental effects. Transitivity of the proper carrier group forces equal weights once the single primitive orbit is selected. The carrier itself is realized as a quasiperiodic structure or as a graph: no periodic three-dimensional lattice carries this point group, so the branch presupposes no crystal.

**Proposition 9.4** (Primitive twelve-port branch prediction). *Under these premises, the physical symbol is*

$$\omega^2(k, \hat{k}) = \frac{1}{2a^2} \sum_{i=1}^{12} \left[ 1 - \cos(ak u_i \cdot \hat{k}) \right]. \quad (19)$$

*With*

$$\mathcal{I}_6(n) = \frac{25}{132} \sum_{i=1}^{12} P_6(u_i \cdot n), \quad \mathcal{I}_6(u_i) = 1,$$

*where  $P_6$  is the degree-six Legendre polynomial. The small- $ak$  expansion is*

$$\omega^2 = k^2 - \frac{a^2}{20} k^4 + \frac{a^4}{840} k^6 + \frac{2a^4}{7875} k^6 \mathcal{I}_6(\hat{k}) + O(a^6 k^8). \quad (20)$$

*Writing the same expansion in a transported frame as*

$$\omega^2 = k^2 + C_4 k^4 + B_0 k^6 + B_6 k^6 \mathcal{I}_6(R^{-1} \hat{k}) + O(k^8), \quad R \in \text{SO}(3)/A_5,$$

*the branch predicts*

$$C_4 = -\frac{a^2}{20}, \quad B_0 = \frac{a^4}{840}, \quad B_6 = \frac{2a^4}{7875}, \quad (21)$$

*and hence the scale-free relations*

$$\frac{B_6}{C_4^2} = \frac{32}{315}, \quad \frac{B_0}{C_4^2} = \frac{10}{21}, \quad \frac{B_6}{B_0} = \frac{16}{75}. \quad (22)$$

All intrinsic anisotropic coefficients at  $1 \leq j \leq 5$  vanish. Once a negative  $C_4$  is resolved on this branch, no scale or anisotropic amplitude is free; only one three-parameter orientation class  $R$  in  $\text{SO}(3)/A_5$  is profiled. Under binary refinement,  $C_4(a/2) = C_4(a)/4$  and  $B_0(a/2) = B_0(a)/16$ ,  $B_6(a/2) = B_6(a)/16$  at fixed physical momentum.

*Proof.* The exact carrier moments are

$$\sum_i (u_i \cdot n)^2 = 4, \quad \sum_i (u_i \cdot n)^4 = \frac{12}{5}, \quad \sum_i (u_i \cdot n)^6 = \frac{12}{7} + \frac{64}{175} \mathcal{I}_6(n).$$

Substitution into the cosine series in (19) gives (20). Eliminating  $a$  gives (22). The lower-rank nulls and uniqueness of the rank-six line follow from Theorem 9.3.  $\square$

This proposition is a frozen prospective conditional physical-branch prediction [3]. The minimal locally Lorentz-invariant Standard Model plus General Relativity in local vacuum supplies the baseline  $C_4 = B_0 = B_6 = 0$  for intrinsic propagation. Nonminimal effective operators, media, curvature, or another icosahedral system can imitate some or all of the pattern. A match therefore distinguishes the branch from the minimal baseline without identifying OPH uniquely. Searches for Lorentz violation commonly organize anisotropic coefficients by angular rank  $j$  in the Standard-Model Extension [52, 53], which provides a natural coefficient language for a dataset-specific contract: at leading order the branch populates only the thirteen-component nonbirefringent rank-six multiplet at operator dimension eight, collapsed to one amplitude and one frame orientation, with every anisotropic coefficient at  $1 \leq j \leq 5$  exactly zero in the carrier rest frame and frame boosts admixing neighboring ranks at first order in the frame velocity.

The exposure class is fixed. The WMAP internal-linear-combination map, its cosmic-microwave-background likelihood class, and every data product examined in the dated template search are ineligible for this prediction. That search included the icosahedral rank-six and higher templates and returned a family-wide  $p$ -value of 0.64. The linked  $C_4$ ,  $B_0$ , and  $B_6$  relations have no eligible physical comparison. A separate exposed photon diagnostic identifies  $a = \sqrt{20}/E_{\text{QG},2}$  in the standard subluminal quadratic-dispersion convention. The published bound  $E_{\text{QG},2} > 10^{13}$  GeV gives  $a < 8.825 \times 10^{-29}$  m at 95% confidence; the LHAASO bound  $E_{\text{QG},2} > 6.9 \times 10^{11}$  GeV is a cross-check with a different source-lag treatment [54, 55]. These exposed retrospective limits cannot score the prospective linked-coefficient prediction. The target statement, exact coefficient snapshot, registration manifest, public commit custody, and detached OpenTimestamps calendar attestations fix the prospective content and its decision rule [3].

No empirical verdict is licensed without a dataset-specific registration that fixes one post-freeze release, a joint likelihood or full covariance for the same-sector  $C_4$ ,  $B_0$ , and rank-six coefficient vector, the carrier and observer frames, the  $\text{SO}(3)/A_5$  orientation profile, the boost law, the environmental and instrumental nuisance model, trials accounting, sensitivity floor, and calibrated joint threshold. The append-only corrected rule scores the complete branch manifold

$$C_4 < 0, \quad B_0 = \frac{10}{21} C_4^2, \quad B_6 = \frac{32}{315} C_4^2,$$

with every intrinsic coefficient at  $1 \leq j \leq 5$  equal to zero and the  $j = 6$  vector on the rotated  $\mathcal{I}_6$  orbit. The branch fails at five standard deviations or more if an isolated intrinsic  $C_4$  is positive, an isolated lower-rank coefficient is nonzero, the linked sixth-order terms are excluded at adequate sensitivity, or the calibrated profile likelihood excludes the complete manifold. Support requires exclusion of the zero-coefficient baseline at five standard deviations or more, agreement with the linked branch manifold within two standard deviations, rejection of named systematic alternatives,

and an independent replication. A null result is inconclusive because the branch supplies no positive lower bound on  $a$ . Incomplete covariance, insufficient sixth-order sensitivity after a negative  $C_4$ , or failure to isolate the carrier contribution is also inconclusive. A failed verdict rejects the primitive twelve-port physical propagation branch. It rejects OPH as a whole only if the physical sector bridge, coherent-frame transport, and exclusivity are proved forced.

*Boundary 9.5.* No statement in this section is a source-only determination of a physical constant or mass. The Koide identity and capacity identities are finite theorems; the tau window is a registered conditional test, postdictive in premise ancestry; the lepton enclosures and the screen-grain separation are diagnostics that consume measured inputs; the electroweak chart comparison is a diagnostic whose pole transport imports Standard Model self-energies; the de Sitter reading is conditional on unconstructed dictionaries. The angular-rank theorem is a finite theorem. The primitive-port coefficient relations form a prospective prediction on the named physical branch, with empirical scoring sealed until the required dataset-specific contract exists.

## 10 Physical interpretation maps and their consequences

The preceding theorems concern finite records, explicit response maps, and reconstruction implications. They become claims about nature only when a physical carrier realizes the operations and identifications in Definition 1.2. Table 3 records the resulting implications and the premise that can fail.

Table 3: Physical questions, reconstruction consequences, and required premises.

| Question                                                    | Consequence of the stated realization                                                                                                                                               | Physical premise or open map                                                                                                                             |
|-------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------|
| When is a measurement public?                               | A completed record is a schedule-independent normal form; on the declared algebra-state surface its event projectors obey Theorem 4.1.                                              | A laboratory system must instantiate durable records, readback, protected boundaries, and the algebra-state map.                                         |
| When is event geometry Lorentzian?                          | A common record-germ tower satisfying (E1)–(E7) gives the Lorentzian four-manifold of Theorem 6.2; the finite instrument separately measures inertia $(1, 3)$ on its declared path. | A physical refinement family must supply the common-refinement pseudometric, population, covering charts, compatible tetrads, cone, and causal premises. |
| What equation governs the gravitational response?           | A realized null balance, Ward conservation, the Bianchi identity, and an independent scale identification give the Einstein field equation of Theorem 7.2.                          | The common physical tower, continuum control, clock, stress, area, vacuum, and scale maps are the open realization maps.                                 |
| How do the gravitational and gauge branches share a source? | One carrier can feed the event/modular branch and the twelve-port response branch, yielding both conclusions from a common source.                                                  | Joint physical realization and equality of $G_{\text{packet}}$ with $G_{\text{Tan}}$ are separate identification premises.                               |

Table 3 continued

| Question                                                                                                                | Consequence of the stated realization                                                                                                                                                                                           | Physical premise or open map                                                                                                                 |
|-------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------|
| Why this gauge type and matter image?                                                                                   | The complete A1 response and endogenous A2 transport force the compact local Lie type. The separately supplied matter representation gives the conditional $\mathbb{Z}_6$ quotient and anomaly-free rank-fifteen finite module. | Laboratory currents, quantum fields, physical matter poles, family multiplicity, and masses require further maps.                            |
| What propagation signature distinguishes the primitive-port branch from the minimal locally Lorentz-invariant baseline? | Proposition 9.4 fixes $B_0$ and $B_6$ from $C_4$ and fixes the complete rank-six shape up to one carrier orientation.                                                                                                           | The scalar or polarization-independent sector bridge, coherent frame transport, carrier-term isolation, and exclusivity are branch premises. |

### 10.1 Gravity through the modular-entropy map

The route runs from patches to gravity in one chain: consensus supplies public record classes, a record-germ realization satisfying (E1)–(E7) supplies the event manifold, modular flow and half-sided inclusions supply the local time generator and null translations on the same tower, and the entropy split with its stationarity conditions would then have to establish the null-balance premise that Theorem 7.2 completes. This is a route with a visible break point. The normal-form, entropy, and tomography statements have exact finite layers. The passage to gravity starts only when one physical tower realizes the event, modular, continuum, universal-coupling, vacuum, and scale premises together.

*Intuition.* Under the stated realization map, the gravitational response represents the large-scale compatibility condition associated with repaired records. Local observers extend and compare records; the metric describes the stable causal relation among those records. Modular flow describes how a patch reads change, and entropy stationarity fixes the response needed for the local accounts to stay compatible.

### 10.2 A common source for gravitational and gauge branches

In this framework, the proposed fundamental theory that combines the gravitational and gauge sectors is one observer-consensus tower with two readout branches. The overlap, record, and repair structure feeds the event and modular branch. The twelve-port incidence, complete reversible response, and endogenous overlap transport feed the local gauge-type branch. Trace-balanced blocks supply a separate conditional matter branch. A single self-reading carrier could supply both sets of antecedents.

This common source neither identifies gravity with a gauge boson nor fixes all couplings. It also leaves two gauge constructions distinct:  $G_{\text{Tan}}$  is reconstructed from a sector category, while  $G_{\text{packet}}$  is the maximal faithful image of the explicit finite response and matter module. A completed unification requires joint physical realization and a proof that the corresponding current diagrams commute.

*Intuition.* One object performs two tasks. Its overlap structure organizes events and the gravitational response; its boundary response organizes charges. The proposed architecture takes the



carrier and its self-reading dynamics as their common source, with spacetime and gauge structure appearing as distinct public readouts.

### 10.3 Completed records as measurement events

The finite construction locates a measurement at the transition from a private, repairable record to a protected public normal form. Event projectors, weights, and state updates are then defined on the completed algebra-state surface. A physical application must identify actual durable records and verify the transaction and readback premises. The theorem does not construct the state space of an interacting continuum field theory.

*Intuition.* Within the model, a record is called public when it has survived every authorized comparison and no repair can change it. Probability describes the completed record surface; state update moves from one public record class to the next. The construction locates observation in the bookkeeping that patches perform on their shared boundaries.

### 10.4 Event geometry from record germs

The spherical support identifies the Lorentz group and the hyperbolic space of timelike directions without supplying physical events. Events enter only through the direct limit of separating record germs. Theorem 6.2 states a sufficient gluing criterion: four-component charts with open images, compatible affine transitions, tetrads, a Lorentzian quadratic form, and causal reachability must all occur on one populated refinement limit. The criterion packages these inputs into one Lorentzian manifold; it does not select four dimensions or Lorentz signature from weaker data. The finite instrument measures rank and inertia on a prescribed chart, while its negative cone margins and absent cofinal limit prevent continuum promotion.

*Intuition.* Record germs say when two finite histories describe the same event and how nearby events can be distinguished. A spacetime appears only when those local descriptions cover the completed event set, agree on overlaps, and carry one compatible causal cone. The manifold records the stable way in which public events fit together.

### 10.5 Gauge type and the finite matter image

The twelve-port coefficients split under the alternating-group action into bands of dimensions 1, 3, 3, and 5. A1 supplies their complete compact response algebra, while A2 implements every proper carrier action internally on that same response. Theorem 8.2 forces one central direction together with simple ideals of dimensions three and eight. The response map  $K$  is an exact conditional matrix witness for this algebra. The matter space  $V$  is a separate supplied representation; its exterior algebra gives the familiar fifteen-state branching, and the exhaustive scan proves the two-mask selection inside its declared ten-summand menu. Physical gauge fields, matter poles, families, and masses require current and continuum realization maps.

*Intuition.* The carrier exposes twelve response channels. Icosahedral symmetry groups them into four inequivalent bands. Complete reversible response closes their public tangent under commutators, and observer agreement requires carrier rechartings to act inside that same response. Compactness and the single fixed line then leave the Standard Model gauge Lie type. A separate matter module tests which charges coexist without anomalies; it does not turn the finite channels into observed particles.

The conditional implications share premises. Failure of the common physical tower removes the event-manifold and Einstein conclusions together. Failure of laboratory-current attachment removes the gauge and matter interpretations together. The finite normal-form, invariant-theory,

and algebraic results retain their stated domains. Table 3 summarizes the principal physical interpretation claims made here.

## 11 Machine verification and reproducibility

Each result maps to an artifact class:

- a Lean library with no admitted propositions in the dependency closure reported here, including formal premise boundaries and countermodels;
- exact-arithmetic code receipts: integer and rational computations, interval certificates with outward rounding, and  $\mathbb{F}_2$  rank computations, each emitting a canonical hashed payload;
- simulation receipt bundles pinning the applicable source or module revision, configuration, seed where stochastic, grids, tolerances, and output hashes, with deterministic reserialization distinguished from fresh source replay;
- verifiers that recompute verdicts from clause vectors and reject receipts whose stored verdict disagrees, so a caller cannot assert a truth flag directly;
- adversarial negative controls: every certificate ships with mutations that must be detected, and a control that cannot fail is treated as a defect of the certificate, not as support;
- provenance rules under which machine-readable ledgers assign every quantitative row a typed class (source result, reconstruction implication, diagnostic, registered test, or rejected candidate in the claim ledgers; tiered ancestry labels in the postdiction ledger) and reject a stronger classification unsupported by the recorded dependencies.

The repository, receipts, schemas, and rebuild instructions are public [1, 2]. The formal and executable stack checks the claims assigned to it and enforces their stated boundaries. Analytic arguments in this manuscript retain their displayed mathematical proofs, and no part of the stack establishes physical realization.

### 11.1 Artifact map

Table 4 identifies the principal artifacts for each result. Repository continuous integration checks the theorem-count floor, rejects admitted proofs in the public library, and rebuilds the repository-local finite certificates used here. Simulator replay is governed by the simulator receipts and reproduction commands cited below.

Table 4: Principal verification artifacts and their scope.

| Result                                                       | Principal artifacts                                                                                                                                                                                                         |
|--------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Observable normalizer, stability, refinement, and complexity | <code>Lean/ObservableNormalForms/</code> and the exact proofs and reductions accompanying <code>extra/observable_normal_forms.tex</code>                                                                                    |
| Transactional diamond and consensus normal form              | general local-diamond and Newman arguments proved in this manuscript; the observation-relative endpoint-uniqueness module of the Lean library; the finite transaction-model verifier with its receipt and negative controls |

Table 4 continued

| Result                                                         | Principal artifacts                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
|----------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Event algebra, Born, Lüders, Tsirelson                         | <code>Lean/EventAlgebra/</code> (Basic, Lueders, Tsirelson, ExpectationBound, partition and state modules); per-module axiom audits                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| Central defects and edge-center entropy                        | overlap-cocycle and one-sided reduction proofs in this manuscript; <code>Lean/ObserverPatchHolography/EinsteinBranch/Entropy.lean</code> for scalar entropy bookkeeping after the split; <code>Lean/ObserverPatchHolography/CollarStates.lean</code> for the identity-channel model and its no-go boundary                                                                                                                                                                                                                                                                                                                                                                              |
| Port action, A1–A2 Lie type, and conditional matrix algebra    | files under <code>Lean/Screen/</code> : <code>A5PortAction.lean</code> , <code>A5Commutant.lean</code> , <code>A5IncidenceResponse.lean</code> , <code>A2HolonomyBridge.lean</code> for the fixed-space and endogenous-transport interfaces, and <code>Compact12.lean</code> for the abstract matrix algebra; the conditional port map and $A_5$ covariance are checked by <code>code/a5_closure/port_current_inner_certificate.py</code> , <code>code/a5_closure/manifests/port_current_response_reference.json</code> , and <code>code/a5_closure/receipts/port_current_inner_reference.receipt.json</code> ; compact-simple classification is the analytic input used in Theorem 8.2 |
| Global form, $\mathbb{Z}_6$ kernel, and exterior selection     | <code>Lean/Screen/Z6Exact.lean</code> for the abstract lattice quotient, <code>TraceBalancedKernel.lean</code> for central-parameter arithmetic, and <code>ExteriorSelection.lean</code> for the declared finite scan                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| Electroweak chart and certified $W/Z$ pole stack               | <code>code/particles/calibration/wz_upstream_completion/outputs/certified_wz_contours.json</code> and <code>code/particles/calibration/wz_upstream_completion/outputs/certified_second_sheet_poles.json</code> , their schemas, fail-closed checkers, and byte-exact producer replays                                                                                                                                                                                                                                                                                                                                                                                                   |
| Icosahedral angular-rank theorem and primitive-port prediction | <code>Lean/Screen/A5PrimitivePortPrediction.lean</code> for equal weights, rational coefficient ratios, and refinement arithmetic, excluding the physical-sector bridge; <code>code/a5_fingerprint/runtime/a5_multiple_fixed_point_receipt.json</code> , <code>code/a5_fingerprint/runtime/spin_six_universality_receipt.json</code> , <code>code/a5_fingerprint/runtime/spin_six_primitive_port_prediction_receipt.json</code> , their mutation tests, and the custody-bound prediction register                                                                                                                                                                                       |
| Record-germ event-manifold gluing                              | analytic theorem proved in this manuscript; finite chart, transition, and signature receipts test only selected antecedents                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |

Table 4 continued

| Result                                                        | Principal artifacts                                                                                                                                                                                                                                                                                                  |
|---------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Null tomography, small-ball arithmetic, and tensor completion | <code>Lean/ObserverPatchHolography/EinsteinBranch/Tensor.lean</code> (null directions, design/decoder, injectivity, and metric ambiguity); <code>Lean/docs/EINSTEIN_BRANCH_INDEX.md</code> ; analytic determinant and norm calculations in this manuscript; explicit null-balance, Ward, Bianchi, and scale premises |
| Signature ladder and control                                  | <code>evidence/einstein_convergence/</code> (four-row manifest, hashed arrays; pinned simulator revision and replay comparison procedure in its README)                                                                                                                                                              |
| Finite local source domain                                    | staged receipts under <code>data/local_domain/</code> and the bundle verifier under <code>oph_fpe/local_domain/</code> in the simulator repository                                                                                                                                                                   |

## 12 Empirical meaning and falsifiability

The custody-bound register fixes target definitions, exclusion thresholds, precision floors, exposure classes, and decision rules before eligible comparison [3]. Proposition 9.4 is the prospective discriminator in this paper. Its coefficient relation is fixed, its related cosmic-microwave-background exposure is excluded, and its dataset-specific comparison is sealed under the conditions stated in Section 9.6. A qualifying failure rejects the primitive twelve-port physical propagation branch. OPH-wide scope requires a theorem that makes this physical branch forced and exclusive.

The gravitational and gauge conclusions have separate empirical boundaries. The Einstein composition requires one physical refinement tower satisfying its event, modular, stress, entropy, continuum, vacuum, and scale premises together. The gauge and matter conclusions require laboratory-current attachment and a continuum operator limit. Failure to construct those maps leaves the corresponding finite theorems intact and removes their claimed physical application. Non-identifiability theorems determine which finite source interfaces cannot supply the missing information.

## 13 Discussion and comparison

Operational reconstructions derive quantum structure from information and composition principles [15, 16, 17]. OPH makes a narrower statement at the quantum stage: a completed finite record, once represented by an algebra-state pair, carries the standard event, conditioning, expectation, and correlation identities. The distinctive input is the observer-indexed normal form and its transaction contract. Unlike those reconstructions, which derive the quantum state space itself from operational axioms, OPH takes the finite algebra-state representation as an explicit input; its contribution is the completed-record surface on which the identities operate and the consensus theorem that defines completion.

Two further neighbors calibrate the record story. Quantum Darwinism locates classical objectivity in redundant environmental records accessible to many observers [29]; the completed-consensus record plays the same public role, with redundancy replaced by protected overlap agreement and an explicit repair dynamics. The cellular-automaton program reconstructs quantum mechanics from a deterministic finite substrate [30]; OPH shares the finite substrate and differs by locating the quantum identities on the completed record surface rather than in an underlying ontological basis. Wigner’s-friend extensions constrain theories whose agents reason about one another’s unfinished

measurements [31]; the public records of this framework are post-consensus objects by construction, and the no-go premises concern pre-consensus perspectives that the completed surface does not represent.

Relational quantum mechanics treats physical facts as relational [27]; the thermal-time program reads time from modular structure [28]. OPH combines these themes by attaching public facts to protected overlap records and consuming normalized modular flow only on a typed common tower. Its local-diamond proof also has a direct distributed-systems neighbor in asynchronous agreement [24, 25].

The gravitational branch belongs to the family of thermodynamic and entanglement-based routes to geometry [38, 39, 42]. Bisognano–Wichmann modular flow and half-sided modular inclusions supply established analytic ingredients [35, 36, 37]. The contribution of Theorem 7.2 is the explicit composition contract: event, modular, stress, entropy, asymptotic, vacuum, and scale premises appear in one statement, while the finite null ambiguity identifies exactly where a metric-proportional term enters. The route from a null-projected balance and the Bianchi identity to the full equation, with  $\Lambda$  as an integration constant, is familiar from trace-free and unimodular formulations [40, 41]; the contribution here is the finite nine-direction design with exact determinant and decoder constants and the premise contract (G1)–(G6) stated as one implication.

The gauge section contains two reconstruction routes and one conditional global-form witness. Doplicher–Roberts/Tannaka reconstruction supplies the structural sector group [46, 47]. Independently, complete A1 response and endogenous A2 transport force the local Standard Model gauge Lie algebra. The explicit matrix map and exterior-module scan give the conditional packet group. Equality of the sector group and packet group is a separate commuting-square premise. This separation prevents the categorical reconstruction, local Lie-type theorem, and physical global group from being conflated.

For a foundations readership, the methodological claim is also concrete. Finite theorems, certified computations, reconstruction implications, and physical realization maps are different mathematical objects. Keeping those types visible makes every emergence claim auditable and gives each claim a specific failure mode. A3 also places agreement constraints before state selection. The physical adequacy of that ordering depends on the realization premises stated in Section 12.

## 14 Conclusion

For the specified finite observer-patch architecture, observable fibers give a canonical partial normalizer, stability moduli control approximate and refined outputs, and succinct boundary problems have explicit complexity barriers. On the transactional carrier, semantic dependency closure and revalidation yield the local diamond, so quadratic descent gives a schedule-independent public record. The completed finite algebra-state surface then carries the standard probability, conditioning, expectation, and correlation identities.

The principal finite gauge result is the A1–A2 forcing theorem. Complete reversible response supplies a compact commutator-closed twelve-dimensional current, and endogenous overlap transport makes the proper carrier action inner on that same current. The one-dimensional fixed space excludes the centerless  $\mathfrak{su}(2)^4$  alternative, leaving  $\mathfrak{u}(1) \oplus \mathfrak{su}(2) \oplus \mathfrak{su}(3)$ . The declared four-band map is a conditional matrix witness for this type. Trace balance and the exterior module give a conditional maximal faithful matter image with common  $\mathbb{Z}_6$  kernel and an anomaly-free rank-fifteen representation; the exhaustive scan isolates its charge-conjugate pair within the stated class. The physical global quotient is not selected by this result.

The quantitative results include one prospective physical-branch prediction. The positive-

chamber circulant identity forces the Koide relation exactly at tracial balance and yields a registered conditional tau test with a custody-bound kill band; the capacity identity fixes the de Sitter transfer sign; the interval-certified screen-grain root and the charged-lepton enclosures are diagnostics whose transport maps are typed as open realization maps. The source-emitted electroweak chart coordinates carry a certified strict one-loop pole stack, a principal-sheet zero-exclusion certificate together with exactly one second-sheet pole of winding number one on the declared continuation, with the imported Standard Model self-energy inputs displayed as its premise. On the primitive twelve-port propagation branch,  $C_4$  fixes  $B_0$ ,  $B_6$ , and the complete rotated angular-rank-six template through the scale-free ratios (22). The minimal locally Lorentz-invariant Standard Model plus General Relativity predicts zero intrinsic coefficients on the stated local-vacuum baseline. The branch prediction is fixed before an eligible comparison, with its physical-sector, coherent-frame, isolation, and exclusivity premises exposed. The sixth-order isotropic term is smaller than the quartic correction by  $(ak)^2/42$ , so a decisive test requires sensitivity beyond a measurement of  $C_4$  alone. No comparison-ready dataset is asserted.

The geometry and dynamics results have a different status. A Lorentz-group action and its hyperbolic homogeneous space follow from the oriented spherical support; their physical interpretation requires event, clock, and frame maps. A Lorentzian event manifold follows only under the displayed event premises. The Einstein field equation follows only from one common tower satisfying the modular, stress, entropy, asymptotic, vacuum, and scale premises. The physical tower, continuum limit, laboratory-current attachment, quantum-field realization, and the listed interpretation maps are the open realization maps of the program. The paper therefore offers a finite reconstruction framework with exact results and explicit physical hypotheses; it does not claim a completed derivation of all physics, particle masses, a cosmology, or a proof that the simulator describes our universe. Its closure hypothesis, three axioms, named dynamics, finite consequences, and physical realization maps are distinct parts of one logical construction.

The ambition of the program is stated here at full strength, in its proper place after the results and their boundaries. The goal is a complete description of physical reality derived from consistency requirements alone: every dimensionless constant, every structural property, and every dynamical law recovered as the unique content of a self-consistent, self-reading observer system, with no free parameter and no external input. On the closure hypothesis of Section 1.3, that goal addresses two classical questions at once. The answer offered to why anything exists is that a timeless self-referential structure requires no external cause: existence coincides with self-consistency, and observers arise inside the structure because a self-explaining structure must contain its own description. The answer offered to why exactly this world is that the consistency requirements are far more selective than they first appear: the results assembled here force the local Standard Model gauge type from twelve ports and two axioms, force the rigid spin-six fingerprint from the same incidence, fix the Koide relation exactly at the certified balance, and pose exact closure equations for the fine-structure coupling and the cosmological capacity whose displayed candidates stand within  $2.5 \times 10^{-6}$  and one percent of the measured coordinates as typed diagnostics.

The distance between that ambition and the present results is recorded in this paper's own typing rather than left to interpretation. The finite theorems are proved; the quantitative closures are typed as diagnostics or registered tests; the physical realization maps are named, with no construction asserted; and the frozen branch prediction is armed with scale-free coefficient relations and a registered kill rule. The program therefore submits its ambition in the only scientifically admissible form: a sequence of exact claims each of which can fail, a registered fingerprint that experiments can match or refute, and a boundary ledger stating exactly what is unproved. Whether the full description exists is undecided; the contribution of this paper is to make the question precise, its completed stages machine-audited, and its registered surfaces falsifiable.

## Reproducibility statement

All theorems, certificates, simulation configurations, receipts, and the formal library are public in the project repositories [1, 2]. The paper release manifest records the release identifier (r2011) together with PDF hashes and sizes. Computational receipt bundles pin the applicable source or module revision, configuration, seed where stochastic, and output hashes. Clean-checkout rebuild instructions and negative controls accompany the artifacts. The claim ledger and prediction register are machine readable [3].

The Lean source tree, pinned toolchain, exact verifiers, manifests, receipts, negative controls, tests, finite signature ladder, and local-domain evidence are identified in Table 4. The complete simulation source is maintained in the public simulator repository at the evidence-producing revisions [2].

This manuscript synthesizes and strengthens results whose longer proofs also appear in public component preprints on observable normal forms, finite event algebras, consensus, Einstein reconstruction, and gauge structure [5, 6, 7, 10, 11]. The component relationship is disclosed here so that overlap is visible.

## Statements and Declarations

**Funding.** The authors declare that no external grants or dedicated third-party research funding were received for the preparation of this manuscript.

**Author affiliation.** Bernhard Mueller is affiliated with Pragma Research Inc., Washington State, United States.

**Author contributions.** Bernhard Mueller developed the OPH concept and mathematical synthesis, wrote the main manuscript, and coordinated the formal and computational evidence. Alexander Osika contributed the physical-realization and hardware boundary, prototype framing, and manuscript review. Mario Ponder contributed the finite port-current construction and verifier, gauge and matter certificates, and proof review of the edge-entropy and Einstein-normalization branches. Kai Xue contributed consistency checks, simulator review, visualization review, and prototype-engineering review. Ben Cassie contributed implementation-claim and physical-evidence review. Peter Nguyen contributed proof auditing, branch-boundary analysis, compact-gauge and global-quotient analysis, and manuscript consistency review. Maarten Antoine Visser contributed consensus-protocol review and physics feedback on the foundations and emergent-spacetime framing. Kale Arnav Anirudha contributed consensus-paper material, asynchronous-agreement analysis, repair-map definitions, and formal review of the protocol surface. David Matscheko contributed proof review of consensus repair, observable normal forms, modular and Einstein algebra, edge entropy, hypercharge, and the  $\mathbb{Z}_6$  quotient. Jonathan Hill contributed Lean formalization and proof auditing of observable normal forms, refinement and repair results, complexity classifications, and artifact coverage. William T. Glynn contributed simulator development, computational implementation, reproducible-build infrastructure, and release-validation work in the main OPH research repository. All authors reviewed and approved the manuscript. All authors agree to be accountable for all aspects of the work.

**Corresponding author.** Correspondence should be addressed to Bernhard Mueller.

Email: [bernhard@floatingpragma.ai](mailto:bernhard@floatingpragma.ai).

**Data availability.** The finite witnesses, simulation configurations, receipts, claim ledgers, and release manifests supporting this study are available in the public OPH and simulation repositories [1, 2]. Individual receipts and their bundle documentation provide the source and output provenance

for computational claims. The artifact map identifies the finite signature-ladder and local-domain evidence cited in the manuscript.

**Code availability.** The Lean sources, exact-arithmetic verifiers, finite scans, simulation code, tests, and rebuild instructions are available in the same public repositories [1, 2]. The computational claims in this manuscript identify their principal files in Section 11.

**Ethics approval and consent to participate.** Not applicable. The study involved no human participants, human data, or animals.

**Consent for publication.** Not applicable.

**Competing interests.** Bernhard Mueller is affiliated with Pragma Research Inc. The authors participate in OPH-related research, software, simulation, or prototype-development programs and may receive professional or reputational benefit from this work. These relationships are disclosed. The authors declare no other competing interests.

## References

- [1] FloatingPragma (2026). Observer Patch Holography: Lean library, exact code receipts, finite certificates, and claim ledgers. <https://github.com/FloatingPragma/observer-patch-holography>
- [2] Mueller, B. (2026). OPH physics simulator and reproduction source. <https://github.com/muellerberndt/oph-physics-sim>. Signature-ladder revision: <https://github.com/muellerberndt/oph-physics-sim/tree/4f0169c0e319468112d076afe6ce5aa14fbfe0d5>. Finite local-domain revision: <https://github.com/muellerberndt/oph-physics-sim/tree/d99ca548a4853e83f819a3a2c9d813f7a3429bdb>.
- [3] FloatingPragma (2026). Precommitted comparison certificates, custody records, and the frozen prediction register. Immutable source receipt: [https://github.com/FloatingPragma/observer-patch-holography/blob/66176656dc1143f9ec50ba1a6e409c403545857f/code/a5\\_fingerprint/runtime/spin\\_six\\_primitive\\_port\\_prediction\\_receipt.json](https://github.com/FloatingPragma/observer-patch-holography/blob/66176656dc1143f9ec50ba1a6e409c403545857f/code/a5_fingerprint/runtime/spin_six_primitive_port_prediction_receipt.json). Append-only custody packet: [https://github.com/FloatingPragma/oph-meta/tree/8cc5261653e37cbca0e6017fcc95a9fe7f649963/falsification/frozen\\_targets/fz11\\_2026-07-31](https://github.com/FloatingPragma/oph-meta/tree/8cc5261653e37cbca0e6017fcc95a9fe7f649963/falsification/frozen_targets/fz11_2026-07-31). Koide custody packet: [https://github.com/FloatingPragma/oph-meta/tree/df097d8fe7c38d008a1ba7827f7d573286ae2012/falsification/frozen\\_targets/fz10\\_2026-07-28](https://github.com/FloatingPragma/oph-meta/tree/df097d8fe7c38d008a1ba7827f7d573286ae2012/falsification/frozen_targets/fz10_2026-07-28). Live register: [https://github.com/FloatingPragma/observer-patch-holography/blob/main/claims/frozen\\_prediction\\_register.json](https://github.com/FloatingPragma/observer-patch-holography/blob/main/claims/frozen_prediction_register.json).
- [4] FloatingPragma (2026). *The OPH Axiom Reference*. [https://github.com/FloatingPragma/observer-patch-holography/blob/ff1981be069d3e5fa70d953e27b3786dd4c2b99d/docs/AXIOM\\_REFERENCE.md](https://github.com/FloatingPragma/observer-patch-holography/blob/ff1981be069d3e5fa70d953e27b3786dd4c2b99d/docs/AXIOM_REFERENCE.md)
- [5] Mueller, B., Matscheko, D., and Hill, J. (2026). *Observation-Determined Normal Forms: Stability, Obstructions, and Refinement in Constraint and Rewrite Systems*. Public manuscript. [https://github.com/FloatingPragma/observer-patch-holography/blob/ff1981be069d3e5fa70d953e27b3786dd4c2b99d/extra/observable\\_normal\\_forms.pdf](https://github.com/FloatingPragma/observer-patch-holography/blob/ff1981be069d3e5fa70d953e27b3786dd4c2b99d/extra/observable_normal_forms.pdf)
- [6] Mueller, B. (2026). *Verified Projection-Event Calculus in Lean 4: Bundled Arbitrary-Partition Pinching, Lüders Retractions, and CHSH Interoperability*. Public manuscript and formal artifact. [https://github.com/FloatingPragma/observer-patch-holography/blob/ff1981be069d3e5fa70d953e27b3786dd4c2b99d/extra/machine\\_checked\\_finite\\_event\\_algebras.pdf](https://github.com/FloatingPragma/observer-patch-holography/blob/ff1981be069d3e5fa70d953e27b3786dd4c2b99d/extra/machine_checked_finite_event_algebras.pdf)



- [7] Mueller, B., Xue, K., Anirudha, K. A., Matscheko, D., and Hill, J. (2026). *Reality as a Consensus Protocol: The Fixed-Point Computation That Implements Physics*. Public manuscript. [https://github.com/FloatingPragma/observer-patch-holography/blob/ff1981be069d3e5fa70d953e27b3786dd4c2b99d/paper/reality\\_as\\_consensus\\_protocol.pdf](https://github.com/FloatingPragma/observer-patch-holography/blob/ff1981be069d3e5fa70d953e27b3786dd4c2b99d/paper/reality_as_consensus_protocol.pdf)
- [8] Mueller, B., Osika, A., Xue, K., Cassie, B., Visser, M. A., and Matscheko, D. (2026). *Federated Echosphedral Screen Microphysics: Patch Hardware, Records, and Observer Synchronization in Observer-Patch Holography*. Public manuscript. [https://github.com/FloatingPragma/observer-patch-holography/blob/7260dc7c4ac05f84c021167f1e095a58998573f1/paper/screen\\_microphysics\\_and\\_observer\\_synchronization.pdf](https://github.com/FloatingPragma/observer-patch-holography/blob/7260dc7c4ac05f84c021167f1e095a58998573f1/paper/screen_microphysics_and_observer_synchronization.pdf)
- [9] Mueller, B., Osika, A., Ponder, M., Xue, K., Cassie, B., Nguyen, P., Visser, M. A., Anirudha, K. A., Matscheko, D., Hill, J., and Glynn, W. T. (2026). *From Observer Consensus to Standard Physics*. Public manuscript. [https://wkaxfdgxoqmgshymt.supabase.co/storage/v1/object/public/papers/from\\_observer\\_consensus\\_to\\_standard\\_physics.pdf](https://wkaxfdgxoqmgshymt.supabase.co/storage/v1/object/public/papers/from_observer_consensus_to_standard_physics.pdf)
- [10] Mueller, B., Osika, A., Ponder, M., Xue, K., Nguyen, P., Visser, M. A., and Matscheko, D. (2026). *Recovering Observer Spacetime and Einstein Dynamics from Overlap Consistency*. Public manuscript. [https://github.com/FloatingPragma/observer-patch-holography/blob/ff1981be069d3e5fa70d953e27b3786dd4c2b99d/paper/recovering\\_observer\\_spacetime\\_and\\_einstein\\_dynamics\\_from\\_overlap\\_consistency.pdf](https://github.com/FloatingPragma/observer-patch-holography/blob/ff1981be069d3e5fa70d953e27b3786dd4c2b99d/paper/recovering_observer_spacetime_and_einstein_dynamics_from_overlap_consistency.pdf)
- [11] Mueller, B., Osika, A., Ponder, M., Xue, K., Nguyen, P., Visser, M. A., and Matscheko, D. (2026). *Deriving Standard Model Gauge Structure from Observer Overlap Consistency*. Public manuscript. [https://github.com/FloatingPragma/observer-patch-holography/blob/ff1981be069d3e5fa70d953e27b3786dd4c2b99d/paper/deriving\\_standard\\_model\\_gauge\\_structure\\_from\\_observer\\_overlap\\_consistency.pdf](https://github.com/FloatingPragma/observer-patch-holography/blob/ff1981be069d3e5fa70d953e27b3786dd4c2b99d/paper/deriving_standard_model_gauge_structure_from_observer_overlap_consistency.pdf)
- [12] Mueller, B., Osika, A., Ponder, M., Xue, K., Visser, M. A., and Matscheko, D. (2026). *Deriving the Particle Zoo from Observer Consistency*. Public manuscript. [https://github.com/FloatingPragma/observer-patch-holography/blob/7260dc7c4ac05f84c021167f1e095a58998573f1/paper/deriving\\_the\\_particle\\_zoo\\_from\\_observer\\_consistency.pdf](https://github.com/FloatingPragma/observer-patch-holography/blob/7260dc7c4ac05f84c021167f1e095a58998573f1/paper/deriving_the_particle_zoo_from_observer_consistency.pdf)
- [13] Mueller, B. (2026). *The Positive-Chamber Koide Identity for Icosahedral Face Circulants*. Public manuscript and formal artifact. [https://github.com/FloatingPragma/observer-patch-holography/blob/7260dc7c4ac05f84c021167f1e095a58998573f1/extra/koide\\_identity\\_from\\_positive\\_c3\\_face\\_circulants.pdf](https://github.com/FloatingPragma/observer-patch-holography/blob/7260dc7c4ac05f84c021167f1e095a58998573f1/extra/koide_identity_from_positive_c3_face_circulants.pdf)
- [14] Mueller, B. (2026). *The de Sitter Time-Advance Sign from a Finite Screen with Fixed Capacity*. Public manuscript. [https://github.com/FloatingPragma/observer-patch-holography/blob/7260dc7c4ac05f84c021167f1e095a58998573f1/extra/de\\_sitter\\_time\\_advance\\_sign\\_from\\_fixed\\_screen\\_capacity.pdf](https://github.com/FloatingPragma/observer-patch-holography/blob/7260dc7c4ac05f84c021167f1e095a58998573f1/extra/de_sitter_time_advance_sign_from_fixed_screen_capacity.pdf)
- [15] Hardy, L. (2001). “Quantum theory from five reasonable axioms.” arXiv:quant-ph/0101012. <https://arxiv.org/abs/quant-ph/0101012>
- [16] Chiribella, G., D’Ariano, G. M., and Perinotti, P. (2011). “Informational derivation of quantum theory.” *Physical Review A* **84**, 012311. <https://doi.org/10.1103/PhysRevA.84.012311>
- [17] Masanes, L., and Müller, M. P. (2011). “A derivation of quantum theory from physical requirements.” *New Journal of Physics* **13**, 063001. <https://doi.org/10.1088/1367-2630/13/6/063001>

- [18] Born, M. (1926). “Zur Quantenmechanik der Stoßvorgänge.” *Zeitschrift für Physik* **37**, 863–867. <https://doi.org/10.1007/BF01397477>
- [19] Lüders, G. (1950). “Über die Zustandsänderung durch den Meßprozeß.” *Annalen der Physik* **443**, 322–328. <https://doi.org/10.1002/andp.19504430510>
- [20] Clauser, J. F., Horne, M. A., Shimony, A., and Holt, R. A. (1969). “Proposed experiment to test local hidden-variable theories.” *Physical Review Letters* **23**(15), 880–884. <https://doi.org/10.1103/PhysRevLett.23.880>
- [21] Cirel’son, B. S. (1980). “Quantum generalizations of Bell’s inequality.” *Letters in Mathematical Physics* **4**(2), 93–100. <https://doi.org/10.1007/BF00417500>
- [22] Umegaki, H. (1954). “Conditional expectation in an operator algebra, I.” *Tôhoku Mathematical Journal* **6**(2–3), 177–181. <https://doi.org/10.2748/tmj/1178245177>
- [23] Takesaki, M. (1972). “Conditional expectations in von Neumann algebras.” *Journal of Functional Analysis* **9**(3), 306–321. [https://doi.org/10.1016/0022-1236\(72\)90004-3](https://doi.org/10.1016/0022-1236(72)90004-3)
- [24] Lamport, L., Shostak, R., and Pease, M. (1982). “The Byzantine generals problem.” *ACM Transactions on Programming Languages and Systems* **4**(3), 382–401. <https://doi.org/10.1145/357172.357176>
- [25] Fischer, M. J., Lynch, N. A., and Paterson, M. S. (1985). “Impossibility of distributed consensus with one faulty process.” *Journal of the ACM* **32**(2), 374–382. <https://doi.org/10.1145/3149.214121>
- [26] Newman, M. H. A. (1942). “On theories with a combinatorial definition of equivalence.” *Annals of Mathematics* **43**(2), 223–243. <https://doi.org/10.2307/1968867>
- [27] Rovelli, C. (1996). “Relational quantum mechanics.” *International Journal of Theoretical Physics* **35**, 1637–1678. <https://doi.org/10.1007/BF02302261>
- [28] Connes, A., and Rovelli, C. (1994). “Von Neumann algebra automorphisms and time-thermodynamics relation in generally covariant quantum theories.” *Classical and Quantum Gravity* **11**, 2899–2917. <https://doi.org/10.1088/0264-9381/11/12/007>
- [29] Zurek, W. H. (2003). “Decoherence, einselection, and the quantum origins of the classical.” *Reviews of Modern Physics* **75**, 715–775. <https://doi.org/10.1103/RevModPhys.75.715>
- [30] ’t Hooft, G. (2016). *The Cellular Automaton Interpretation of Quantum Mechanics*. Fundamental Theories of Physics 185. Springer, Cham. <https://doi.org/10.1007/978-3-319-41285-6>
- [31] Frauchiger, D., and Renner, R. (2018). “Quantum theory cannot consistently describe the use of itself.” *Nature Communications* **9**, 3711. <https://doi.org/10.1038/s41467-018-05739-8>
- [32] Ratcliffe, J. G. (2019). *Foundations of Hyperbolic Manifolds*, 3rd ed. Graduate Texts in Mathematics 149. Springer, Cham. <https://doi.org/10.1007/978-3-030-31597-9>
- [33] Serre, J.-P. (1977). *Linear Representations of Finite Groups*. Graduate Texts in Mathematics 42. Springer, New York. <https://doi.org/10.1007/978-1-4684-9458-7>

- [34] Cohan, N. V. (1958). “The spherical harmonics with the symmetry of the icosahedral group.” *Proceedings of the Cambridge Philosophical Society* **54**(1), 28–38. <https://doi.org/10.1017/S0305004100033156>
- [35] Bisognano, J. J., and Wichmann, E. H. (1975). “On the duality condition for a Hermitian scalar field.” *Journal of Mathematical Physics* **16**, 985–1007. <https://doi.org/10.1063/1.522605>
- [36] Bisognano, J. J., and Wichmann, E. H. (1976). “On the duality condition for quantum fields.” *Journal of Mathematical Physics* **17**, 303–321. <https://doi.org/10.1063/1.522898>
- [37] Wiesbrock, H.-W. (1993). “Half-sided modular inclusions of von Neumann algebras.” *Communications in Mathematical Physics* **157**, 83–92. <https://doi.org/10.1007/BF02098019>
- [38] Jacobson, T. (1995). “Thermodynamics of spacetime: The Einstein equation of state.” *Physical Review Letters* **75**, 1260–1263. <https://doi.org/10.1103/PhysRevLett.75.1260>
- [39] Jacobson, T. (2016). “Entanglement equilibrium and the Einstein equation.” *Physical Review Letters* **116**, 201101. <https://doi.org/10.1103/PhysRevLett.116.201101>
- [40] van der Bij, J. J., van Dam, H., and Ng, Y. J. (1982). “The exchange of massless spin-two particles.” *Physica A* **116**, 307–320. [https://doi.org/10.1016/0378-4371\(82\)90247-3](https://doi.org/10.1016/0378-4371(82)90247-3)
- [41] Ellis, G. F. R., van Elst, H., Murugan, J., and Uzan, J.-P. (2011). “On the trace-free Einstein equations as a viable alternative to general relativity.” *Classical and Quantum Gravity* **28**, 225007. <https://doi.org/10.1088/0264-9381/28/22/225007>
- [42] Van Raamsdonk, M. (2010). “Building up spacetime with quantum entanglement.” *General Relativity and Gravitation* **42**, 2323–2329. <https://doi.org/10.1007/s10714-010-1034-0>
- [43] Donnelly, W. (2012). “Decomposition of entanglement entropy in lattice gauge theory.” *Physical Review D* **85**, 085004. <https://doi.org/10.1103/PhysRevD.85.085004>
- [44] Casini, H., Huerta, M., and Rosabal, J. A. (2014). “Remarks on entanglement entropy for gauge fields.” *Physical Review D* **89**, 085012. <https://doi.org/10.1103/PhysRevD.89.085012>
- [45] Hall, B. C. (2015). *Lie Groups, Lie Algebras, and Representations: An Elementary Introduction*, 2nd ed. Graduate Texts in Mathematics 222. Springer, Cham. <https://doi.org/10.1007/978-3-319-13467-3>
- [46] Doplicher, S., and Roberts, J. E. (1989). “A new duality theory for compact groups.” *Inventiones Mathematicae* **98**, 157–218. <https://doi.org/10.1007/BF01388849>
- [47] Doplicher, S., and Roberts, J. E. (1990). “Why there is a field algebra with a compact gauge group describing the superselection structure in particle physics.” *Communications in Mathematical Physics* **131**, 51–107. <https://doi.org/10.1007/BF02097680>
- [48] Georgi, H., and Glashow, S. L. (1974). “Unity of all elementary-particle forces.” *Physical Review Letters* **32**(8), 438–441. <https://doi.org/10.1103/PhysRevLett.32.438>
- [49] Bouchiat, C., Iliopoulos, J., and Meyer, Ph. (1972). “An anomaly-free version of Weinberg’s model.” *Physics Letters B* **38**(7), 519–523. [https://doi.org/10.1016/0370-2693\(72\)90532-1](https://doi.org/10.1016/0370-2693(72)90532-1)

- [50] Tong, D. (2017). “Line operators in the Standard Model.” *Journal of High Energy Physics* **07**, 104. [https://doi.org/10.1007/JHEP07\(2017\)104](https://doi.org/10.1007/JHEP07(2017)104)
- [51] Witten, E. (1982). “An SU(2) anomaly.” *Physics Letters B* **117**, 324–328. [https://doi.org/10.1016/0370-2693\(82\)90728-6](https://doi.org/10.1016/0370-2693(82)90728-6)
- [52] Kostelecký, V. A., and Mewes, M. (2009). “Electrodynamics with Lorentz-violating operators of arbitrary dimension.” *Physical Review D* **80**, 015020. <https://doi.org/10.1103/PhysRevD.80.015020>
- [53] Kostelecký, V. A., and Russell, N. (2026). “Data tables for Lorentz and CPT violation.” 2026 edition, arXiv:0801.0287. <https://arxiv.org/abs/0801.0287>. Original journal article: <https://doi.org/10.1103/RevModPhys.83.11>
- [54] Xi, Y., and Shu, F.-W. (2025). “Constraints on Lorentz invariance violation from GRB 221009A using the DisCan method.” *Chinese Physics C* **49**, 125101. <https://doi.org/10.1088/1674-1137/adfa01>
- [55] LHAASO Collaboration (2024). “Stringent tests of Lorentz invariance violation from LHAASO observations of GRB 221009A.” *Physical Review Letters* **133**, 071501. <https://doi.org/10.1103/PhysRevLett.133.071501>
- [56] Mohr, P. J., Newell, D. B., Taylor, B. N., and Tiesinga, E. (2025). “CODATA recommended values of the fundamental physical constants: 2022.” *Reviews of Modern Physics* **97**, 025002. <https://doi.org/10.1103/RevModPhys.97.025002>
- [57] Takahashi, F., et al. (Particle Data Group) (2026). “Review of Particle Physics.” *International Journal of Modern Physics A* **41**, 2630011. <https://doi.org/10.1142/S0217751X26300115>. 2026 particle listings: [https://pdg.lbl.gov/2026/listings/particle\\_properties.html](https://pdg.lbl.gov/2026/listings/particle_properties.html)

## AI Assistance Disclosure

This research project used the latest research-grade commercial models, including Anthropic’s Fable and OpenAI’s GPT-5.6-Sol, for research support, software development, editing, and synthesis. The authors are responsible for the paper’s claims, methods, and final text.