

[PROBLEM P10]

Adherence and Complication Recognition During Post-Operative Recovery

01	PROBLEM SUMMARY
Patients may receive dense written guidance while tired, medicated, or anxious.	
02	CONTEXT & AFFECTED GROUPS
After discharge, patients and caregivers become responsible for wound care, medication, movement restrictions, diet, exercises, follow-up visits, and monitoring for complications. Instructions can vary by procedure and change across different stages of recovery.	
03	DETAILED PROBLEM SPECIFICATION
Patients may receive dense written guidance while tired, medicated, or anxious. Daily tasks can be forgotten, performed incorrectly, or stopped early when symptoms improve. Normal recovery symptoms may be difficult to distinguish from warning signs such as infection, blood clots, uncontrolled pain, bleeding, dehydration, or wound separation. Recovery plans also depend on age, existing conditions, caregiver availability, and access to transport or follow-up care.	
04	STAKES & CONSEQUENCES IF UNADDRESSED
Missed routines and delayed recognition of complications can result in preventable pain, slower healing, readmission, additional procedures, and higher costs. Care teams may receive incomplete or late information about changes occurring between scheduled appointments. EduTech	
05	KEY PROBLEM SCOPE & OBJECTIVES
- Patients may receive dense written guidance while tired, medicated, or anxious. - Daily tasks can be forgotten, performed incorrectly, or stopped early when symptoms improve. - Normal recovery symptoms may be difficult to distinguish from warning signs such as infection, blood clots, uncontrolled pain, bleeding, dehydration, or wound separation. - Recovery plans also depend on age, existing conditions, caregiver availability, and access to transport or follow-up care.	