

## Row Space, Column Space, and Null Space

### Matrix Spaces

#### Definition 1

For an  $m \times n$  matrix

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

the vectors

$$\begin{aligned} \mathbf{r}_1 &= [a_{11} \quad a_{12} \quad \cdots \quad a_{1n}] \\ \mathbf{r}_2 &= [a_{21} \quad a_{22} \quad \cdots \quad a_{2n}] \\ &\vdots \\ \mathbf{r}_m &= [a_{m1} \quad a_{m2} \quad \cdots \quad a_{mn}] \end{aligned}$$

in  $R^n$  formed from the rows of  $A$  are called the **row vectors** of  $A$ , and the vectors

$$\mathbf{c}_1 = \begin{bmatrix} a_{11} \\ a_{21} \\ \vdots \\ a_{m1} \end{bmatrix}, \quad \mathbf{c}_2 = \begin{bmatrix} a_{12} \\ a_{22} \\ \vdots \\ a_{m2} \end{bmatrix}, \dots, \quad \mathbf{c}_n = \begin{bmatrix} a_{1n} \\ a_{2n} \\ \vdots \\ a_{mn} \end{bmatrix}$$

in  $R^m$  formed from the columns of  $A$  are called the **column vectors** of  $A$ .

#### EXAMPLE 1 | Row and Column Vectors of a $2 \times 3$ Matrix

Let

$$A = \begin{bmatrix} 2 & 1 & 0 \\ 3 & -1 & 4 \end{bmatrix}$$

The row vectors of  $A$  are

$$\mathbf{r}_1 = [2 \quad 1 \quad 0] \quad \text{and} \quad \mathbf{r}_2 = [3 \quad -1 \quad 4]$$

and the column vectors of  $A$  are

$$\mathbf{c}_1 = \begin{bmatrix} 2 \\ 3 \end{bmatrix}, \quad \mathbf{c}_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}, \quad \text{and} \quad \mathbf{c}_3 = \begin{bmatrix} 0 \\ 4 \end{bmatrix}$$

## Row Space, Column Space

### Definition 2

If  $A$  is an  $m \times n$  matrix, then the subspace of  $R^n$  spanned by the row vectors of  $A$  is denoted by  $\text{row}(A)$  and is called the **row space** of  $A$ , and the subspace of  $R^m$  spanned by the column vectors of  $A$  is denoted by  $\text{col}(A)$  and is called the **column space** of  $A$ . The solution space of the homogeneous system of equations  $A\mathbf{x} = \mathbf{0}$ , which is a subspace of  $R^n$ , is denoted by  $\text{null}(A)$  and is called the **null space** of  $A$ .

### Theorem 4.8.1

A system of linear equations  $A\mathbf{x} = \mathbf{b}$  is consistent if and only if  $\mathbf{b}$  is in the column space of  $A$ .

### EXAMPLE 2 | A Vector $\mathbf{b}$ in the Column Space of $A$

Let  $A\mathbf{x} = \mathbf{b}$  be the linear system

$$\begin{bmatrix} -1 & 3 & 2 \\ 1 & 2 & -3 \\ 2 & 1 & -2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 1 \\ -9 \\ -3 \end{bmatrix}$$

Show that  $\mathbf{b}$  is in the column space of  $A$  by expressing it as a linear combination of the column vectors of  $A$ .

**Solution** Solving the system by Gaussian elimination yields (verify)

$$x_1 = 2, \quad x_2 = -1, \quad x_3 = 3$$

It follows from this and Formula (2) that

$$2 \begin{bmatrix} -1 \\ 1 \\ 2 \end{bmatrix} - \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix} + 3 \begin{bmatrix} 2 \\ -3 \\ -2 \end{bmatrix} = \begin{bmatrix} 1 \\ -9 \\ -3 \end{bmatrix}$$



## The Relationship Between $A\mathbf{x} = \mathbf{0}$ and $A\mathbf{x} = \mathbf{b}$

In this subsection we will explore the relationship between the solutions of a homogeneous linear system  $A\mathbf{x} = \mathbf{0}$  and the solutions (if any) of the nonhomogeneous linear system  $A\mathbf{x} = \mathbf{b}$  with the same coefficient matrix. These are called **corresponding linear systems**. By way of example, we will consider the following linear systems that we first discussed in Examples 5 and 6 of Section 1.2 and then again in Example 3 of Section 4.6.

$$\begin{bmatrix} 1 & 3 & -2 & 0 & 2 & 0 \\ 2 & 6 & -5 & -2 & 4 & -3 \\ 0 & 0 & 5 & 10 & 0 & 15 \\ 2 & 6 & 0 & 8 & 4 & 18 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \\ x_6 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} 1 & 3 & -2 & 0 & 2 & 0 \\ 2 & 6 & -5 & -2 & 4 & -3 \\ 0 & 0 & 5 & 10 & 0 & 15 \\ 2 & 6 & 0 & 8 & 4 & 18 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \\ x_6 \end{bmatrix} = \begin{bmatrix} 0 \\ -1 \\ 5 \\ 6 \end{bmatrix}$$

In Section 1.2 we found the general solutions of these systems to be

**homogeneous**  $\longrightarrow x_1 = -3r - 4s - 2t, \quad x_2 = r, \quad x_3 = -2s, \quad x_4 = s, \quad x_5 = t, \quad x_6 = 0$

**nonhomogeneous**  $\longrightarrow x_1 = -3r - 4s - 2t, \quad x_2 = r, \quad x_3 = -2s, \quad x_4 = s, \quad x_5 = t, \quad x_6 = \frac{1}{3}$

which we can express in column-vector form as

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \\ x_6 \end{bmatrix} = \begin{bmatrix} -3r - 4s - 2t \\ r \\ -2s \\ s \\ t \\ 0 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \\ x_6 \end{bmatrix} = \begin{bmatrix} -3r - 4s - 2t \\ r \\ -2s \\ s \\ t \\ \frac{1}{3} \end{bmatrix}$$



By splitting the entries on the right apart and collecting terms with like parameters we can rewrite these general solutions as

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \\ x_6 \end{bmatrix} = r \begin{bmatrix} -3 \\ 1 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} + s \begin{bmatrix} -4 \\ 0 \\ -2 \\ 1 \\ 0 \\ 0 \end{bmatrix} + t \begin{bmatrix} -2 \\ 0 \\ 0 \\ 0 \\ 1 \\ 0 \end{bmatrix} \quad (3)$$

Homogeneous Case

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \\ x_6 \end{bmatrix} = r \begin{bmatrix} -3 \\ 1 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} + s \begin{bmatrix} -4 \\ 0 \\ -2 \\ 1 \\ 0 \\ 0 \end{bmatrix} + t \begin{bmatrix} -2 \\ 0 \\ 0 \\ 0 \\ 1 \\ 0 \end{bmatrix} + \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ \frac{1}{3} \end{bmatrix} \quad (4)$$

Nonhomogeneous Case

In Example 3 of Section 4.6 we observed that the three vectors on the right side of (3) are linearly independent and therefore form a basis for the solution space of the homogeneous system. Thus, as illustrated in (5), the general solution  $\mathbf{x}$  of the nonhomogeneous system can be divided into two parts, a basis  $\mathbf{x}_h$  for the null space of the homogeneous system and a term  $\mathbf{x}_0$  that is a solution of the nonhomogeneous system (in this case, the solution resulting from setting the parameters to zero).

$$\underbrace{\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \\ x_6 \end{bmatrix}}_{\mathbf{x}} = \begin{bmatrix} -3r - 4s - 2t \\ r \\ -2s \\ s \\ t \\ \frac{1}{3} \end{bmatrix} = \underbrace{\begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ \frac{1}{3} \end{bmatrix}}_{\mathbf{x}_0} + r \underbrace{\begin{bmatrix} -3 \\ 1 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}}_{\mathbf{x}_h} + s \underbrace{\begin{bmatrix} -4 \\ 0 \\ -2 \\ 1 \\ 0 \\ 0 \end{bmatrix}}_{\mathbf{x}_h} + t \underbrace{\begin{bmatrix} -2 \\ 0 \\ 0 \\ 0 \\ 1 \\ 0 \end{bmatrix}}_{\mathbf{x}_h} \quad (5)$$

This example illustrates the following general theorem.

### Theorem 4.8.2

If  $\mathbf{x}_0$  is any solution of a consistent linear system  $A\mathbf{x} = \mathbf{b}$ , and if  $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\}$  is a basis for the null space of  $A$ , then every solution of  $A\mathbf{x} = \mathbf{b}$  can be expressed in the form


$$\mathbf{x} = \mathbf{x}_0 + c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_k\mathbf{v}_k \quad (6)$$

Conversely, for all choices of scalars  $c_1, c_2, \dots, c_k$ , the vector  $\mathbf{x}$  in this formula is a solution of  $A\mathbf{x} = \mathbf{b}$ .

The vector  $\mathbf{x}_0$  in Formula (6) is called a **particular solution of  $A\mathbf{x} = \mathbf{b}$** , and the remaining part of the formula is called the **general solution of  $A\mathbf{x} = \mathbf{0}$** . With this terminology Theorem 4.8.2 can be rephrased as:

*The general solution of a consistent linear system can be expressed as the sum of a particular solution of that system and the general solution of the corresponding homogeneous system.*

Geometrically, the solution set of  $A\mathbf{x} = \mathbf{b}$  can be viewed as the translation by  $\mathbf{x}_0$  of the solution space of  $A\mathbf{x} = \mathbf{0}$  (Figure 4.8.1).



**FIGURE 4.8.1** The solution space of  $A\mathbf{x} = \mathbf{b}$  is a translation of the solution space of  $A\mathbf{x} = \mathbf{0}$ .

## Bases for Row Spaces, Column Spaces, and Null Spaces

### Theorem 4.8.3

- (a) Row equivalent matrices have the same row space.
- (b) Row equivalent matrices have the same null space.

### Theorem 4.8.4

If a matrix  $R$  is in row echelon form, then the row vectors with the leading 1's (the nonzero row vectors) form a basis for the row space of  $R$ , and the column vectors with the leading 1's of the row vectors form a basis for the column space of  $R$ .



### EXAMPLE 3 | Bases for the Row and Column Spaces of a Matrix in Row Echelon Form

Find bases for the row and column spaces of the matrix

$$R = \begin{bmatrix} 1 & -2 & 5 & 0 & 3 \\ 0 & 1 & 3 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

**Solution** Since the matrix  $R$  is in row echelon form, it follows from Theorem 4.8.4 that the vectors

$$\mathbf{r}_1 = [1 \quad -2 \quad 5 \quad 0 \quad 3]$$

$$\mathbf{r}_2 = [0 \quad 1 \quad 3 \quad 0 \quad 0]$$

$$\mathbf{r}_3 = [0 \quad 0 \quad 0 \quad 1 \quad 0]$$

form a basis for the row space of  $R$ , and the vectors

$$\mathbf{c}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \quad \mathbf{c}_2 = \begin{bmatrix} -2 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \quad \mathbf{c}_4 = \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \\ 0 \end{bmatrix}$$

form a basis for the column space of  $R$ .

### EXAMPLE 4 | Basis for a Row Space by Row Reduction

Find a basis for the row space of the matrix

$$A = \begin{bmatrix} 1 & -3 & 4 & -2 & 5 & 4 \\ 2 & -6 & 9 & -1 & 8 & 2 \\ 2 & -6 & 9 & -1 & 9 & 7 \\ -1 & 3 & -4 & 2 & -5 & -4 \end{bmatrix}$$

**Solution** Since elementary row operations do not change the row space of a matrix, we can find a basis for the row space of  $A$  by finding a basis for the row space of any row echelon form of  $A$ . Reducing  $A$  to row echelon form, we obtain (verify)

$$R = \begin{bmatrix} 1 & -3 & 4 & -2 & 5 & 4 \\ 0 & 0 & 1 & 3 & -2 & -6 \\ 0 & 0 & 0 & 0 & 1 & 5 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

By Theorem 4.8.4, the nonzero row vectors of  $R$  form a basis for the row space of  $R$  and hence form a basis for the row space of  $A$ . These basis vectors are

$$\mathbf{r}_1 = [1 \quad -3 \quad 4 \quad -2 \quad 5 \quad 4]$$

$$\mathbf{r}_2 = [0 \quad 0 \quad 1 \quad 3 \quad -2 \quad -6]$$

$$\mathbf{r}_3 = [0 \quad 0 \quad 0 \quad 0 \quad 1 \quad 5]$$



### Theorem 4.8.5

If  $A$  and  $B$  are row equivalent matrices, then:

- (a) A given set of column vectors of  $A$  is linearly independent if and only if the corresponding column vectors of  $B$  are linearly independent.
- (b) A given set of column vectors of  $A$  forms a basis for the column space of  $A$  if and only if the corresponding column vectors of  $B$  form a basis for the column space of  $B$ .

### EXAMPLE 5 | Basis from the Columns of $A$

Find a basis for the column space of the matrix

$$A = \begin{bmatrix} 1 & -3 & 4 & -2 & 5 & 4 \\ 2 & -6 & 9 & -1 & 8 & 2 \\ 2 & -6 & 9 & -1 & 9 & 7 \\ -1 & 3 & -4 & 2 & -5 & -4 \end{bmatrix}$$

that consists of column vectors of  $A$ .

**Solution** We observed in Example 4 that the matrix

$$R = \begin{bmatrix} 1 & -3 & 4 & -2 & 5 & 4 \\ 0 & 0 & 1 & 3 & -2 & -6 \\ 0 & 0 & 0 & 0 & 1 & 5 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

is a row echelon form of  $A$ . Keeping in mind that  $A$  and  $R$  can have different column spaces, we cannot find a basis for the column space of  $A$  directly from the column vectors of  $R$ . However, it follows from Theorem 4.8.5(b) that if we can find a set of column vectors of  $R$  that forms a basis for the column space of  $R$ , then the *corresponding* column vectors of  $A$  will form a basis for the column space of  $A$ .

Since the first, third, and fifth columns of  $R$  contain the leading 1's of the row vectors, the vectors

$$\mathbf{c}'_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \quad \mathbf{c}'_3 = \begin{bmatrix} 4 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \quad \mathbf{c}'_5 = \begin{bmatrix} 5 \\ -2 \\ 1 \\ 0 \end{bmatrix}$$

form a basis for the column space of  $R$ . Thus, the corresponding column vectors of  $A$ , which are

$$\mathbf{c}_1 = \begin{bmatrix} 1 \\ 2 \\ 2 \\ -1 \end{bmatrix}, \quad \mathbf{c}_3 = \begin{bmatrix} 4 \\ 9 \\ 9 \\ -4 \end{bmatrix}, \quad \mathbf{c}_5 = \begin{bmatrix} 5 \\ 8 \\ 9 \\ -5 \end{bmatrix}$$

form a basis for the column space of  $A$ .

### EXAMPLE 6 | Basis from the Rows of $A$

Find a basis for the row space of

$$A = \begin{bmatrix} 1 & -2 & 0 & 0 & 3 \\ 2 & -5 & -3 & -2 & 6 \\ 0 & 5 & 15 & 10 & 0 \\ 2 & 6 & 18 & 8 & 6 \end{bmatrix}$$

consisting entirely of row vectors from  $A$ .

**Solution** We will transpose  $A$ , thereby converting the row space of  $A$  into the column space of  $A^T$ ; then we will use the method of Example 5 to find a basis for the column space of  $A^T$ ; and then we will transpose again to convert column vectors back to row vectors.

Transposing  $A$  yields

$$A^T = \begin{bmatrix} 1 & 2 & 0 & 2 \\ -2 & -5 & 5 & 6 \\ 0 & -3 & 15 & 18 \\ 0 & -2 & 10 & 8 \\ 3 & 6 & 0 & 6 \end{bmatrix}$$

and then reducing this matrix to row echelon form we obtain

$$\begin{bmatrix} 1 & 2 & 0 & 2 \\ 0 & 1 & -5 & -10 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

The first, second, and fourth columns contain the leading 1's, so the corresponding column vectors in  $A^T$  form a basis for the column space of  $A^T$ ; these are

$$\mathbf{c}_1 = \begin{bmatrix} 1 \\ -2 \\ 0 \\ 0 \\ 3 \end{bmatrix}, \quad \mathbf{c}_2 = \begin{bmatrix} 2 \\ -5 \\ -3 \\ -2 \\ 6 \end{bmatrix}, \quad \text{and} \quad \mathbf{c}_4 = \begin{bmatrix} 2 \\ 6 \\ 18 \\ 8 \\ 6 \end{bmatrix}$$

Transposing again and adjusting the notation appropriately yields the basis vectors

$$\mathbf{r}_1 = [1 \quad -2 \quad 0 \quad 0 \quad 3], \quad \mathbf{r}_2 = [2 \quad -5 \quad -3 \quad -2 \quad 6], \\ \mathbf{r}_4 = [2 \quad 6 \quad 18 \quad 8 \quad 6]$$

for the row space of  $A$ .



## EXAMPLE 8 | Basis and Linear Combinations

(a) Find a subset of the vectors

$$\begin{aligned}\mathbf{v}_1 &= (1, -2, 0, 3), & \mathbf{v}_2 &= (2, -5, -3, 6), \\ \mathbf{v}_3 &= (0, 1, 3, 0), & \mathbf{v}_4 &= (2, -1, 4, -7), & \mathbf{v}_5 &= (5, -8, 1, 2)\end{aligned}$$

that forms a basis for the subspace of  $\mathbb{R}^4$  spanned by these vectors.

(b) Express each vector not in the basis as a linear combination of the basis vectors.

**Solution (a)** We begin by constructing a matrix that has  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_5$  as its column vectors:

$$\begin{bmatrix} 1 & 2 & 0 & 2 & 5 \\ -2 & -5 & 1 & -1 & -8 \\ 0 & -3 & 3 & 4 & 1 \\ 3 & 6 & 0 & -7 & 2 \end{bmatrix} \quad (9)$$

$\uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow$   
 $\mathbf{v}_1 \quad \mathbf{v}_2 \quad \mathbf{v}_3 \quad \mathbf{v}_4 \quad \mathbf{v}_5$

The first part of our problem can be solved by finding a basis for the column space of this matrix. Reducing the matrix to *reduced* row echelon form and denoting the column vectors of the resulting matrix by  $\mathbf{w}_1, \mathbf{w}_2, \mathbf{w}_3, \mathbf{w}_4$ , and  $\mathbf{w}_5$  yields

$$\begin{bmatrix} 1 & 0 & 2 & 0 & 1 \\ 0 & 1 & -1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \quad (10)$$

$\uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow$   
 $\mathbf{w}_1 \quad \mathbf{w}_2 \quad \mathbf{w}_3 \quad \mathbf{w}_4 \quad \mathbf{w}_5$

The leading 1's occur in columns 1, 2, and 4, so by Theorem 4.8.4,

$$\{\mathbf{w}_1, \mathbf{w}_2, \mathbf{w}_4\}$$

is a basis for the column space of (6), and consequently,

$$\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_4\}$$

is a basis for the column space of (9).

**Solution (b)** We will start by expressing  $\mathbf{w}_3$  and  $\mathbf{w}_5$  as linear combinations of the basis vectors  $\mathbf{w}_1, \mathbf{w}_2, \mathbf{w}_4$ . The simplest way of doing this is to express  $\mathbf{w}_3$  and  $\mathbf{w}_5$  in terms of basis vectors with numerically smaller subscripts. Accordingly, we will express  $\mathbf{w}_3$  as a linear combination of  $\mathbf{w}_1$  and  $\mathbf{w}_2$ , and we will express  $\mathbf{w}_5$  as a linear combination of the vectors  $\mathbf{w}_1, \mathbf{w}_2$ , and  $\mathbf{w}_4$ . By inspection of (10), these linear combinations are

$$\begin{aligned}\mathbf{w}_3 &= 2\mathbf{w}_1 - \mathbf{w}_2 \\ \mathbf{w}_5 &= \mathbf{w}_1 + \mathbf{w}_2 + \mathbf{w}_4\end{aligned}$$

We call these the **dependency equations**. The corresponding relationships in (9) are

$$\begin{aligned}\mathbf{v}_3 &= 2\mathbf{v}_1 - \mathbf{v}_2 \\ \mathbf{v}_5 &= \mathbf{v}_1 + \mathbf{v}_2 + \mathbf{v}_4\end{aligned}$$