



< All iCloud

Ad



Polar co-ordinates in Double integration :-

$$A = \frac{1}{2} \int_{\theta_1}^{\theta_2} r^2 d\theta$$



Polar rectangle.

$$\iint_R f(x, y) dA$$

$$x = r \cos \theta$$

$$y = r \sin \theta$$

$$x^2 + y^2 = r^2$$

$$\tan \theta = \frac{y}{x}$$

$$dA = r dr d\theta$$

< All iCloud

Ad



Figure ①

∴ always take the constant as outer limit

∴ across the region along the radius

$$\int_{\theta=\alpha}^{\theta=\beta} \int_{r=a}^{r=b} f(r \cos \theta, r \sin \theta) \, dr \, d\theta$$

Figure ②


$$\int_{\theta=\alpha}^{\theta=\beta} \int_{r=g_1(\theta)}^{r=g_2(\theta)} f(r \cos \theta, r \sin \theta) \, r \, dr \, d\theta$$

$$Q_9 = \iint_R xy \, dA$$
, R : The region in Q_1 bound
 by $x^2 + y^2 = 4$, $x = y$ & $x = 0$

Solⁿ

$$x^2 = 4$$

$$x = 2$$



$$\theta = \tan^{-1}\left(\frac{y}{x}\right)$$

$$\theta = \tan^{-1}(1)$$

$$\theta = \pi/4$$

$$\iint_{\substack{\theta: \pi/4 \quad r=0 \\ \theta: \pi/2 \quad r=2}} r \cos \theta \cdot r \sin \theta \, r \, dr \, d\theta$$

$$\int_{\pi/4}^{\pi/2} \int_0^2 r^3 \cos \theta \sin \theta \, dr \, d\theta$$

All iCloud

Ad



$$\int_{\pi/4}^{\pi/2} \left[\frac{r^4}{4} \cos \theta \sin \theta \right]_0^2 d\theta$$

$$\int_{\pi/4}^{\pi/2} \left[\frac{16}{4} \cos \theta \sin \theta \right] - 0 d\theta$$

$$4 \int_{\pi/4}^{\pi/2} \cos \theta \sin \theta d\theta$$

$$4 \left[\frac{\sin^2 \theta}{2} \right]_{\pi/4}^{\pi/2}$$

$$4 \left[\frac{\sin^2(\pi/2)}{2} - \frac{\sin^2(\pi/4)}{2} \right]$$

$$4 \left[\frac{1}{2} - \frac{1}{4} \right]$$

$$4 \left[\frac{1}{4} \right] = \textcircled{1}$$

ANS

$$\therefore \int f(x) \cdot f'(x) dx = \left[\frac{f(x)^{n+1}}{n+1} \right] + C$$

< All iCloud

Ad




Q) Find volume between $z = 9 - x^2 - y^2$ & the xy -plane as bounded by the cylinder

$$x^2 + y^2 = 1$$



$$r^2 = 1$$

$$r = 1$$



$$\int_0^{2\pi} \int_0^1 (9 - r^2) r \, dr \, d\theta$$

$$\int_0^{2\pi} \int_0^1 (9r - r^3) \, dr \, d\theta$$


$$\int_0^{2\pi} \left[\frac{9r^2}{2} - \frac{r^4}{4} \right]_0^1 d\theta$$

< All iCloud

Aa



Q) Volume b/w $z = \frac{y^2}{x^2 + y^2}$, above xy -plane
 & b/w cylinders $x^2 + y^2 = 1$ & $x^2 + y^2 = 2$



$$\int_{\theta=0}^{2\pi} \int_{r=1}^{\sqrt{2}} \frac{\sin^2 \theta}{r^2} \cdot r \, dr \, d\theta$$

$$\int_0^{2\pi} \int_1^{\sqrt{2}} \sin^2 \theta \, r \, dr \, d\theta$$

Start from
 origin, as the
 arrow touches
 the shaded
 region
 that will
 become
 our
 initial
 r

< All iCloud

Ad



$$\int_0^{2\pi} \sin^2 \theta \left[\frac{r^2}{2} \right]_1^{\sqrt{2}} d\theta$$

$$\int_0^{2\pi} \sin^2 \theta \left[\frac{2}{2} - \frac{1}{2} \right] d\theta$$

$$\frac{1}{2} \int_0^{2\pi} \sin^2 \theta d\theta$$

$$\frac{1}{2} \int_0^{2\pi} \frac{1 - \cos 2\theta}{2} d\theta$$

$$\frac{1}{4} \left[\theta - \frac{\sin 2\theta}{2} \right]_0^{2\pi}$$

$$\frac{1}{4} \left[(2\pi - 0) - (0 - 0) \right]$$

$$\frac{2\pi}{4}$$

=

$$\pi/2$$

ANS

< All iCloud

Ad



$$\equiv \int \int_R y \, dA, \quad R: \text{The region bound by}$$

$$x^2 + y^2 = 2x \quad \& \quad y = x$$

$$x^2 - 2x + y^2 = 0$$

$$(x-1)^2 - 1 + y^2 = 0$$


$$(x-1)^2 + y^2 = 1$$

$$y \leq x$$

$$x^2 + y^2 = 2x$$

$$r^2 = 2r \cos \theta$$

$$r = 2 \cos \theta$$



$$\theta = \pi/2 \quad r = 2 \cos \theta$$

$$\int \int r \sin \theta \, r \, dr \, d\theta$$

$$\theta = \pi/4 \quad \theta = 0$$

$$2 \cos \theta$$

$$\pi/2$$

$$\int \sin \theta \, r^2 \, dr \, d\theta$$

$$\text{Ex: } x^2 + y^2 = 2x$$

we need to
convert it in
the form of
polar to
make it
in the integral

< All iCloud

Ad



Ex 2 Set the region

$$R: x^2 + y^2 = 4$$

$$x^2 + y^2 = 2y \text{ in Quad I}$$


$$r = 2$$

$$x^2 + y^2 - 2y = 0$$

$$x^2 + (y-1) = 1$$

$$r^2 = 2r \sin \theta$$

$$r = 2 \sin \theta$$



$$\theta = \pi/2 \quad r = 2$$

$$\theta = 0$$

$$r = 2 \sin \theta$$

$$f(r \cos \theta, r \sin \theta) \, r \, dr \, d\theta$$

AM

< All iCloud

Ad



$$\theta = \pi/4$$

$$\theta = \pi/2$$

Limits:

$$\theta = \pi/4 \quad u = 1/\sqrt{2}$$

$$\theta = \pi/2 \quad u = 0$$

$$\pi/2 > \pi/4$$

$$+ \frac{8}{3} \int_0^{1/\sqrt{2}} u^3 + du$$

$$\frac{8}{3} \left[\frac{u^4}{4} \right]_0^{1/\sqrt{2}}$$

$$\frac{8}{3} \left[\frac{1}{16} \right]$$

$$\frac{1}{6}$$

Ans

< All iCloud

Ad



$$\theta = \pi/4$$

$$2\cos\theta$$

$$\int_{\pi/4}^{\pi/2} \int_0^{2\cos\theta} \sin\theta \cdot r^2 dr d\theta$$

$$\int_{\pi/4}^{\pi/2} \sin\theta \left[\frac{r^3}{3} \right]_0^{2\cos\theta} d\theta$$

$$\int_{\pi/4}^{\pi/2} \sin\theta \left[\frac{8\cos^3\theta}{3} - 0 \right] d\theta$$

$$\frac{8}{3} \int_{\pi/4}^{\pi/2} \sin\theta \cos^3\theta d\theta$$

$$\text{let } u = \cos\theta$$

$$du = -\sin\theta d\theta$$

$$\sin\theta d\theta = -du$$

$$\text{Ex: } x^2 + y^2 = 2x$$

we need to
convert it in
the form of
polar to
make it
in the integral

$$x^2 + y^2 = 2x \quad y^2 = 2x - x^2$$

$$x^2 + y^2 = 2x$$

$$2x^2 = 2x$$

$$x^2 - x = 0$$

$$x = 1$$

$$y = 1$$

$$(1, 1)$$

$$\tan\theta = \frac{y}{x}$$

$$\tan\theta = \frac{1}{1}$$

$$x = 0$$

$$y = 0$$

$$(0, 0)$$

$$\tan\theta = \frac{y}{x}$$

$$\tan\theta = \frac{0}{0}$$

< All iCloud

Ad




Intersection

$$4 = x^2 + y^2$$

$$\downarrow$$

$$4 = r^2$$

$$r = 2$$



$$V = \int \int_R f \, dA$$

$$8 - 2x^2 - 2y^2$$

$$\downarrow$$

$$\vdots$$

$$\text{top}$$

$$\vdots$$

$$\text{bottom}$$

$$= \begin{matrix} 8 - 2(x^2 + y^2) \\ 8 - 2r^2 \end{matrix}$$

$$V = \int_{\theta=0}^{2\pi} \int_{r=0}^2 (8 - 2r^2) r \, dr \, d\theta$$

< All iCloud

Ad



(\hat{n}) b/c
when $\theta = 0$
and $r = 2$

but when
 $\theta = \hat{n}$
 $r = 0$

$r = -2$

and in polar
when r is -ve
we flip it
means r becomes
positive

Full
rotation
 $0 \rightarrow \hat{n}$

$r = 2$

(Top function - Bottom function)

These two will intersect
and it gives
the level
curve

$\frac{z}{2}$

Volume

b/w

$$z = 9 - 2x^2 - 2y^2$$

$$z = 1$$

Bottom

Top

$$9 - 2x^2 - 2y^2 = 1$$

$$8 = 2(x^2 + y^2)$$

$$4 = x^2 + y^2$$

Intersection

< All iCloud

Ad



$$z = 12 - 3x - 4y$$

$$f(x,y) = 12 - 3x - 4y$$

eg Volume below $3x + 4y + z = 12$ bound by
 z region b/w $x^2 + y^2 = 2x$ & above xy -plane


polar

$$r^2 = 2r \cos \theta$$

$$r = 2 \cos \theta$$

$$x^2 + 2x + y^2 = 0$$

$$(x+1)^2 + y^2 = 1$$



$$\iint_R f(x,y) dA = \int_{\theta=0}^{\pi} \int_{r=0}^{r=2 \cos \theta} (12 - 3r \cos \theta - 4r \sin \theta) r dr d\theta$$

< All iCloud

Add



$$V = \int_0^{2\pi} \int_0^1 r \sqrt{2-r^2} - r^2 dr d\theta$$

$$V = \int_0^{2\pi} \int_0^1 r \sqrt{2-r^2} dr d\theta - \int_0^{2\pi} \int_0^1 r^2 dr d\theta$$

↓

$$\text{let } u = 2-r^2$$
$$\frac{du}{dr} = -2r$$
$$\frac{du}{-2} = r dr$$

$$r=0 \rightarrow u=2$$
$$r=1 \rightarrow u=1$$

< All iCloud

Aa

Q.5
sol

volume

b/w

$$x^2 + y^2 + z^2 = 2 \quad \& \quad z = \sqrt{x^2 + y^2}$$

$$z = \sqrt{2 - x^2 - y^2}$$

$$z = \sqrt{x^2 + y^2}$$

$$\sqrt{2 - x^2 - y^2} = \sqrt{x^2 + y^2}$$


$$2 - x^2 - y^2 = x^2 + y^2$$

$$2 = 2x^2 + 2y^2$$

$$1 = x^2 + y^2$$

intersection:

$$\boxed{x^2 + y^2 = 1}$$

1. let $D(x, y)$

↳ to divide top & bottom

$$\sqrt{2 - r^2} - \sqrt{r^2}$$

$$\sqrt{2 - r^2} - r$$

< All iCloud

Ad



$$V = \int_0^{2\pi} \left[\frac{8r^2}{2} - \frac{2r^4}{4} \right]_0^2 d\theta$$

$$V = \int_0^{2\pi} \left(4(2)^2 - \frac{2(2)^4}{4} \right) d\theta$$

$$V = \int_0^{2\pi} (16 - 8) d\theta$$

$$V = 8 \int_0^{2\pi} d\theta \quad \Rightarrow \quad 8 \left[\theta \right]_0^{2\pi}$$


$$8 [2\pi - 0] = \boxed{16\pi} \quad \text{ANS}$$

< All iCloud

Ad



Eg find Area of 'R' where 'R' is the
region bound by $r = 3 \cos \theta$



$$A = \int \int_R 1 \, dA$$

$$A = \int_{\theta=0}^{\theta=\pi/2} \int_{r=0}^{r=3\cos\theta} r \, dr \, d\theta$$

$$\Rightarrow \frac{9\pi}{4}$$

< All iCloud

Ad



$$V = 4\pi \left(\frac{\sqrt{2} - 1}{3} \right)$$

ANS
7

(x)

$$V = \iint_R f(x,y) dA$$

 $f(x,y) = \text{Height}$

$$V = A \cdot H$$

$$A = \frac{V}{H}$$

\therefore volume is
equal to
Area when
height is
equal to
one

$$A = \frac{\iint_R f(x,y) dA}{f(x,y)}$$

$$A = \iint_R dA$$

\rightarrow Give area of
a Region: R

< All iCloud

Ad



$$V = \int_0^{2\pi} \left(\int_1^2 \sqrt{4} \left(1 + \frac{du}{2} \right) d\theta \right)$$

$$V = \int_0^{2\pi} \left(\int_1^2 u^{1/2} du d\theta \right)$$

$$V = \frac{1}{2} \int_0^{2\pi} \left[\frac{2}{3} u^{3/2} \right]_1^2 d\theta$$

$$V = \frac{1}{3} \int_0^{2\pi} (2^{3/2} - 1) d\theta$$

$$V = \frac{1}{3} [2\sqrt{2} - 1] [\theta]_0^{2\pi}$$

$$V = \frac{1}{3} [2\sqrt{2} - 1] [2\pi]$$

$$V = \frac{4\sqrt{2}}{3} \pi - \frac{2\pi}{3} = \frac{2\pi}{3} (2\sqrt{2} - 1)$$

$$= \int_0^{2\pi} \left[\frac{2^3}{3} \right]_0^1 d\theta$$

$$= \int_0^{2\pi} \frac{1}{3} d\theta$$

$$= \frac{1}{3} (2\pi)$$

$$= \frac{2\pi}{3}$$

< All iCloud

Ad

eg

$$\int_{-1}^{+1} \frac{\sqrt{1-y^2}}{1+x^2+y^2} dx dy$$

right side
semi
circle

$$x=0$$


$$x = \sqrt{1-y^2}$$

$$y = -1$$

$$y = 1$$

$$x^2 + y^2 = 1$$

$$r=1$$



$$\theta = \pi/2$$

$$r = 1$$

$$\int_{\theta = -\pi/2}^{\pi/2} \int_{r=0}^1$$

$$\frac{1}{1+r^2}$$

$$r dr d\theta$$

< All iCloud


Aa



cardioid.

circle

Q) Find Area of the region bound by
 $r = 2$ & $r = 2(1 + \cos \theta)$ on QUAD-I



< All iCloud

Ad



\therefore The θ range from 0 to $\pi/2$ b/c if we
 get our range from 0 to 2π which means
 the upper positive area will cancel
 the lower negative area which results
 in total area to be zero so for

avoiding this we find the area of our
 upper circle by θ range: 0 to $\pi/2$
 and double the answer b/c both
 are symmetric.

when ever you are given a region
 in the polar form like: $x = r \cos \theta$
 or $y = r \sin \theta$

< All iCloud

Ad



$$\int_{-\pi/2}^{\pi/2} \frac{1}{2} [\ln a - 0] d\theta$$

$$\frac{\ln a}{2} \int_{-\pi/2}^{\pi/2} d\theta$$

$$\frac{\ln a}{2} \left[\theta \right]_{-\pi/2}^{\pi/2}$$

$$\frac{\ln 2}{2} \left[\pi/2 - (-\pi/2) \right] = \frac{\ln 2}{2} [\pi]$$

$$\frac{\pi \ln 2}{2}$$

Ans

< All iCloud

Aa



$$-\pi/2$$

let

$$u = 1 + r^2$$

$$\frac{du}{dr} = 2r$$

$$du/2 = r dr$$

$$r=0$$

$$u=1$$

$$r=1$$

$$u=2$$

$$\int_{-\pi/2}^{\pi/2} \int_1^2$$

$$\frac{du/2}{u} d\theta$$

$$\int_{-\pi/2}^{\pi/2} \frac{1}{2} \left[\ln u \right]_1^2 d\theta$$

$$\int_{-\pi/2}^{\pi/2} \frac{1}{2} [\ln 2 - 0] d\theta$$

< All iCloud

Ad

 $2\pi/2$ to $\pi/2$ ↑
wrongb/c
we measure
angle only in
clockwise direction. (θ)
↓

$$y = 1 \rightarrow y = r \sin \theta$$

$$1 = r \sin \theta$$

$$\therefore r = 1$$

$$\sin \theta = 1 \Rightarrow \theta = \pi/2$$

$$y = -1 \rightarrow y = r \sin \theta$$

$$-1 = r \sin \theta$$

$$\theta = \frac{-1}{1} = -\pi/2 \quad \text{or } 3\pi/2$$

 $\int_{-\pi/2}^{\pi/2}$ \int_0^1

$$\frac{r}{1+r^2}$$

or \cos

< All iCloud

Ad



$$\int_0^{\pi/4} 2 \sec \theta \, d\theta$$

$$2 \left[\ln | \sec \theta - \tan \theta | \right]_0^{\pi/4}$$

$$2 \left[\ln | \sqrt{2} - 1 | - \ln | 1 - 0 | \right]$$

$$2 \left[\ln (\sqrt{2} - 1) - 0 \right]$$

$$\boxed{2 \ln (\sqrt{2} - 1)}$$

Ans

< All iCloud

Ad



$$\int_{\theta=0}^{\theta=\pi/4} \int_{r=\sec\theta}^{r=3\sec\theta} \frac{1}{\sqrt{r^2}} r dr d\theta$$

$$\text{col} = \int_0^{\pi/4} \int_{\sec\theta}^{3\sec\theta} \frac{1}{r} r dr d\theta$$

$$\int_0^{\pi/4} \left[r \right]_{\sec\theta}^{3\sec\theta} d\theta$$


$$\int_0^{\pi/4} (3\sec\theta - \sec\theta) d\theta$$

< All iCloud

Ad



Q) $\int_1^3 \int_0^x \frac{1}{\sqrt{x^2+y^2}} dy dx$



$$x \cos \theta = 3$$

$$x = \frac{3}{\cos \theta} = 3 \sec \theta$$

$$x = 1$$

$$x \cos \theta = 1$$

$$x = \frac{1}{\cos \theta}$$

$$x = \sec \theta$$

$$y = x \Rightarrow \tan(\theta) = \theta \quad \tan 1 = \theta$$

$$\theta = \tan^{-1} 1$$

$$y = 0 \Rightarrow \tan(\theta) = 0 \rightarrow \theta = 0^\circ$$