



Teiji Kunihiro
Yuta Kikuchi
Kyosuke Tsumura

Geometrical Formulation of Renormalization-Group Method as an Asymptotic Analysis

With Applications to Derivation
of Causal Fluid Dynamics



Proper length of the identical bodies

$$l = \frac{PP'}{OC} = \frac{P'P'}{OC}$$

Minkowski showed that:



Springer

Fundamental Theories of Physics

Volume 206

Series Editors

Henk van Beijeren, Utrecht, The Netherlands

Philippe Blanchard, Bielefeld, Germany

Bob Coecke, Oxford, UK

Dennis Dieks, Utrecht, The Netherlands

Bianca Dittrich, Waterloo, ON, Canada

Ruth Durrer, Geneva, Switzerland

Roman Frigg, London, UK

Christopher Fuchs, Boston, MA, USA

Domenico J. W. Giulini, Hanover, Germany

Gregg Jaeger, Boston, MA, USA

Claus Kiefer, Cologne, Germany

Nicolaas P. Landsman, Nijmegen, The Netherlands

Christian Maes, Leuven, Belgium

Mio Murao, Tokyo, Japan

Hermann Nicolai, Potsdam, Germany

Vesselin Petkov, Montreal, QC, Canada

Laura Ruetsche, Ann Arbor, MI, USA

Mairi Sakellariadou, London, UK

Alwyn van der Merwe, Greenwood Village, CO, USA

Rainer Verch, Leipzig, Germany

Reinhard F. Werner, Hanover, Germany

Christian Wüthrich, Geneva, Switzerland

Lai-Sang Young, New York City, NY, USA

The international monograph series “Fundamental Theories of Physics” aims to stretch the boundaries of mainstream physics by clarifying and developing the theoretical and conceptual framework of physics and by applying it to a wide range of interdisciplinary scientific fields. Original contributions in well-established fields such as Quantum Physics, Relativity Theory, Cosmology, Quantum Field Theory, Statistical Mechanics and Nonlinear Dynamics are welcome. The series also provides a forum for non-conventional approaches to these fields. Publications should present new and promising ideas, with prospects for their further development, and carefully show how they connect to conventional views of the topic. Although the aim of this series is to go beyond established mainstream physics, a high profile and open-minded Editorial Board will evaluate all contributions carefully to ensure a high scientific standard.

More information about this series at <https://link.springer.com/bookseries/6001>

Teiji Kunihiro · Yuta Kikuchi · Kyosuke Tsumura

Geometrical Formulation of Renormalization-Group Method as an Asymptotic Analysis

With Applications to Derivation of Causal
Fluid Dynamics

Teiji Kunihiro
Yukawa Institute for Theoretical Physics
Kyoto University
Sakyo-ku, Kyoto, Japan

Yuta Kikuchi
Brookhaven National Laboratory
Upton, NY, USA

Kyosuke Tsumura
Analysis Technology Center
Fujifilm Corporation
Minamiashigara, Kanagawa, Japan

ISSN 0168-1222

ISSN 2365-6425 (electronic)

Fundamental Theories of Physics

ISBN 978-981-16-8188-2

ISBN 978-981-16-8189-9 (eBook)

<https://doi.org/10.1007/978-981-16-8189-9>

© Springer Nature Singapore Pte Ltd. 2022

This work is subject to copyright. All rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

The publisher, the authors and the editors are safe to assume that the advice and information in this book are believed to be true and accurate at the date of publication. Neither the publisher nor the authors or the editors give a warranty, expressed or implied, with respect to the material contained herein or for any errors or omissions that may have been made. The publisher remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

This Springer imprint is published by the registered company Springer Nature Singapore Pte Ltd.

The registered company address is: 152 Beach Road, #21-01/04 Gateway East, Singapore 189721, Singapore

Preface

The purpose of this monograph is twofold: The one is to present a comprehensive account of the so-called Renormalization-Group (RG) method and its extension, which we named the doublet scheme, in a geometrical point of view with various examples. The second is the application of the doublet scheme in the RG method to the derivation of the so-called second-order (causal) fluid dynamics from the relativistic as well as non-relativistic Boltzmann equations with quantum statistics. All the contents of the monograph are virtually based on the authors original work except for some review parts.

The RG method is a global and asymptotic analysis of differential equations, which was developed by an Illinois group and others some thirty years ago. We introduce the method in a purely mathematical way on the basis of the classical theory of envelopes, a notion in elementary differential geometry.

Then a focus is put on the fact that the RG method provides us with a powerful and transparent method for the reduction of dynamics, which includes an elementary method of a construction of the invariant/attractive manifolds and reduced dynamical equations written in terms of the variables that constitute the coordinates of the manifolds. They are the key concepts in the reduction theory of dynamical systems, and thereby naturally lead to a foundation to existing theories in the specific physical systems such as Krylov-Bogoliubov-Mitropolsky theory for non-linear oscillators and Kuramoto's reduction theory for evolution equations. Examples treated include stochastic equations like Langevin and Fokker-Planck equations.

Although the RG method is applicable to discrete systems and thereby provides us with an optimized discretization scheme of differential equations, we have omitted the once prepared part on the discrete systems partly because of its irrelevance to the second part and partly for the sake of making the already lengthy monograph too voluminous. We hope that there will be a chance to publish the omitted part somewhere else in the future.

The usual reduction theory including the RG method based on the perturbation theory utilizes the zero modes of the linear operator in the unperturbed equation. However, in the derivation of the so-called causal second-order fluid dynamics, which has a nature of the mesoscopic dynamics of the given system, from the Boltzmann

equation, one needs to extend the invariant manifold so as to incorporate appropriate excited modes as the additional coordinate variables. In this monograph, a general reduction theory is presented for constructing the mesoscopic dynamics with inclusion of appropriate excited modes from the given microscopic equation, which is formulated as an extension of the RG method and called the doublet scheme.

In the second part, we work out for the derivation of the second-order or causal fluid dynamics on the basis of the doublet scheme: Thus we obtain not only the fluid dynamic equations that are uniquely those in the energy frame but also the microscopic formulae of the transport coefficients and the relaxation times in Kubo-like formulae that admit natural physical interpretations. It is shown that the resultant fluid dynamics is not only causal but also stable dissipative (non-)relativistic fluid dynamics. The derivation of the mesoscopic dynamics beyond the fluid dynamics in the non-relativistic regime is also one of the hot topics in physics of cold atoms. The present monograph includes numerical analyses on these interesting physical systems. We also provide an accurate and efficient numerical method for computing the transport equations and relaxation times using the microscopic expressions. The numerical method utilizes the double exponential formulae for integrations and the direct matrix inversion method. The numerical calculations are fully worked out for typical model systems composed of classical particles, a fermion system with a Yukawa interaction, and boson system described by a chiral Lagrangian. The numerical calculations are also presented for the non-relativistic system and a critical comparison is made with the relaxation-time approximation, which is commonly used in the current literature.

The presentation of the monograph, at least in the first part, is intended to be as pedagogical as possible so that not only researchers who are not familiar with the RG theory in physics but also motivated undergraduate students with mathematical backgrounds such as introductory calculus including linear differential equations and linear algebra may appreciate and understand the method.

Conversely, the monograph is not intended to be a systematic review of the current status of the studies of the RG method and causal fluid dynamics either with respect to its foundations or applications. Since the literature on these subjects is huge now that making a systematic review on this subject is not the intention of the authors and beyond the authors' ability. Therefore we apologize to those whose important articles are not cited in this monograph in advance.

Sakyo-ku, Kyoto, Japan
 Upton, NY, USA
 Minamiashigara, Kanagawa, Japan
 June 2021

Teiji Kunihiro
 Yuta Kikuchi
 Kyosuke Tsumura

Acknowledgements

One of the authors (T.K.) is grateful to Glenn Paquette who gave a nice seminar talk on the Illinois Renormalization-Group (RG) method at the Department of Applied Mathematics and Informatics, Ryukoku University, in the end of 1994, which was presented in a pedagogical way mostly with the use of elementary examples; however, honestly speaking, virtually all the audience of the seminar, the major part of which consisted of experts of differential equations/dynamical systems, did not understand what the lecturer was doing. This ‘extraordinary’ phenomenon may have reflected the fact that the content of the seminar, *i.e.*, the RG method, was so original and new. In any event, the seminar and the subsequent conversation with a couple of other attendants of the seminar strongly motivated T.K. to make an effort to understand what is being done in a purely mathematical way in the Illinois RG method.

He is indebted to Shoji Yotsutani for his question made to him, which prompted him to realize the significance of the ‘initial’ condition in the formulation of the RG method in a purely mathematical manner. He is also deeply indebted to late Professor Masaya Yamaguti for his interest in his work on the envelope formulation of the RG method and encouragement.

T.K. acknowledges Junta Matsukidaira and Shin-Ichiro Ei and Kazuyuki Fujii and Yoshitaka Hatta for their collaboration on the works on which this monograph is partly based.

We thank Steven Strogatz for his interest in this project to writing this monograph and elucidating questions/comments on a part of the manuscript in the final stage.

Finally, but not least, we are grateful to Yoshiki Kuramoto, Takao Ohta, and Alexander Gorban for their interest in our work on the RG method and encouragement.

Contents

1	Introduction: Reduction of Dynamics, Notion of Effective Theories, and Renormalization Groups	1
1.1	Reduction of Dynamics of a Simple Equation and the Notion of Effective Theory	1
1.2	Notion of Effective Theories and Renormalization Group in Physical Sciences	3
1.3	The Renormalization Group Method in Global and Asymptotic Analysis	4
1.4	Derivation of Stochastic Equations and Fluid Dynamic Limit of Boltzmann Equation	6
Part I	Geometrical Formulation of Renormalization-Group Method and Its Extention for Global and Asymptotic Analysis with Examples	
2	Naïve Perturbation Method for Solving Ordinary Differential Equations and Notion of Secular Terms	11
2.1	Introduction	11
2.2	A Simple Example: Damped Oscillator	11
2.3	Motion of a Particle in an Anharmonic Potential: Duffing Equation	17
2.3.1	Exact Solution of Duffing Equation	18
2.3.2	Naïve Perturbation Theory Applied to Duffing Equation	20
2.4	van der Pol Equation	23
2.5	Concluding Remarks	24
	Appendix: Method of Variation of Constants	25
3	Conventional Resummation Methods for Differential Equations	29
3.1	Introduction	29

3.2	Solvability Condition of Linear Equations and Appearance of Secular Terms: Fredholm's Alternative	29
3.3	Solvability Condition of Linear Differential Equations with Hermitian Operator	32
3.4	Lindstedt-Poincaré Method: Duffing Equation Revisited	35
3.5	Krylov-Bogoliubov-Mitropolsky Method	37
3.5.1	Generalities	38
3.5.2	Damped Oscillator	40
3.5.3	Duffing Equation	42
3.5.4	The van der Pol Equation	43
3.6	Multiple-Scale Method	45
3.6.1	Duffing Equation	46
3.6.2	Bifurcation in the Lorenz Model	48
Appendix 1:	Asymmetric Matrices, Left and Right Eigenvectors, and Symmetrized Inner Product	55
Appendix 2:	Solvability Condition of Linear Equations: Fredholm's Alternative Theorem	61
4	Renormalization Group Method for Global Analysis:	
	A Geometrical Formulation and Simple Examples	65
4.1	Introduction	65
4.2	Classical Theory of Envelopes and Its Adaptation for Global Analysis of Differential Equations	66
4.2.1	Envelope Curve in Two-Dimensional Space	66
4.2.2	Envelope Curves/Trajectories in n -Dimensional Space	70
4.2.3	Adaptation of the Envelope Theory in a Form Applicable to Dynamical Equations	72
4.3	Damped Oscillator in RG Method	73
4.3.1	Treatment as a Second-Order Differential Equation for Single Dependent Variable x	73
4.3.2	Treatment of Damped Oscillator as a System of First-Order Equations	79
4.4	RG/E Analysis of a Boundary-Layer Problem Without Matching; A System Treatment	84
4.5	The van der Pol Equation in RG Method	88
4.6	Jump Phenomenon in Forced Duffing Equation	91
4.7	Proof of a Global Validness of the Envelope Function as The approximate Solution to the Differential Equation	97
Appendix 1:	An RG Analysis of The Boundary-Layer Problem in the Scalar Representation	98
Appendix 2:	Useful Formulae of Solutions for Inhomogeneous Differential Equations for the RG Method	100