

**RISK: URGENT — CONSULT DOCTOR IMMEDIATELY**

**Chronic conditions:** Diabetes, Hypertension

## Vitals Summary

| Metric         | Readings | Average     | Min | Max | Status           |
|----------------|----------|-------------|-----|-----|------------------|
| Blood Pressure | 4        | 150/98 mmHg | 120 | 190 | FLAG             |
| SpO2           | 4        | 82%         | —   | —   | FLAG             |
| Pulse Rate     | 4        | 74 bpm      | 60  | 90  | OK               |
| Temperature    | 4        | 38.6°C      | —   | —   | OK               |
| Weight         | 4        | 82.7 kg     | —   | —   | change: +12.0 kg |

## Daily Wellness Summary

Symptom days — Headache: 1d | Fever: 0d | Chest Pain: 0d

## Health Goals Progress

**Sleep Goal** — —/7.5 hours (0%)

## AI Health Analysis

[illegible]

*Right now, prioritize complete physical rest and recovery. Once your vitals are medically stabilized, work on gradually improving your sleep to 7.5 hours and slowly increasing your daily step count.*

## Recommendations

Your weight loss and sleep goals are currently on hold as you must prioritize immediate medical stabilization of your oxygen, blood pressure, and fever.

1. Go to the nearest hospital emergency department immediately for oxygen therapy due to critically low SpO2 levels (81.8%).
2. Seek immediate medical attention to safely lower your blood pressure, which has reached a crisis level of 190/120 mmHg.
3. Consult your physician for a complete review of your diabetes and hypertension medications.
4. Monitor and record your blood pressure, temperature, and SpO2 levels 3 to 4 times daily once stabilized.
5. Ensure complete physical rest, stay hydrated with safe drinking water, and avoid any physical exertion until cleared by a doctor.

■ **Attention Required**

- Critically low average oxygen saturation (81.8%) requires immediate emergency medical evaluation and oxygen administration.
- Hypertensive crisis level blood pressure (190/120 mmHg) puts you at high risk for stroke or cardiovascular events; seek immediate care.
- High fever (38.6°C) combined with diabetes, hypertension, headache, and nausea requires urgent diagnostic workup.

Flag Details

| Flag Type                 | Count |
|---------------------------|-------|
| elevated_bp               | 1     |
| borderline_spo2           | 1     |
| fever                     | 3     |
| high_bp                   | 2     |
| critical_low_spo2         | 2     |
| low_spo2                  | 1     |
| hypertensive_crisis       | 1     |
| significant_weight_change | 1     |