

Combinators for Algebraic Structures (CAS)

Version 1

Timothy G. Griffin

July 12, 2016

Abstract

The document presents a small language of combinators for construction semigroups and semirings.

1 Motivation

We assume $2 \leq |S|$ for all carrier sets S .

2 Semigroup Combinators

2.1 Adding an identity

$$\text{AddId}(\alpha, (S, \bullet)) \equiv (S \uplus \{\alpha\}, \bullet_{\alpha}^{\text{id}})$$

where

$$a \bullet_{\alpha}^{\text{id}} b \equiv \begin{cases} a & (\text{if } b = \text{inr}(\alpha)) \\ b & (\text{if } a = \text{inr}(\alpha)) \\ \text{inl}(x \bullet y) & (\text{if } a = \text{inl}(x), b = \text{inl}(y)) \end{cases}$$

2.2 Adding an annihilator

$$\text{AddAn}(\omega, (S, \bullet)) \equiv (S \uplus \{\omega\}, \bullet_{\omega}^{\text{an}})$$

where

$$a \bullet_{\omega}^{\text{an}} b \equiv \begin{cases} \text{inr}(\omega) & (\text{if } b = \text{inr}(\omega)) \\ \text{inr}(\omega) & (\text{if } a = \text{inr}(\omega)) \\ \text{inl}(x \bullet y) & (\text{if } a = \text{inl}(x), b = \text{inl}(y)) \end{cases}$$

2.3 Direct Product

Let (S, \bullet) and (T, \diamond) be semigroups.

The direct product is denoted

$$(S, \bullet) \times (T, \diamond) \equiv (S \times T, \star)$$

where

$$\star = \bullet \times \diamond$$

is defined as

$$(s_1, t_1) \star (s_2, t_2) = (s_1 \bullet s_2, t_1 \diamond t_2).$$

2.4 Lexicographic Product

Suppose that semigroup (S, \bullet) is commutative, idempotent, and selective and that (T, \diamond) is a semigroup.

$$(S, \bullet) \vec{\times} (T, \diamond) \equiv (S \times T, \star)$$

where $\star \equiv \bullet \vec{\times} \diamond$ is defined as

$$(s_1, t_1) \star (s_2, t_2) = \begin{cases} (s_1 \bullet s_2, t_1 \diamond t_2) & s_1 = s_1 \bullet s_2 = s_2 \\ (s_1 \bullet s_2, t_1) & s_1 = s_1 \bullet s_2 \neq s_2 \\ (s_1 \bullet s_2, t_2) & s_1 \neq s_1 \bullet s_2 = s_2 \end{cases}$$

Examples for $(\mathbb{N}, \min) \vec{\times} (\mathbb{N}, \min)$.

$$\begin{aligned} (1, 17) \star (2, 3) &= (1, 17) \\ (2, 17) \star (2, 3) &= (2, 3) \\ (2, 3) \star (2, 3) &= (2, 3) \end{aligned}$$

Examples for $(\mathbb{N}, \min) \vec{\times} (\mathbb{N}, \max)$.

$$\begin{aligned} (1, 17) \star (2, 3) &= (1, 17) \\ (2, 17) \star (2, 3) &= (2, 17) \\ (2, 3) \star (2, 3) &= (2, 3) \end{aligned}$$

Examples for $(\mathbb{N}, \max) \vec{\times} (\mathbb{N}, \min)$.

$$\begin{aligned} (1, 17) \star (2, 3) &= (2, 3) \\ (2, 17) \star (2, 3) &= (2, 3) \\ (2, 3) \star (2, 3) &= (2, 3) \end{aligned}$$

2.5 Lifted Product

Assume (S, \bullet) is a semigroup. Let $\text{lift}(S, \bullet) \equiv (\text{fin}(2^S), \hat{\bullet})$ where

$$X \hat{\bullet} Y = \{x \bullet y \mid x \in X, y \in Y\}.$$

Example.

$$\{1, 3, 17\} \hat{+} \{1, 3, 17\} = \{2, 4, 6, 18, 20, 34\}$$

2.6 Minimal set union

Let \leq be a partial order on S . For $X \subseteq S$, define

$$\min_{\leq}(X) \equiv \{x \in X \mid \forall y \in X, \neg(y < x)\}.$$

Define

$$\mathcal{P}_{\text{fin}}(S, \leq) \equiv \{X \subseteq S \mid X \text{ finite and } \min_{\leq}(X) = X\}$$

and

$$A \cup_{\min}^{\leq} B \equiv \min_{\leq}(A \cup B)$$

$$\text{MinUnion}(S, \leq) \equiv (\mathcal{P}_{\text{fin}}(S, \leq), \cup_{\min}^{\leq})$$

NOTE: perhaps we can always define \leq in terms of a (S, oplus) , and so get rid of the class of structures (S, \leq) .

2.7 Minimal set lift

$$A \otimes_{\min}^{\leq} B \equiv \min_{\leq}(\{a \otimes b \mid a \in A, b \in B\})$$

$$\text{MinLift}(S, \leq, \otimes) \equiv (\mathcal{P}_{\text{fin}}(S, \leq), \otimes_{\min}^{\leq})$$

NOTE: perhaps we can always define \leq in terms of a (S, oplus) , and so get rid of the class of structures (S, \leq) .

3 Pre-order Combinators

4 Pre-order Semigroup Combinators

5 Bi-Semigroup Combinators

5.1 Adding a zero

$$\text{AddZero}(\bar{0}, (S, \oplus, \otimes)) \equiv (S \uplus \{\bar{0}\}, \oplus_0^{\text{id}}, \otimes_0^{\text{an}})$$

5.2 Adding a one

$$\text{AddOne}(\bar{1}, (S, \oplus, \otimes)) \equiv (S \uplus \{\bar{1}\}, \oplus_1^{\text{an}}, \otimes_1^{\text{id}})$$

5.3 Direct Product

$$(S, \oplus_S, \otimes_S) \vec{\times} (T, \oplus_T, \otimes_T) \equiv (S \times T, \oplus_S \times \oplus_T, \otimes_S \times \otimes_T)$$

5.4 Lexicographic Product

$$(S, \oplus_S, \otimes_S) \vec{\times} (T, \oplus_T, \otimes_T) \equiv (S \times T, \oplus_S \vec{\times} \oplus_T, \otimes_S \times \otimes_T)$$

5.5 \mathcal{M} and \mathcal{N}

Suppose $D \in \{L, R\}$. Let $\leq \equiv \leq_{\oplus}^D$ and define

$$\mathcal{M}(D, (S, \oplus, \otimes)) \equiv (\mathcal{P}_{\text{fin}}(S, \leq), \otimes_{\min}^{\leq}, \cup_{\min}^{\leq})$$

$$\mathcal{N}(D, (S, \oplus, \otimes)) \equiv (\mathcal{P}_{\text{fin}}(S, \leq), \cup_{\min}^{\leq}, \otimes_{\min}^{\leq})$$

Recall: $a \leq_{\oplus}^L b \equiv a = a \oplus b$ and $a \leq_{\oplus}^R b \equiv b = a \oplus b$.

6 Properties

(S, \bullet)

6.0.1 Initial set of semigroup properties

associative	$\mathbf{AS} \equiv \forall a, b, c \in S, a \bullet (b \bullet c) = (a \bullet b) \bullet c$
identity	$\mathbf{ID} \equiv \exists \alpha \in S, \forall a \in S, a = \alpha \bullet a = a \bullet \alpha$
annihilator	$\mathbf{AN} \equiv \exists \omega \in S, \forall a \in S, \omega = \omega \bullet a = a \bullet \omega$
commutative	$\mathbf{CM} \equiv \forall a, b \in S, a \bullet b = b \bullet a$
selective	$\mathbf{SL} \equiv \forall a, b \in S, a \bullet b \in \{a, b\}$
idempotent	$\mathbf{IP} \equiv \forall a \in S, a \bullet a = a$

6.0.2 Initial set of order properties

(S, \leq)

reflexive	$\mathbf{RX} \equiv \forall a \in S, a \leq a$
transitive	$\mathbf{TR} \equiv \forall a, b, c \in S, a \leq b \wedge b \leq c \implies a \leq c$
anti-symmetric	$\mathbf{AY} \equiv \forall a, b \in S, a \leq b \wedge b \leq a \implies a = b$
total	$\mathbf{TO} \equiv \forall a, b \in S, a \leq b \vee b \leq a$
bottom	$\mathbf{BM} \equiv \exists b \in S, \forall a \in S, b \leq a$
top	$\mathbf{TP} \equiv \exists t \in S, \forall a \in S, a \leq t$

6.0.3 Initial set of bi-semigroup properties

(S, \oplus, \otimes)

left distributivity	$\mathbf{LD} \equiv \forall a, b, c \in S, a \otimes (b \oplus c) = (a \otimes b) \oplus (a \otimes c)$
right distributivity	$\mathbf{RD} \equiv \forall a, b, c \in S, (a \oplus b) \otimes c = (a \otimes c) \oplus (b \otimes c)$
zero annihilates	$\mathbf{ZA} \equiv \exists \bar{0} \in S, \mathbf{IID}(S, \oplus, \bar{0}) \wedge \mathbf{IAN}(S, \otimes, \bar{0})$
one annihilates	$\mathbf{OA} \equiv \exists \bar{1} \in S, \mathbf{IID}(S, \otimes, \bar{1}) \wedge \mathbf{IAN}(S, \oplus, \bar{1})$

where

$$\begin{aligned}\mathbb{IID}(S, \bullet, \alpha) &\equiv \forall a \in S, a = \alpha \bullet a = a \bullet \alpha \\ \mathbb{IAN}(S, \bullet, \omega) &\equiv \forall a \in S, \omega = \omega \bullet a = a \bullet \omega\end{aligned}$$

6.0.4 Initial set of order semigroup properties

(S, \leq, \otimes)

$$\begin{array}{l|l} \text{left monotonicity} & \text{LM} \equiv \forall a, b, c \in S, a \leq b \implies (a \otimes c) \leq (b \otimes c) \\ \text{right monotonicity} & \text{RM} \equiv \forall a, b, c \in S, a \leq b \implies (c \otimes a) \leq (c \otimes b)\end{array}$$

7 Closure

To close we need extra properties.

For semigroups.

			required for
is right	$\mathbb{IR} \equiv \forall s, t \in S, s \bullet t = t$		$\mathbb{SL}(\times)$
is left	$\mathbb{IL} \equiv \forall s, t \in S, s \bullet t = s$		$\mathbb{SL}(\times)$
left cancellative	$\mathbb{LC} \equiv \forall a, b, c \in S, c \bullet a = c \bullet b \implies a = b$		$\mathbb{LD}(\vec{\times})$
right cancellative	$\mathbb{RC} \equiv \forall a, b, c \in S, a \bullet c = b \bullet c \implies a = b$		$\mathbb{RD}(\vec{\times})$
left constant	$\mathbb{LK} \equiv \forall a, b, c \in S, c \bullet a = c \bullet b$		$\mathbb{LD}(\vec{\times})$
right constant	$\mathbb{RK} \equiv \forall a, b, c \in S, a \bullet c = b \bullet c$		$\mathbb{RD}(\vec{\times})$
anti-left	$\mathbb{AL} \equiv \forall a, b \in S, a \bullet b \neq a$		$\mathbb{LC}(\text{AddId}), \mathbb{LAB}(\vec{\times}, \times)$
anti-right	$\mathbb{AR} \equiv \forall a, b \in S, b \bullet a \neq a$		$\mathbb{RC}(\text{AddId}), \mathbb{RAB}(\vec{\times}, \times)$

For bi-semigroups (S, \oplus, \otimes)

			required for
left absorption	$\mathbb{LAB} \equiv \forall a, b \in S, a = (b \otimes a) \oplus a = a \oplus (b \otimes a)$		$\mathbb{LD}(\text{AddOne})$
right absorption	$\mathbb{RAB} \equiv \forall a, b \in S, a = (a \otimes b) \oplus a = a \oplus (a \otimes b)$		$\mathbb{LD}(\text{AddOne})$

7.1 A few usefull implications

$$\begin{aligned}\mathbb{SL} &\implies \mathbb{IP} \\ \mathbb{RAB}(S, \oplus, \otimes) \wedge \mathbb{ID}(S, \oplus) &\implies \mathbb{IP}(S, \otimes) \\ \mathbb{LAB}(S, \oplus, \otimes) \wedge \mathbb{ID}(S, \oplus) &\implies \mathbb{IP}(S, \otimes) \\ \mathbb{LD}(S, \oplus, \otimes) \wedge \mathbb{OA}(S, \oplus, \otimes) &\implies \mathbb{RAB}(S, \oplus, \otimes) \\ \mathbb{RD}(S, \oplus, \otimes) \wedge \mathbb{OA}(S, \oplus, \otimes) &\implies \mathbb{LAB}(S, \oplus, \otimes)\end{aligned}$$

7.2 AddId

$AS(AddId(\alpha, (S, \bullet)))$	\Leftrightarrow	$AS(S, \bullet)$
$ID(AddId(\alpha, (S, \bullet)))$	\Leftrightarrow	$TRUE$
$AN(AddId(\alpha, (S, \bullet)))$	\Leftrightarrow	$AN(S, \bullet)$
$CM(AddId(\alpha, (S, \bullet)))$	\Leftrightarrow	$CM(S, \bullet)$
$IP(AddId(\alpha, (S, \bullet)))$	\Leftrightarrow	$IP(S, \bullet)$
$SL(AddId(\alpha, (S, \bullet)))$	\Leftrightarrow	$SL(S, \bullet)$
$IR(AddId(\alpha, (S, \bullet)))$	\Leftrightarrow	$FALSE$
$IL(AddId(\alpha, (S, \bullet)))$	\Leftrightarrow	$FALSE$
$LC(AddId(\alpha, (S, \bullet)))$	\Leftrightarrow	$LC(S, \bullet) \wedge AL(S, \bullet)$
$RC(AddId(\alpha, (S, \bullet)))$	\Leftrightarrow	$RC(S, \bullet) \wedge AR(S, \bullet)$
$LK(AddId(\alpha, (S, \bullet)))$	\Leftrightarrow	$FALSE$
$RK(AddId(\alpha, (S, \bullet)))$	\Leftrightarrow	$FALSE$
$AL(AddId(\alpha, (S, \bullet)))$	\Leftrightarrow	$FALSE$
$AR(AddId(\alpha, (S, \bullet)))$	\Leftrightarrow	$FALSE$

7.3 AddAn

$AS(AddAn(\alpha, (S, \bullet)))$	\Leftrightarrow	$AS(S, \bullet)$
$ID(AddAn(\alpha, (S, \bullet)))$	\Leftrightarrow	$ID(S, \bullet)$
$AN(AddAn(\alpha, (S, \bullet)))$	\Leftrightarrow	$TRUE$
$CM(AddAn(\alpha, (S, \bullet)))$	\Leftrightarrow	$CM(S, \bullet)$
$IP(AddAn(\alpha, (S, \bullet)))$	\Leftrightarrow	$IP(S, \bullet)$
$SL(AddAn(\alpha, (S, \bullet)))$	\Leftrightarrow	$SL(S, \bullet)$
$IR(AddAn(\alpha, (S, \bullet)))$	\Leftrightarrow	$FALSE$
$IL(AddAn(\alpha, (S, \bullet)))$	\Leftrightarrow	$FALSE$
$LC(AddAn(\alpha, (S, \bullet)))$	\Leftrightarrow	$FALSE$
$RC(AddAn(\alpha, (S, \bullet)))$	\Leftrightarrow	$FALSE$
$LK(AddAn(\alpha, (S, \bullet)))$	\Leftrightarrow	$FALSE$
$RK(AddAn(\alpha, (S, \bullet)))$	\Leftrightarrow	$FALSE$
$AL(AddAn(\alpha, (S, \bullet)))$	\Leftrightarrow	$FALSE$
$AR(AddAn(\alpha, (S, \bullet)))$	\Leftrightarrow	$FALSE$

7.4 \times

$$\begin{aligned}
\text{AS}((S, \bullet) \times (T, \diamond)) &\Leftrightarrow \text{AS}(S, \bullet) \wedge \text{AS}(T, \diamond) \\
\text{ID}((S, \bullet) \times (T, \diamond)) &\Leftrightarrow \text{ID}(S, \bullet) \wedge \text{ID}(T, \diamond) \\
\text{AN}((S, \bullet) \times (T, \diamond)) &\Leftrightarrow \text{AN}(S, \bullet) \wedge \text{AN}(T, \diamond) \\
\text{CM}((S, \bullet) \times (T, \diamond)) &\Leftrightarrow \text{CM}(S, \bullet) \wedge \text{CM}(T, \diamond) \\
\text{IP}((S, \bullet) \times (T, \diamond)) &\Leftrightarrow \text{IP}(S, \bullet) \wedge \text{IP}(T, \diamond) \\
\text{SL}((S, \bullet) \times (T, \diamond)) &\Leftrightarrow (\text{IR}(S, \bullet) \wedge \text{IR}(T, \diamond)) \vee (\text{IL}(S, \bullet) \wedge \text{IL}(T, \diamond)) \\
\text{IR}((S, \bullet) \times (T, \diamond)) &\Leftrightarrow \text{IR}(S, \bullet) \wedge \text{IR}(T, \diamond) \\
\text{IL}((S, \bullet) \times (T, \diamond)) &\Leftrightarrow \text{IL}(S, \bullet) \wedge \text{IL}(T, \diamond) \\
\text{LC}((S, \bullet) \times (T, \diamond)) &\Leftrightarrow \text{LC}(S, \bullet) \wedge \text{LC}(T, \diamond) \\
\text{RC}((S, \bullet) \times (T, \diamond)) &\Leftrightarrow \text{RC}(S, \bullet) \wedge \text{RC}(T, \diamond) \\
\text{LK}((S, \bullet) \times (T, \diamond)) &\Leftrightarrow \text{LK}(S, \bullet) \wedge \text{LK}(T, \diamond) \\
\text{RK}((S, \bullet) \times (T, \diamond)) &\Leftrightarrow \text{RK}(S, \bullet) \wedge \text{RK}(T, \diamond) \\
\text{AL}((S, \bullet) \times (T, \diamond)) &\Leftrightarrow \text{AL}(S, \bullet) \vee \text{AL}(T, \diamond) \\
\text{AR}((S, \bullet) \times (T, \diamond)) &\Leftrightarrow \text{AR}(S, \bullet) \vee \text{AR}(T, \diamond)
\end{aligned}$$

7.5 $\vec{\times}$ (semigroup)

Assuming $\text{AS}(S, \bullet)$, $\text{CM}(S, \bullet)$, and $\text{SL}(S, \bullet)$.

$$\begin{aligned}
\text{AS}((S, \bullet) \vec{\times} (T, \diamond)) &\Leftrightarrow \text{AS}(T, \diamond) \\
\text{ID}((S, \bullet) \vec{\times} (T, \diamond)) &\Leftrightarrow \text{ID}(S, \bullet) \wedge \text{ID}(T, \diamond) \\
\text{AN}((S, \bullet) \vec{\times} (T, \diamond)) &\Leftrightarrow \text{AN}(S, \bullet) \wedge \text{AN}(T, \diamond) \\
\text{CM}((S, \bullet) \vec{\times} (T, \diamond)) &\Leftrightarrow \text{CM}(T, \diamond) \\
\text{IP}((S, \bullet) \vec{\times} (T, \diamond)) &\Leftrightarrow \text{IP}(T, \diamond) \\
\text{SL}((S, \bullet) \vec{\times} (T, \diamond)) &\Leftrightarrow \text{SL}(T, \diamond) \\
\text{IR}((S, \bullet) \vec{\times} (T, \diamond)) &\Leftrightarrow \text{FALSE} \\
\text{IL}((S, \bullet) \vec{\times} (T, \diamond)) &\Leftrightarrow \text{FALSE} \\
\text{LC}((S, \bullet) \vec{\times} (T, \diamond)) &\Leftrightarrow \text{FALSE} \\
\text{RC}((S, \bullet) \vec{\times} (T, \diamond)) &\Leftrightarrow \text{FALSE} \\
\text{LK}((S, \bullet) \vec{\times} (T, \diamond)) &\Leftrightarrow \text{FALSE} \\
\text{RK}((S, \bullet) \vec{\times} (T, \diamond)) &\Leftrightarrow \text{FALSE} \\
\text{AL}((S, \bullet) \vec{\times} (T, \diamond)) &\Leftrightarrow \text{FALSE} \\
\text{AR}((S, \bullet) \vec{\times} (T, \diamond)) &\Leftrightarrow \text{FALSE}
\end{aligned}$$

7.6 lift

$\text{AS}(\text{lift}(S, \bullet))$	\Leftrightarrow	$\text{AS}(S, \bullet)$
$\text{ID}(\text{lift}(S, \bullet))$	\Leftrightarrow	$\text{ID}(S, \bullet) \ (\hat{\alpha} = \{\alpha\})$
$\text{AN}(\text{lift}(S, \bullet))$	\Leftrightarrow	$\text{TRUE} \ (\omega = \{\})$
$\text{CM}(\text{lift}(S, \bullet))$	\Leftrightarrow	$\text{CM}(S, \bullet)$
$\text{SL}(\text{lift}(S, \bullet))$	\Leftrightarrow	$\text{IL}(S, \bullet) \vee \text{IR}(S, \bullet) \vee (\text{IP}(S, \bullet) \wedge S = 2)$
$\text{IP}(\text{lift}(S, \bullet))$	\Leftrightarrow	$\text{SL}((S, \bullet))$
$\text{IL}(\text{lift}(S, \bullet))$	\Leftrightarrow	FALSE
$\text{IR}(\text{lift}(S, \bullet))$	\Leftrightarrow	FALSE
$\text{LC}(\text{lift}(S, \bullet))$	\Leftrightarrow	FALSE
$\text{RC}(\text{lift}(S, \bullet))$	\Leftrightarrow	FALSE
$\text{LK}(\text{lift}(S, \bullet))$	\Leftrightarrow	FALSE
$\text{RK}(\text{lift}(S, \bullet))$	\Leftrightarrow	FALSE
$\text{AL}(\text{lift}(S, \bullet))$	\Leftrightarrow	FALSE
$\text{AR}(\text{lift}(S, \bullet))$	\Leftrightarrow	FALSE

7.7 MinUnion

Assume $\text{BM}(S, \leq)$ and $\text{AY}(S, \leq)$. (AY does not seem to be required for associativity, idempotence... Don't think this is needed?)

$\text{AS}(\text{MinUnion}(S, \leq))$	\Leftrightarrow	TRUE
$\text{ID}(\text{MinUnion}(S, \leq))$	\Leftrightarrow	$\text{TRUE} \ (\alpha = \{\})$
$\text{AN}(\text{MinUnion}(S, \leq))$	\Leftrightarrow	$\text{TRUE} \ (\omega = \{\perp\})$
$\text{CM}(\text{MinUnion}(S, \leq))$	\Leftrightarrow	TRUE
$\text{SL}(\text{MinUnion}(S, \leq))$	\Leftrightarrow	$\text{TO}(S, \leq)$
$\text{IP}(\text{MinUnion}(S, \leq))$	\Leftrightarrow	TRUE
$\text{IL}(\text{MinUnion}(S, \leq))$	\Leftrightarrow	FALSE
$\text{IR}(\text{MinUnion}(S, \leq))$	\Leftrightarrow	FALSE
$\text{LC}(\text{MinUnion}(S, \leq))$	\Leftrightarrow	FALSE
$\text{RC}(\text{MinUnion}(S, \leq))$	\Leftrightarrow	FALSE
$\text{LK}(\text{MinUnion}(S, \leq))$	\Leftrightarrow	FALSE
$\text{RK}(\text{MinUnion}(S, \leq))$	\Leftrightarrow	FALSE
$\text{AL}(\text{MinUnion}(S, \leq))$	\Leftrightarrow	FALSE
$\text{AR}(\text{MinUnion}(S, \leq))$	\Leftrightarrow	FALSE

7.8 MinLift

Assume $\text{LM}(S, \leq, \otimes)$, $\text{RM}(S, \leq, \otimes)$, and $\text{AY}(S, \leq)$. (Needed for associativity)
Perhaps others constraints are needed to make progress on SL and IP . Lattice Theory to the rescue?

$$\begin{aligned}
\text{AS}(\text{MinLift}(S, \leq, \otimes)) &\Leftrightarrow \text{AS}(S, \otimes) \\
\text{ID}(\text{MinLift}(S, \leq, \otimes)) &\Leftrightarrow \text{ID}(S, \otimes) \\
\text{AN}(\text{MinLift}(S, \leq, \otimes)) &\Leftrightarrow \text{TRUE} \quad (\omega = \{\}) \\
\text{CM}(\text{MinLift}(S, \leq, \otimes)) &\Leftrightarrow \text{CM}(S, \otimes) \\
\text{SL}(\text{MinLift}(S, \leq, \otimes)) &\Leftrightarrow ??? \quad (\text{SOMEDAY !!!}) \\
\text{IP}(\text{MinLift}(S, \leq, \otimes)) &\Leftrightarrow ??? \quad (\text{SOMEDAY !!!}) \\
\text{IL}(\text{MinLift}(S, \leq, \otimes)) &\Leftrightarrow \text{FALSE} \\
\text{IR}(\text{MinLift}(S, \leq, \otimes)) &\Leftrightarrow \text{FALSE} \\
\text{LC}(\text{MinLift}(S, \leq, \otimes)) &\Leftrightarrow \text{FALSE} \\
\text{RC}(\text{MinLift}(S, \leq, \otimes)) &\Leftrightarrow \text{FALSE} \\
\text{LK}(\text{MinLift}(S, \leq, \otimes)) &\Leftrightarrow \text{FALSE} \\
\text{RK}(\text{MinLift}(S, \leq, \otimes)) &\Leftrightarrow \text{FALSE} \\
\text{AL}(\text{MinLift}(S, \leq, \otimes)) &\Leftrightarrow \text{FALSE} \\
\text{AR}(\text{MinLift}(S, \leq, \otimes)) &\Leftrightarrow \text{FALSE}
\end{aligned}$$

7.9 AddZero

$$\begin{aligned}
\text{LD}(\text{AddZero}(\bar{0}, (S, \oplus, \otimes))) &\Leftrightarrow \text{LD}(S, \oplus, \otimes) \\
\text{RD}(\text{AddZero}(\bar{0}, (S, \oplus, \otimes))) &\Leftrightarrow \text{RD}(S, \oplus, \otimes) \\
\text{ZA}(\text{AddZero}(\bar{0}, (S, \oplus, \otimes))) &\Leftrightarrow \text{TRUE} \\
\text{OA}(\text{AddZero}(\bar{0}, (S, \oplus, \otimes))) &\Leftrightarrow \text{OA}(S, \oplus, \otimes) \\
\text{RAB}(\text{AddZero}(\bar{0}, (S, \oplus, \otimes))) &\Leftrightarrow \text{RAB}(S, \oplus, \otimes) \\
\text{LAB}(\text{AddZero}(\bar{0}, (S, \oplus, \otimes))) &\Leftrightarrow \text{LAB}(S, \oplus, \otimes)
\end{aligned}$$

7.10 AddOne

$$\begin{aligned}
\text{LD}(\text{AddOne}(\bar{1}, (S, \oplus, \otimes))) &\Leftrightarrow \text{LD}(S, \oplus, \otimes) \wedge \text{RAB}(S, \oplus, \otimes) \wedge \text{IP}(S, \oplus) \\
\text{RD}(\text{AddOne}(\bar{1}, (S, \oplus, \otimes))) &\Leftrightarrow \text{RD}(S, \oplus, \otimes) \wedge \text{LAB}(S, \oplus, \otimes) \wedge \text{IP}(S, \oplus) \\
\text{ZA}(\text{AddOne}(\bar{1}, (S, \oplus, \otimes))) &\Leftrightarrow \text{ZA}(S, \oplus, \otimes) \\
\text{OA}(\text{AddOne}(\bar{1}, (S, \oplus, \otimes))) &\Leftrightarrow \text{TRUE} \\
\text{RAB}(\text{AddOne}(\bar{1}, (S, \oplus, \otimes))) &\Leftrightarrow \text{RAB}(S, \oplus, \otimes) \wedge \text{IP}(S, \oplus) \\
\text{LAB}(\text{AddOne}(\bar{1}, (S, \oplus, \otimes))) &\Leftrightarrow \text{LAB}(S, \oplus, \otimes) \wedge \text{IP}(S, \oplus)
\end{aligned}$$

7.11 Direct Product, (\times, \times)

$$\begin{aligned}
\text{LD}((S, \oplus_S, \otimes_S) \times (T, \oplus_T, \otimes_T)) &\Leftrightarrow \text{LD}(S, \oplus_S, \otimes_S) \wedge \text{LD}(T, \oplus_T, \otimes_T) \\
\text{RD}((S, \oplus_S, \otimes_S) \times (T, \oplus_T, \otimes_T)) &\Leftrightarrow \text{RD}(S, \oplus_S, \otimes_S) \wedge \text{RD}(T, \oplus_T, \otimes_T) \\
\text{ZA}((S, \oplus_S, \otimes_S) \times (T, \oplus_T, \otimes_T)) &\Leftrightarrow \text{ZA}(S, \oplus_S, \otimes_S) \wedge \text{ZA}(T, \oplus_T, \otimes_T) \\
\text{OA}((S, \oplus_S, \otimes_S) \times (T, \oplus_T, \otimes_T)) &\Leftrightarrow \text{OA}(S, \oplus_S, \otimes_S) \wedge \text{OA}(T, \oplus_T, \otimes_T) \\
\text{RAB}((S, \oplus_S, \otimes_S) \times (T, \oplus_T, \otimes_T)) &\Leftrightarrow \text{RAB}(S, \oplus_S, \otimes_S) \wedge \text{RAB}(T, \oplus_T, \otimes_T) \\
\text{LAB}((S, \oplus_S, \otimes_S) \times (T, \oplus_T, \otimes_T)) &\Leftrightarrow \text{LAB}(S, \oplus_S, \otimes_S) \wedge \text{LAB}(T, \oplus_T, \otimes_T)
\end{aligned}$$

7.12 Lexicographic Product, $(\vec{\times}, \times)$

Assume $\text{AS}(S, \oplus_S)$, $\text{CM}(S, \oplus_S)$, and $\text{SL}(S, \oplus_S)$.

$$\begin{aligned}
\text{LD}((S, \oplus_S, \otimes_S) \vec{\times} (T, \oplus_T, \otimes_T)) &\Leftrightarrow \text{LD}(S, \oplus_S, \otimes_S) \wedge \text{LD}(T, \oplus_T, \otimes_T) \\
&\quad \wedge (\text{LC}(S, \otimes_S) \vee \text{LK}(T, \otimes_T)) \\
\text{RD}((S, \oplus_S, \otimes_S) \vec{\times} (T, \oplus_T, \otimes_T)) &\Leftrightarrow \text{RD}(S, \oplus_S, \otimes_S) \wedge \text{RD}(T, \oplus_T, \otimes_T) \\
&\quad \wedge (\text{RC}(S, \otimes_S) \vee \text{RK}(T, \otimes_T)) \\
\text{ZA}((S, \oplus_S, \otimes_S) \vec{\times} (T, \oplus_T, \otimes_T)) &\Leftrightarrow \text{ZA}(S, \oplus_S, \otimes_S) \wedge \text{ZA}(T, \oplus_T, \otimes_T) \\
\text{OA}((S, \oplus_S, \otimes_S) \vec{\times} (T, \oplus_T, \otimes_T)) &\Leftrightarrow \text{OA}(S, \oplus_S, \otimes_S) \wedge \text{OA}(T, \oplus_T, \otimes_T) \\
\text{RAB}((S, \oplus_S, \otimes_S) \vec{\times} (T, \oplus_T, \otimes_T)) &\Leftrightarrow \text{RAB}(S, \oplus_S, \otimes_S) \wedge (\text{RAB}(T, \oplus_T, \otimes_T) \vee \text{AR}(S, \otimes_S)) \\
\text{LAB}((S, \oplus_S, \otimes_S) \vec{\times} (T, \oplus_T, \otimes_T)) &\Leftrightarrow \text{LAB}(S, \oplus_S, \otimes_S) \wedge (\text{LAB}(T, \oplus_T, \otimes_T) \vee \text{AL}(S, \otimes_S))
\end{aligned}$$

7.13 \mathcal{N}

Which constraints should we impose on (S, \oplus, \otimes) in order to make this easier, cleaner? And how can we do this without giving up too much in terms of expressive power?

$$\begin{aligned}
\text{LD}(\mathcal{N}(L, (S, \oplus, \otimes))) &\Leftrightarrow ??? \\
\text{RD}(\mathcal{N}(L, (S, \oplus, \otimes))) &\Leftrightarrow ??? \\
\text{ZA}(\mathcal{N}(L, (S, \oplus, \otimes))) &\Leftrightarrow ??? \\
\text{OA}(\mathcal{N}(L, (S, \oplus, \otimes))) &\Leftrightarrow ??? \\
\text{RAB}(\mathcal{N}(L, (S, \oplus, \otimes))) &\Leftrightarrow ??? \\
\text{LAB}(\mathcal{N}(L, (S, \oplus, \otimes))) &\Leftrightarrow ???
\end{aligned}$$

$$\begin{aligned}
\text{LD}(\mathcal{N}(R, (S, \oplus, \otimes))) &\Leftrightarrow ??? \\
\text{RD}(\mathcal{N}(R, (S, \oplus, \otimes))) &\Leftrightarrow ??? \\
\text{ZA}(\mathcal{N}(R, (S, \oplus, \otimes))) &\Leftrightarrow ??? \\
\text{OA}(\mathcal{N}(R, (S, \oplus, \otimes))) &\Leftrightarrow ??? \\
\text{RAB}(\mathcal{N}(R, (S, \oplus, \otimes))) &\Leftrightarrow ??? \\
\text{LAB}(\mathcal{N}(R, (S, \oplus, \otimes))) &\Leftrightarrow ???
\end{aligned}$$

7.14 \mathcal{M}

$$\begin{aligned}
\text{LD}(\mathcal{M}(L, (S, \oplus, \otimes))) &\Leftrightarrow ??? \\
\text{RD}(\mathcal{M}(L, (S, \oplus, \otimes))) &\Leftrightarrow ??? \\
\text{ZA}(\mathcal{M}(L, (S, \oplus, \otimes))) &\Leftrightarrow ??? \\
\text{OA}(\mathcal{M}(L, (S, \oplus, \otimes))) &\Leftrightarrow ??? \\
\text{RAB}(\mathcal{M}(L, (S, \oplus, \otimes))) &\Leftrightarrow ??? \\
\text{LAB}(\mathcal{M}(L, (S, \oplus, \otimes))) &\Leftrightarrow ???
\end{aligned}$$

$$\begin{aligned}
\text{LD}(\mathcal{M}(R, (S, \oplus, \otimes))) &\Leftrightarrow ??? \\
\text{RD}(\mathcal{M}(R, (S, \oplus, \otimes))) &\Leftrightarrow ??? \\
\text{ZA}(\mathcal{M}(R, (S, \oplus, \otimes))) &\Leftrightarrow ??? \\
\text{OA}(\mathcal{M}(R, (S, \oplus, \otimes))) &\Leftrightarrow ??? \\
\text{RAB}(\mathcal{M}(R, (S, \oplus, \otimes))) &\Leftrightarrow ??? \\
\text{LAB}(\mathcal{M}(R, (S, \oplus, \otimes))) &\Leftrightarrow ???
\end{aligned}$$

8 Other Problems

8.1 Matrix algebras

8.2 swap

We might want to define

$$\text{swap}(S, \oplus, \otimes) \equiv (S, \otimes, \oplus).$$

The disadvantage of this is that we would then have to introduce (and close with respect to) these “dual” properties:

$$\begin{aligned} \text{LD}^\delta(S, \oplus, \otimes) &\equiv \forall a, b, c \in S, a \oplus (b \otimes c) = (a \oplus b) \otimes (a \oplus c) \\ \text{RD}^\delta(S, \oplus, \otimes) &\equiv \forall a, b, c \in S, (a \otimes b) \oplus c = (a \oplus c) \otimes (b \oplus c) \\ \text{RAB}^\delta(S, \oplus, \otimes) &\equiv \forall a, b \in S, a = (a \oplus b) \otimes a = a \otimes (a \oplus b) \\ \text{LAB}^\delta(S, \oplus, \otimes) &\equiv \forall a, b \in S, a = (b \oplus a) \otimes a = a \otimes (b \oplus a) \end{aligned}$$

8.3 k -best paths

8.4 Semi-direct product

8.5 General reductions