EASY NOTES

MA PSYCHOLOGY 1ST YEAR

PAPER 06: MPC-06

(Statistics in Psychology)

(A Series of Important Topics for Examination. Based on Previous 10 Years Question Papers)

For Term End Examination - June 2025

Achiever's Hive

EASY NOTES

by

Ms. Neha Pandey
(Psychologist & Teacher, Founder of Achiever's Hive)

First Published: October 2024

Free distribution of this document, except the author, will be considered as COPYRIGHT Violation.

All rights reserved. No part of this document can be reproduced, stored in or introduced into a retrieval system or **transmitted** in any form or by any means (electrical, mechanical, photocopying, recording or otherwise)

All data were deemed correct at time of creation. Author is not liable for errors or omissions.

SHARING THIS DOCUMENT IS NOT ONLY ILLEGAL BUT UNETHICAL AS WELL.

Please report such sharing to us on:
WhatsApp: +91 8979564963
Email: achievershive.in@gmail.com

Professional Guidance for:

MAPC 1st Year Practical: MPCL 007 Practical File

MAPC 2nd Year Practical: MPCE 014 / MPCE 024 / MPCE 034 Practical File

Internship: Internship Report with Case Study

WhatsApp: +91-8979564963

TABLE OF CONTENTS

Sr. No.	Topic (Theory)	Pg. No.
1.	Levels Of Measurement	5
2.	Normal Distribution: Characteristics & Applications	7
3.	Divergence In Normality	11
4.	Parametric Statistics	14
5.	Non-Parametric Test	18
6.	Descriptive Statistics	21
7.	Inferential Statistics	25
8.	Measures Of Central Tendency	29
9.	Measures of Dispersion/Measures of Variability	32
10.	Partial & Part Correlation	37
11.	Point Biserial Correlation vs Phi-Coefficient	43
12.	Point & Interval Estimation	47
13.	Type I & Type II Errors	50
14.	Levels Of Significance	53
15.	Skewness	54
16.	Hypothesis Testing	57
17.	Standard Error	58
18.	Steps In Computation Of Regression	60

Achiever's Hive

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

Sr. No.	Topic (Theory)	Pg. No.
19.	Interactional Effect	62
20.	Correlation	64
21.	Two Way Anova	68

Sr. No.	Topic (Numerical)	Pg. No.
1.	Mann Whitney U Test	72
2.	Chi Square	74
3.	Regression Equation	76
4.	Kendall's Tau	78
5.	Spearman's Rho	80
6.	Pearson Product Moment Correlation	82
7.	One Way Anova -1	84
8.	One Way Anova -2 (With Critical Values)	89
9.	T-Test	95

Achiever's Hive

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

1. Levels of Measurement

Levels of measurement, also known as scales of measurement, describe how data can be categorized and quantified. Proposed by psychologist Stanley Smith Stevens, these levels are foundational in statistics and research, as they determine what kind of mathematical operations and statistical analyses are appropriate for a dataset. There are four levels of measurement, each with increasing complexity: **nominal, ordinal, interval,** and **ratio**.

1. Nominal Level

Definition: The nominal level of measurement categorizes data into distinct, mutually exclusive groups that have no inherent order or ranking.

Characteristics:

- **Categories**: Data are simply labels or names, with no quantitative value.
- **No Order**: The groups or categories have no natural order.
- **Examples**: Gender (male, female), eye colour (blue, brown, green), types of fruit (apple, banana, orange).
- **Statistical Analysis**: Frequencies, mode, and chi-square tests (to test for associations between nominal variables).

2. Ordinal Level

Definition: The ordinal level of measurement categorizes data into ordered groups, where the order matters but the intervals between the values are not consistent or meaningful.

Characteristics:

- **Order**: Data have a ranked or ordered sequence.
- **Unequal Intervals**: The differences between ranks may not be equal or measurable.
- **Examples**: Education level (high school, bachelor's, master's), customer satisfaction ratings (satisfied, neutral, dissatisfied), socioeconomic status (low, middle, high).
- **Statistical Analysis**: Mode, median, percentiles, and non-parametric tests like the Mann-Whitney U test.

3. Interval Level

Definition: The interval level of measurement provides not only order but also equal intervals between values. However, it lacks an absolute zero point, which means ratios (multiplications and divisions) cannot be meaningfully computed.

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

Characteristics:

- **Order and Equal Intervals**: Data have a defined order, and the intervals between values are meaningful and equal.
- **No True Zero**: There is no true zero point, meaning zero does not represent the absence of the characteristic.
- **Examples**: Temperature in Celsius or Fahrenheit (e.g., 20°C, 30°C, 40°C), IQ scores, calendar years.
- **Statistical Analysis**: Mean, standard deviation, correlation, and t-tests (as long as the data are normally distributed).

4. Ratio Level

Definition: The ratio level of measurement has all the features of interval data (order, equal intervals) and includes a true zero point, allowing for meaningful ratios.

Characteristics:

- **Order, Equal Intervals, and True Zero**: The presence of a true zero means that values can be compared in terms of ratios.
- **Examples**: Height, weight, age, income, time, distance.
- **Statistical Analysis**: All mathematical and statistical operations are permissible, including geometric and harmonic means, t-tests, ANOVA, and regression.

Summary of Levels of Measurement

Level	Order	Equal Intervals	True Zero	Examples
Nominal	No	No	No	Gender, nationality, types of pets
Ordinal	Yes	No	No	Satisfaction ratings, education level
Interval	Yes	Yes	No	Temperature (°C, °F), IQ scores
Ratio	Yes	Yes	Yes	Age, weight, income, distance

Importance of Levels of Measurement

Understanding the level of measurement is crucial because it dictates the type of statistical tests and data manipulation that are appropriate. Using the wrong statistical methods with a particular level of data can lead to inaccurate or misleading results. For instance,


This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

calculating the mean of nominal data (like computing an "average gender") is meaningless, while calculating the mean of interval or ratio data (like average age) is standard practice.

In sum, the levels of measurement serve as a guideline for researchers to appropriately handle data, ensuring accurate analysis and interpretation.

2. Normal Distribution

The **normal distribution**, also known as the **Gaussian distribution**, is one of the most fundamental and widely used probability distributions in statistics. It describes a continuous probability distribution that is symmetric and bell-shaped, where most observations cluster around the central peak, and probabilities taper off symmetrically on both sides. In many natural and social phenomena, normal distribution approximates the distribution of data, making it a crucial concept in statistical analysis.



Characteristics of Normal Distribution

I. **Bell-Shaped Curve**: The normal distribution has a symmetrical bell-shaped curve where the mean, median, and mode of the data all fall at the same point, right at the center of the distribution.

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

II. **Symmetry**: The distribution is symmetric around its mean, meaning that the left and right sides of the curve are mirror images of each other. This symmetry implies that about half the data values are less than the mean and half are greater than the mean.

III. **Mean, Median, and Mode Are Equal**: In a perfectly normal distribution, the mean, median, and mode all coincide at the center of the distribution, highlighting its symmetry.

IV. **Asymptotic**: The tails of the normal distribution curve approach, but never touch, the horizontal axis. This implies that extreme values (very high or very low) are possible but increasingly unlikely as they deviate further from the mean.

V. **Defined by Two Parameters**:
- **Mean (μ)**: The central tendency of the distribution, which determines the position of the centre of the curve.
- **Standard Deviation (σ)**: Measures the spread or dispersion of the data. A larger standard deviation results in a flatter and wider curve, while a smaller standard deviation produces a steeper and narrower curve.

VI. **68-95-99.7 Rule** (Empirical Rule): For a normal distribution:
- About **68%** of data falls within one standard deviation (±1σ) from the mean.
- About **95%** of data falls within two standard deviations (±2σ) from the mean.
- About **99.7%** of data falls within three standard deviations (±3σ) from the mean.

VII. **Continuous Probability Distribution**: The normal distribution is continuous, meaning that it deals with continuous data and the probability for any single exact point (e.g., the probability of a specific value) is zero. Instead, probabilities are calculated over intervals.

The normal distribution curve has one peak, or mode, at the mean, indicating that most values are clustered around the central point.

Mathematical Formula for Normal Distribution

The probability density function (PDF) of the normal distribution is given by:

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

$$f(x|\mu,\sigma) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

Where:

- x is the variable.
- μ is the mean of the distribution.
- σ is the standard deviation of the distribution.
- σ^2 is the variance.
- e is Euler's number, approximately equal to 2.71828.
- π is approximately equal to 3.14159.

This formula defines the bell-shaped curve of a normal distribution.

Applications of Normal Distribution

I. Natural and Social Phenomena:

Many biological, psychological, and social variables (e.g., heights, IQ scores, blood pressure) approximately follow a normal distribution. The normal distribution helps in understanding and predicting real-world phenomena.

II. Statistical Inference:

Central Limit Theorem (CLT): One of the most important applications of normal distribution is in the **central limit theorem**, which states that the distribution of the sample means approaches a normal distribution as the sample size becomes large, regardless of the

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

population's original distribution. This is foundational for hypothesis testing, confidence intervals, and other inferential statistics.

III. Hypothesis Testing:

Normal distribution is essential for conducting various statistical tests such as **t-tests**, **Z-tests**, and **ANOVA**. These tests assume that the underlying data is normally distributed or approximately normal for large samples.

IV. Quality Control:

In manufacturing and quality control processes, normal distribution is used to monitor the performance of systems through control charts. Companies use the distribution to detect variations in production processes and ensure that outputs fall within acceptable limits (e.g., Six Sigma).

V. Finance and Economics:

Normal distribution is widely used in finance to model stock prices, returns, and risk. For example, in portfolio theory, stock returns are often assumed to follow a normal distribution, allowing for the calculation of risk (standard deviation) and expected returns.

VI. Grading and Standardized Testing:

In education, standardized test scores like the SAT, GRE, and IQ tests are often normalized, meaning the results are adjusted to follow a normal distribution. This enables comparison across large populations and different administrations of the test.

VII. Prediction and Forecasting:

Normal distribution is applied in forecasting models, especially where predictions involve large datasets. The assumption of normality allows for the calculation of confidence intervals, error margins, and the likelihood of future events.

VIII. Behavioral Sciences:

In psychology and other behavioral sciences, traits such as intelligence, personality measures, and certain cognitive abilities are often assumed to be normally distributed, facilitating comparison and analysis across populations.

IX. Medicine and Health Sciences:

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

Medical researchers use normal distribution to interpret data like blood pressure, cholesterol levels, or drug efficacy rates, allowing for the identification of abnormal values or outliers that may indicate a health problem.

X. Sampling and Estimation:

In many sampling techniques, it is assumed that the distribution of sample means is normally distributed for sufficiently large sample sizes. This assumption simplifies the process of estimating population parameters (such as the population mean) from sample data.

The normal distribution is a cornerstone of probability theory and statistics, used extensively in various fields to model real-world data, make predictions, and conduct statistical analyses. Its key features, including its bell-shaped curve, symmetry, and the 68-95-99.7 rule, provide valuable insights into the behavior of random variables. Through its applications in hypothesis testing, quality control, finance, and scientific research, the normal distribution enables meaningful interpretations and decisions based on statistical data.

3. Divergence in Normality

In statistics, **divergence in normality** refers to deviations or departures from a normal (Gaussian) distribution in a dataset. Since many statistical methods, including parametric tests like t-tests, ANOVA, and linear regression, assume that the underlying data is normally distributed, divergence from normality can impact the reliability and validity of these analyses. Recognizing and addressing these divergences is essential to ensure accurate results and interpretations.

Types of Divergence in Normality

A. Skewness: Skewness describes the asymmetry of the data distribution around the mean.

- **Positive Skew (Right Skew)**: The distribution has a longer right tail, with most values concentrated on the left. This is common in income or age data where a few high values pull the mean to the right.
- **Negative Skew (Left Skew)**: The distribution has a longer left tail, with most values on the right. This is less common but can appear in data like test scores where most students score high with a few lower outliers.

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

B. **Kurtosis**: Kurtosis measures the "tailedness" or the extremity of deviations within a distribution.
   - **Leptokurtic**: Distributions with high kurtosis have heavier tails and a sharper peak than normal distributions. This indicates more frequent extreme values, suggesting outliers or high variance.
   - **Platykurtic**: Distributions with low kurtosis have lighter tails and a flatter peak, meaning fewer extreme values.

C. **Outliers**: Extreme values that are significantly different from the rest of the data can cause divergence from normality by stretching or skewing the distribution.

D. **Multimodality**: While a normal distribution is unimodal (has a single peak), some distributions may have multiple peaks (multimodal), indicating the presence of subgroups within the data.

E. **Bimodality**: This is a specific case of multimodality where there are two clear peaks. Bimodal distributions often indicate that data is composed of two different populations or categories.

Why Divergence from Normality Matters

A. **Impact on Statistical Tests**: Many statistical tests assume normality for accurate calculations. When data diverges from normality, it can:
   - Affect p-values and confidence intervals, increasing the likelihood of Type I or Type II errors.
   - Lead to misleading conclusions, particularly with small sample sizes where the Central Limit Theorem (which states that sample means tend to follow a normal distribution as sample size grows) may not apply as effectively.

B. **Influence on Descriptive Statistics**: Means and standard deviations are sensitive to outliers and skewness. Divergence from normality can make these statistics less representative of the dataset, impacting the interpretation of data.

C. **Modeling and Prediction**: Divergence in normality affects assumptions in predictive models, such as linear regression, that rely on normally distributed residuals (errors). Non-normal residuals can bias parameter estimates and lead to unreliable predictions.

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

Assessing Divergence from Normality

A. **Graphical Methods**:
- **Histograms** and **Density Plots**: Visualize the data's shape to identify skewness, kurtosis, and multimodality.
- **Q-Q (Quantile-Quantile) Plot**: Compares the quantiles of the dataset to a theoretical normal distribution. Points falling along a straight line indicate normality, while deviations from the line signal divergence.

B. **Statistical Tests**:
- **Shapiro-Wilk Test**: A statistical test that evaluates whether a dataset differs significantly from a normal distribution. A p-value below a certain threshold (e.g., 0.05) suggests divergence from normality.
- **Kolmogorov-Smirnov Test**: Another test comparing the sample distribution to a normal distribution. It is sensitive to both the shape and spread of the distribution.
- **Skewness and Kurtosis Coefficients**: Calculate skewness and kurtosis to quantify the deviation. Skewness values outside the range of -1 to 1 and kurtosis values significantly different from 3 (the kurtosis of a normal distribution) suggest divergence.

Addressing Divergence from Normality

A. **Data Transformation**:
- **Log Transformation**: Often used for right-skewed data to reduce skewness.
- **Square Root Transformation**: Useful for moderately skewed data, reducing the impact of large values.
- **Box-Cox Transformation**: A flexible method that applies power transformations to bring data closer to normality.

B. **Non-Parametric Tests**: When data is non-normal, non-parametric tests like the Mann-Whitney U test, Wilcoxon signed-rank test, and Kruskal-Wallis test can be used, as they do not assume normality.

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

C. **Robust Statistical Techniques**: Robust methods, such as using medians and interquartile ranges instead of means and standard deviations, reduce sensitivity to non-normal data.

D. **Bootstrap Methods**: Resampling techniques like bootstrapping generate sampling distributions empirically, helping estimate statistics without assuming normality.

Practical Example

Consider a study examining salaries across a population. Salary data is often right-skewed, with a few extremely high salaries pulling the mean upward. Analyzing this data under the assumption of normality could lead to erroneous conclusions, particularly if relying on mean salary as a measure of central tendency. Here, using a median (a robust statistic) or transforming the data could improve accuracy.

Divergence from normality in statistics is a significant consideration, especially in fields requiring precise analyses. Identifying and addressing non-normality ensures the robustness of statistical conclusions, ultimately leading to more reliable insights and decisions.

4. Parametric Statistics

Parametric statistics refers to statistical techniques that make certain assumptions about the parameters (such as the mean and standard deviation) of the population distribution from which the sample data are drawn. These methods are widely used when the data follows a known and specific distribution, typically the **normal distribution**. Parametric tests, such as t-tests, ANOVA, and regression analysis, are more powerful when the assumptions hold true because they rely on the distribution's parameters to make inferences about the population.

Key Assumptions of Parametric Statistics

For parametric statistical methods to be valid, certain assumptions about the data must be satisfied. The most common assumptions include:

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

1. Normality

Assumption: The data is normally distributed, or the distribution of the population from which the sample is drawn is normal.

Explanation: This means that the data should form a bell-shaped, symmetrical curve (normal distribution). Even if the individual data points are not perfectly normal, the distribution of the sample means (as per the Central Limit Theorem) should be approximately normal, especially for large sample sizes.

2. Homogeneity of Variance (Homoscedasticity)

Assumption: The variance among groups or samples is roughly equal.

Explanation: For example, when comparing two or more groups, the spread of scores (variance) should be similar across these groups. This assumption is crucial for tests like the t-test or ANOVA, where unequal variances (heteroscedasticity) can lead to misleading results.

3. Independence of Observations

Assumption: The data points or observations are independent of each other.

Explanation: This means that the value of one observation should not influence or be related to the value of another observation. Independence is critical in most statistical tests to avoid biased estimates and incorrect inferences.

4. Interval or Ratio Scale of Measurement

Assumption: Data is measured on an interval or ratio scale.

Explanation: Parametric tests require that the data be quantitative and continuous, with equal intervals between values (e.g., temperature, height, income). These scales allow for meaningful calculations like means and standard deviations.

5. Linearity (for some tests)

Assumption: There is a linear relationship between variables (for tests like regression analysis).

Explanation: When conducting tests such as correlation or regression analysis, it is assumed that the relationship between the independent and dependent variables is linear, meaning that the change in one variable is proportional to the change in another.

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

Common Parametric Tests

1. **t-Test**: Compares the means of two groups to determine if they are significantly different.
2. **ANOVA (Analysis of Variance)**: Compares the means of three or more groups to assess if there are significant differences.
3. **Pearson Correlation**: Measures the strength and direction of the linear relationship between two variables.
4. **Regression Analysis**: Assesses the relationship between a dependent variable and one or more independent variables.
5. **Z-Test**: Used to determine if there is a significant difference between sample means and population means, typically when the sample size is large.

When Parametric Tests are Not Suitable

If the assumptions of parametric tests are not met, alternative **non-parametric tests** are used, which do not rely on strict assumptions about the underlying population distribution. These include tests like the Mann-Whitney U test, Wilcoxon signed-rank test, and Kruskal-Wallis test.

Advantages of Parametric Tests

I. **Higher Statistical Power**: Parametric tests are generally more powerful than non-parametric tests, meaning they are more likely to detect a true effect or difference if one exists. This is because they use more information from the data, such as means and standard deviations.
II. **Assumption of Normality Provides Precision**: Parametric tests make specific assumptions about the data (e.g., normal distribution), which allows for more precise and reliable estimates when these assumptions hold true.
III. **Wider Range of Applications**: Parametric tests are suitable for many different types of data and analyses, including comparisons between means (e.g., t-tests, ANOVA), regression analysis, and correlation, making them versatile tools for statistical analysis.
IV. **Incorporate Distribution Parameters**: They utilize the actual values of the data (not just ranks or orderings), which allows for more detailed and nuanced interpretations of results, especially when calculating effect sizes, confidence intervals, and probabilities.

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

V. **Well-Developed Statistical Theories**: Parametric tests have a strong foundation in statistical theory, meaning there are clear guidelines on how to use them, interpret results, and apply them to various situations. This makes them widely accepted and understood across disciplines.
VI. **Smaller Samples May Still Work**: Under certain conditions, parametric tests can still perform adequately even with relatively small sample sizes, provided that the assumptions (like normality) are satisfied.

Disadvantages of Parametric Tests

I. **Assumptions Must Be Met**: Parametric tests require strict assumptions, including normality, homogeneity of variance, and linearity (for regression). When these assumptions are violated, the test results may be invalid, leading to biased or incorrect conclusions.
II. **Sensitive to Outliers**: Parametric tests are more sensitive to outliers than non-parametric tests. Extreme values can skew the results because they rely on means and variances, which are affected by outliers.
III. **Limited Use for Non-Interval Data**: Parametric tests typically require data on an interval or ratio scale. They are not appropriate for ordinal or nominal data, which limits their use for certain types of variables.
IV. **Not Robust to Non-Normal Data**: If the data is not normally distributed and the sample size is small, parametric tests may not perform well. The results could be misleading in terms of significance or effect size.
V. **Large Sample Size Requirement**: In some cases, parametric tests require large sample sizes to ensure that the central limit theorem applies (i.e., the distribution of the sample means is approximately normal). When sample sizes are small, the test's reliability diminishes.
VI. **Less Flexible**: Parametric tests are rigid in their application. If the assumptions are violated, a different approach (such as a non-parametric test) must be used, reducing their flexibility.

Parametric tests are powerful and widely used in statistical analysis, especially when the assumptions of normality, equal variances, and linearity hold true. However, their reliance on these assumptions makes them less suitable for certain data types or situations, especially when dealing with non-normal distributions, small sample sizes, or data with

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

outliers. Understanding when and how to use parametric versus non-parametric tests is critical for accurate data analysis and interpretation.

5. Non-Parametric Tests

Non-parametric tests are statistical methods that do not rely on assumptions about the population's distribution. They are often used when parametric test assumptions, such as normality or homogeneity of variance, are violated, or when dealing with ordinal data, nominal data, or small sample sizes. Non-parametric methods are more flexible because they don't require the data to fit a specific distribution, making them useful for analyzing data that is skewed or has outliers.

Non-parametric tests often rank the data instead of using the actual values, which makes them less sensitive to outliers and non-normal distributions.

Common Non-Parametric Tests

- **Mann-Whitney U Test**: Compares two independent groups when the data does not meet the assumptions of a t-test. It's the non-parametric equivalent of the independent t-test.
- **Wilcoxon Signed-Rank Test**: Compares two related samples or repeated measures (like a paired t-test) to assess differences in their medians.
- **Kruskal-Wallis Test**: Compares three or more independent groups, similar to ANOVA, but without assuming normality.
- **Spearman's Rank Correlation**: Measures the strength and direction of the monotonic relationship between two variables, used when Pearson correlation assumptions are not met.
- **Chi-Square Test**: Tests the relationship between categorical variables, often used for contingency tables to assess independence between variables.
- **Friedman Test**: Compares three or more paired groups, used as a non-parametric alternative to repeated measures ANOVA.

Assumptions of Non-Parametric Tests

Non-parametric tests are more robust because they are less dependent on strict assumptions. However, they still have some key assumptions:

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

1. Ordinal or Nominal Data (or Non-Normally Distributed Interval/Ratio Data)

Assumption: Non-parametric tests are appropriate for ordinal data (ranked data), nominal data (categorical data), or interval/ratio data that are not normally distributed.

Explanation: These tests can handle data measured on less precise scales, and they do not rely on the mean or variance of the data. Non-parametric methods focus on the rank or the median rather than the mean.

2. Independence of Observations

Assumption: The observations or data points should be independent of one another, particularly for tests like the Mann-Whitney U test, Kruskal-Wallis test, or Chi-Square test.

Explanation: Similar to parametric tests, non-parametric tests assume that the data points from different groups or samples are unrelated. For paired tests like the Wilcoxon signed-rank test, however, this assumption applies within the paired data (i.e., observations must be paired correctly and independently).

3. Ranking (for some tests)

Assumption: Non-parametric tests often rely on the ranking of data rather than the actual values (e.g., Mann-Whitney U test, Kruskal-Wallis test).

Explanation: The tests typically rank the data points from the smallest to the largest. Therefore, the actual values are not as important as the order of the data points.

4. Distribution-Free

Assumption: Non-parametric tests do not assume a specific distribution for the population (e.g., normal distribution).

Explanation: This is one of the major benefits of non-parametric tests. They can be used with data that is skewed, has outliers, or does not fit into a normal distribution, as the analysis is based on the ranking of data rather than its distribution.

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

5. Continuous or Discrete Data (for Some Tests)

Assumption: Some non-parametric tests assume that the data is continuous (e.g., Wilcoxon signed-rank test, Mann-Whitney U test).

Explanation: Although many non-parametric tests are suitable for ordinal data, some, like the Mann-Whitney U test and the Wilcoxon signed-rank test, work best with continuous data that is ranked.

Advantages of Non-Parametric Tests

I. **Fewer Assumptions**: Non-parametric tests do not require data to follow a normal distribution, making them more versatile for various types of data.
II. **Robust to Outliers**: Since non-parametric tests often rely on ranks rather than raw data values, they are less sensitive to extreme values or outliers.
III. **Applicable to Small Samples**: They can be used effectively with small sample sizes, where parametric tests might not be reliable.
IV. **Useful for Ordinal and Nominal Data**: Non-parametric tests can be applied to data measured on ordinal or nominal scales, unlike parametric tests, which require interval or ratio data.

Disadvantages of Non-Parametric Tests

I. **Less Power**: Non-parametric tests are generally less powerful than parametric tests when parametric assumptions are met, meaning they may not detect differences when they exist.
II. **Less Information**: By relying on ranks rather than actual data values, non-parametric tests can lose some information that could be important in the analysis.
III. **Limited Interpretations**: The results of non-parametric tests may be less straightforward to interpret than parametric tests, especially in terms of the magnitude of differences or relationships between variables.

Non-parametric tests are essential tools when the assumptions of parametric tests, such as normality and equal variances, are violated. These tests provide flexibility for analyzing non-normally distributed, ordinal, or categorical data, ensuring accurate and reliable results in situations where parametric methods fall short.

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

6. Descriptive Statistics

Descriptive statistics refers to statistical techniques that are used to summarize, organize, and describe data in a meaningful way. Rather than making inferences about a population from a sample (as inferential statistics does), descriptive statistics focus on giving a clear and concise picture of the data at hand. These methods help to simplify large amounts of data into an understandable form through various measures such as central tendency, variability, and distribution.

Types of Descriptive Statistics

Descriptive statistics can be divided into three main categories:

I. **Measures of Central Tendency**

These measures describe the "center" or "average" value of a data set and provide a single value around which the data tends to cluster.

1. **Mean**: The arithmetic average of the data set. Calculated by adding all the values together and dividing by the number of values.

$$\text{Mean} = \frac{\sum_{i=1}^{n} x_i}{n}$$

Where x_i are the data points and n is the number of data points.

2. **Median**: The middle value when the data is arranged in ascending or descending order. If the number of observations is even, the median is the average of the two middle values.

- For an **odd** number of observations: the median is the middle value when the data is sorted.
- For an **even** number of observations: the median is the average of the two middle values.

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

3. **Mode**: The most frequently occurring value in the data set. A data set may have one mode (unimodal), more than one mode (multimodal), or no mode at all.

- The mode is the value that appears most frequently in the data set. A data set may have more than one mode or no mode at all if all values are unique.

II. Measures of Variability (Dispersion)

These measures provide information about the spread or dispersion of the data. They help us understand how much the data points differ from each other and from the central tendency.

1. **Range**: The difference between the highest and lowest values in the data set.

$$\text{Range} = \text{Maximum value} - \text{Minimum value}$$

---

2. **Variance**: The average of the squared differences from the mean. It provides a measure of how much the values in the data set deviate from the mean.

$$\sigma^2 = \frac{\sum_{i=1}^{N}(x_i - \mu)^2}{N}$$

Where x_i are the data points, μ is the mean, and N is the number of data points.

3. **Standard Deviation**: The square root of the variance. It indicates the average distance of each data point from the mean and is expressed in the same units as the data.

$$\sigma = \sqrt{\sigma^2}$$

For a sample:

$$s = \sqrt{s^2}$$

---

4. **Interquartile Range (IQR)**: The range of the middle 50% of the data, calculated as the difference between the first quartile (Q1) and the third quartile (Q3).

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963


$$\text{IQR} = Q_3 - Q_1$$

Where Q_1 is the first quartile (25th percentile) and Q_3 is the third quartile (75th percentile).

III. Measures of Distribution Shape

These measures describe the overall shape of the data distribution, providing insights into the symmetry, skewness, and kurtosis of the data.

1. **Skewness**: Measures the asymmetry of the distribution.




A **positive skew** means the tail is on the right side of the distribution (right-skewed), and most data points are on the left.

A **negative skew** means the tail is on the left side of the distribution (left-skewed), and most data points are on the right.

Achiever's Hive

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

2. **Kurtosis**: Measures the "tailedness" of the distribution.



A distribution with **high kurtosis** has heavy tails, meaning there are more extreme values (outliers).

A distribution with **low kurtosis** has light tails, meaning fewer extreme values.

Visual Representation of Descriptive Statistics

Descriptive statistics are often visualized using graphical tools to give a clearer picture of the data distribution:

- **Histograms**: Used to represent the frequency distribution of continuous data. It shows how often each range of values occurs in a data set.
- **Boxplots**: Display the median, quartiles, and outliers, giving a concise summary of the data's central tendency and variability.
- **Bar Charts**: Used to represent categorical data by showing the frequency of each category.
- **Pie Charts**: Represent proportions of categorical data as slices of a circle.
- **Scatterplots**: Used to visualize the relationship between two quantitative variables, showing how one variable may affect or relate to the other.

Uses of Descriptive Statistics

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

I. **Summarizing Data**: Descriptive statistics help reduce large data sets into a manageable form by summarizing key characteristics, such as the average, spread, and distribution.
II. **Data Interpretation**: By providing a clear picture of the data, descriptive statistics allow researchers, decision-makers, and analysts to interpret the data effectively and make informed decisions.
III. **Exploratory Data Analysis (EDA)**: Before conducting inferential statistics, descriptive statistics help in understanding patterns, trends, and anomalies in the data.
IV. **Comparing Groups**: Measures like mean and standard deviation can be used to compare different groups or variables, allowing for easy identification of significant differences or similarities.

Descriptive statistics play a foundational role in data analysis by summarizing and simplifying large amounts of data into meaningful, easily interpretable figures. They are essential for understanding data distributions, central tendencies, and variability, and are widely used across various fields for initial data exploration and interpretation.

7. Inferential Statistics

Inferential statistics refers to the branch of statistics that allows researchers to make conclusions or inferences about a population based on data collected from a sample. While **descriptive statistics** summarizes the data, inferential statistics goes a step further by making predictions, generalizations, and decisions about the larger population from which the sample is drawn.

Inferential statistics often involve **hypothesis testing, estimation, and making predictions** using sample data. Since it's often impractical or impossible to collect data from every member of a population, inferential methods provide a way to draw conclusions from smaller, manageable datasets.

Key Concepts in Inferential Statistics

I. **Population and Sample**

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

Population: The entire set of individuals or data points that are the subject of study (e.g., all voters in a country).

Sample: A subset of the population selected for the actual analysis. The goal is to infer information about the population from the sample.

II. **Parameter vs. Statistic**

Parameter: A numerical value that describes a characteristic of the entire population (e.g., population mean, population variance).

Statistic: A numerical value that describes a characteristic of a sample (e.g., sample mean, sample variance). Statistics are used to estimate population parameters.

III. **Sampling Distribution**

Definition: The probability distribution of a given statistic based on a random sample. The central limit theorem states that the sampling distribution of the sample mean will be approximately normal if the sample size is sufficiently large, regardless of the population's distribution.

IV. **Hypothesis Testing**

- **Null Hypothesis (H_0)**: A statement of no effect or no difference, serving as the default assumption (e.g., "There is no difference between the means of two groups").
- **Alternative Hypothesis (H_1)**: A statement that contradicts the null hypothesis, proposing an effect or difference (e.g., "There is a significant difference between the means").
- **p-Value**: The probability of obtaining the observed sample results, or something more extreme, if the null hypothesis is true. If the p-value is below a certain threshold (commonly 0.05), the null hypothesis is rejected.
- **Type I Error (α)**: Rejecting the null hypothesis when it is true (a "false positive").
- **Type II Error (β)**: Failing to reject the null hypothesis when it is false (a "false negative").

V. **Confidence Intervals**

Definition: A range of values, derived from the sample, that is likely to contain the population parameter with a certain level of confidence (e.g., 95% confidence interval).

Interpretation: If you calculate a 95% confidence interval for a sample mean, you are 95% confident that the true population mean falls within that interval.

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

VI. **Estimation**

Point Estimation: A single value estimate of a population parameter (e.g., sample mean is used to estimate the population mean).

Interval Estimation: Provides a range of values (confidence interval) for the parameter.

Common Inferential Statistical Methods

I. **t-Test**: Used to compare the means of two groups and determine if the differences are statistically significant.

a) **Independent t-test**: Compares the means of two independent groups.
b) **Paired t-test**: Compares the means of two related groups (e.g., before and after treatment).

II. **Analysis of Variance (ANOVA)**: Used to compare the means of three or more groups. It determines if there are significant differences between the group means.

a) **One-Way ANOVA**: Tests for differences between groups based on one independent variable.
b) **Two-Way ANOVA**: Tests for differences considering two independent variables.

III. **Chi-Square Test**: Used to examine the relationship between categorical variables in a contingency table. It tests whether the observed frequencies differ significantly from expected frequencies. These are commonly used for testing the independence of two variables or the goodness of fit.

IV. **Regression Analysis**:

a) **Simple Linear Regression**: Assesses the relationship between two variables (one independent, one dependent) and predicts the value of the dependent variable based on the independent variable.
b) **Multiple Regression**: Involves two or more independent variables to predict a single dependent variable.

V. **Correlation**:

a) **Pearson's Correlation**: Measures the strength and direction of the linear relationship between two continuous variables.

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

b) **Spearman's Rank Correlation**: A non-parametric test used to measure the strength and direction of the relationship between two ranked variables.

Assumptions in Inferential Statistics

Random Sampling: The data should be collected through random sampling to avoid bias. Random sampling ensures that every member of the population has an equal chance of being selected.

Normal Distribution: Many inferential statistics methods, especially parametric tests like t-tests and ANOVA, assume that the data follows a normal distribution, or that the sampling distribution of the test statistic is approximately normal (especially for large samples, as per the Central Limit Theorem).

Independent Observations: The observations in the sample should be independent of each other, meaning that the value of one observation should not affect the value of another.

Equal Variance (Homoscedasticity): In some tests (e.g., ANOVA), it's assumed that the variances in different groups being compared are approximately equal.

Advantages of Inferential Statistics

I. **Generalization to Populations**: Inferential statistics allow researchers to make generalizations about a population based on sample data, making it feasible to study large populations without collecting data from every individual.
II. **Hypothesis Testing**: It provides a framework for testing hypotheses, helping researchers determine whether their findings are statistically significant.
III. **Precision with Confidence Intervals**: Confidence intervals provide an estimate of the population parameter, giving researchers an understanding of the precision of their estimates.
IV. **Decision-Making**: Inferential statistics are essential in fields like medicine, psychology, economics, and business, where making decisions based on data-driven inferences is critical.

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

Limitations of Inferential Statistics


I. **Dependence on Assumptions**: Inferential methods often rely on certain assumptions (e.g., normality, equal variance). When these assumptions are violated, the results can be inaccurate.
II. **Sampling Error**: Since inferential statistics are based on samples, there is always the risk of sampling error, meaning the sample may not perfectly represent the population.
III. **Generalizability Issues**: Inferences drawn from a sample are only as good as the sample itself. Non-random or biased sampling can lead to incorrect conclusions about the population.
IV. **Type I and Type II Errors**: There's always the risk of making errors in hypothesis testing, such as rejecting a true null hypothesis (Type I error) or failing to reject a false null hypothesis (Type II error).

Inferential statistics is a powerful tool for drawing conclusions about a population based on sample data. By using techniques such as hypothesis testing, confidence intervals, and regression, researchers can make informed predictions and decisions. However, the reliability of these methods depends on proper sampling and adherence to underlying assumptions.

8. Measures of central tendency

Measures of central tendency are statistical tools used to identify a central or typical value around which the data tends to cluster. They provide a summary of a data set by identifying the point at which the data is centered. The three most commonly used measures of central tendency are **mean**, **median**, and **mode**.

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963



1. Mean (Arithmetic Mean)

Definition: The mean is the sum of all the values in a data set divided by the number of values. It is often referred to as the "average."

Formula:

$$\text{Mean} = \frac{\sum_{i=1}^{n} x_i}{n}$$

Where:

- x_i represents each individual data point.
- n is the total number of data points.
- $\sum_{i=1}^{n} x_i$ is the sum of all data points.

Example: For the data set {2, 4, 6, 8, 10}, the mean is: (2+4+6+8+10) / 5 = 6

Advantages:

- Easy to calculate.
- Uses all data points, providing a good overall summary.

Disadvantages:

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

- Sensitive to extreme values (outliers), which can skew the mean.

2. Median

Definition: The median is the middle value in a data set when the values are arranged in ascending or descending order. If the number of data points is even, the median is the average of the two middle numbers.

How to Find:

- Arrange the data in order.
- Identify the middle value (if odd number of data points) or average the two middle values (if even number of data points).

Example:

- For the data set {3, 5, 7, 9, 11}, the median is **7**.
- For the data set {3, 5, 7, 9}, the median is **6.**

Advantages:

- Not affected by outliers or extreme values.
- Useful for skewed distributions.

Disadvantages:

- Does not take into account all values in the data set.
- Not as informative as the mean for symmetrical distributions.

3. Mode

Definition: The mode is the value that appears most frequently in a data set. A data set may have one mode (unimodal), more than one mode (bimodal or multimodal), or no mode at all if no number repeats.

How to Find: Identify the number(s) that occur most often in the data set.

Example: For the data set {1, 2, 2, 3, 4}, the mode is **2**.

For the data set {1, 1, 2, 3, 3, 4}, the data is bimodal, with modes **1** and **3**.

Advantages:

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

- Useful for categorical data where finding an average is not meaningful.
- Provides the most common value in the data set.

Disadvantages:

- May not exist in some data sets.
- Not useful for continuous data or large datasets with no repeating values.

Comparison of the Measures of Central Tendency

- **Mean** is most useful when the data is symmetrically distributed without outliers.
- **Median** is preferable when dealing with skewed data or when outliers are present, as it is less sensitive to extreme values.
- **Mode** is particularly useful for categorical data or identifying the most common value in a distribution.

Each measure of central tendency provides different insights into the data, and the best one to use depends on the nature of the data and the goal of the analysis. Often, using a combination of these measures offers the most complete understanding of the data distribution.

9. Measures of dispersion/ Measures of Variability

Measures of dispersion (also known as measures of variability or spread) describe how spread out the data points in a data set are. They provide insight into the variability of the data, showing how much the data values deviate from the central tendency (like the mean or median). The most common measures of dispersion are **range**, **variance**, **standard deviation**, and **interquartile range**.

1. Range

Definition: The range is the difference between the highest and lowest values in a data set. It gives a basic measure of the spread of data but does not provide information about the distribution of values between the extremes.

Formula:

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

$$\text{Range} = \text{Maximum value} - \text{Minimum value}$$

Example: For the data set {2, 4, 6, 8, 10}, the range is: 10 – 2 = 8

Advantages:

- Simple to calculate.
- Provides a quick sense of data spread.

Disadvantages:

- Sensitive to outliers (extreme values).
- Does not consider how data is distributed between the highest and lowest points.

2. Variance

Definition: Variance measures the average squared deviation of each data point from the mean. It provides a sense of how much the data varies around the mean.

Formula:

$$\sigma^2 = \frac{\sum_{i=1}^{N}(x_i - \mu)^2}{N}$$

Where x_i are the data points, μ is the mean, and N is the number of data points.

Example: For the data set {2, 4, 6, 8, 10}, with a mean of 6:

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

$$s^2 = \frac{\sum_{i=1}^{n}(x_i - \bar{x})^2}{n-1}$$

Step-by-step calculation:

1. Find the squared deviations from the mean for each data point:
   - $(2-6)^2 = (-4)^2 = 16$
   - $(4-6)^2 = (-2)^2 = 4$
   - $(6-6)^2 = 0^2 = 0$
   - $(8-6)^2 = 2^2 = 4$
   - $(10-6)^2 = 4^2 = 16$
2. Sum of squared deviations:
   $16 + 4 + 0 + 4 + 16 = 40$
3. Number of data points (n) is 5, so:
   $s^2 = \frac{40}{5-1} = \frac{40}{4} = 10$

Variance:

$s^2 = 10$

Advantages:

- Takes into account all data points.
- Important in many statistical applications (e.g., in the calculation of standard deviation).

Disadvantages:

- Results in squared units, making it harder to interpret in terms of the original data scale.

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

3. Standard Deviation

Definition: The standard deviation is the square root of the variance, bringing the measure back to the same units as the data. It shows the average deviation of data points from the mean and provides a more interpretable measure of dispersion than variance.

Formula:

3. **Standard Deviation**:

$\sigma = \sqrt{\sigma^2}$

For a sample:

$s = \sqrt{s^2}$

Example: For the data set {2, 4, 6, 8, 10}, the standard deviation is:

Variance:

$s^2 = 10$

Standard Deviation (Sample):

$s = \sqrt{s^2} = \sqrt{10}$

Calculating $\sqrt{10}$:

$s \approx 3.16$

Standard deviation:

$s \approx 3.16$

So, the standard deviation for the data set $2, 4, 6, 8, 10$ is approximately **3.16**.

Advantages:

- Widely used and easy to interpret in terms of the data's original units.
- More robust and informative than the range, as it considers all data points.

Disadvantages:

- Sensitive to outliers.

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

4. Interquartile Range (IQR)

Definition: The interquartile range is the range of the middle 50% of the data, calculated as the difference between the first quartile (Q1) and the third quartile (Q3). It is particularly useful for skewed distributions or data with outliers.

Formula:

$$\text{IQR} = Q_3 - Q_1$$

Q1 (First Quartile): The 25th percentile (the value below which 25% of the data lies).

Q3 (Third Quartile): The 75th percentile (the value below which 75% of the data lies).

Example: For the data set {1, 2, 3, 4, 5, 6, 7, 8, 9, 10}, Q1 = 3, and Q3 = 8, so: IQR is 8-3 = 5

Advantages:

- Not affected by outliers or extreme values.
- Provides a measure of spread for the central portion of the data.

Disadvantages:

- Ignores the dispersion in the tails of the distribution.

5. Mean Absolute Deviation (MAD)

Definition: The mean absolute deviation is the average of the absolute differences between each data point and the mean. It measures the spread of the data in a more intuitive way than variance or standard deviation.

Formula:

$$\text{MAD} = \frac{\sum_{i=1}^{n} |x_i - \bar{x}|}{n}$$

Where:

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

- x_i represents each data point.
- $\bar{x}$ is the mean of the data set.
- n is the number of data points.
- $|x_i - \bar{x}|$ is the absolute deviation of each data point from the mean.

Advantages:

- Simple and easy to interpret.
- Less affected by extreme values than variance or standard deviation.

Disadvantages:

- Not as commonly used in inferential statistics as variance and standard deviation.

Comparison of the Measures of Dispersion

- **Range** is the simplest and gives a quick sense of the spread but is not informative for most distributions.
- **Variance** and **standard deviation** provide more detailed information by considering all data points, but they are sensitive to outliers.
- **IQR** is robust to outliers and provides a good summary for skewed data.
- **MAD** is another intuitive measure that avoids the squaring of deviations but is less common.

Each measure of dispersion provides different insights depending on the nature of the data and the analysis's goals. Often, a combination of these measures is used to fully understand the variability of the data.

10. Partial and Part Correlation

Partial Correlation

Partial correlation measures the strength and direction of the linear relationship between two variables, while controlling for the effect of one or more additional variables. Essentially, it tells you how much of the correlation between two variables is unique, after accounting for the influence of other variables.

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

Key Points of Partial Correlation:

I. Controls for Third Variables: Partial correlation controls the influence of third (or more) variables. For example, if you want to understand the relationship between variables X and Y while removing the influence of variable Z, you would use partial correlation.

II. Range: The value of partial correlation ranges between -1 and 1.

- A value close to 1 indicates a strong positive relationship after controlling for other variables.
- A value close to -1 indicates a strong negative relationship after controlling for other variables.
- A value around 0 indicates no linear relationship after controlling for other variables.

III. Formula:

If we are calculating the partial correlation between X and Y controlling for Z, the formula is:

The formula for the partial correlation between two variables X and Y, controlling for a third variable Z, is:

$$r_{XY \cdot Z} = \frac{r_{XY} - r_{XZ} \cdot r_{YZ}}{\sqrt{(1 - r_{XZ}^2)(1 - r_{YZ}^2)}}$$

Where:

- r_{XY} is the Pearson correlation coefficient between X and Y.
- r_{XZ} is the Pearson correlation coefficient between X and Z.
- r_{YZ} is the Pearson correlation coefficient between Y and Z.

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

Explanation:

- **Numerator**: $r_{XY} - r_{XZ} \cdot r_{YZ}$ represents the difference between the raw correlation of X and Y and the product of their correlations with Z.
- **Denominator**: $\sqrt{(1 - r_{XZ}^2)(1 - r_{YZ}^2)}$ normalizes the result by adjusting for the strength of the correlation between X and Z, and Y and Z.

This formula helps determine the correlation between X and Y after removing the linear effects of Z on both variables.

Some practical examples of partial correlation in different fields:

1. **Psychology: Examining Stress and Academic Performance**

Suppose researchers are studying the correlation between **stress** (X) and **academic performance** (Y) in students. **Sleep quality** (Z) could be a confounding factor because poor sleep might increase stress and decrease academic performance.

By calculating the partial correlation between stress and academic performance while controlling for sleep quality, the researchers can better understand if the relationship between stress and performance is due to stress itself or primarily due to sleep issues.

Interpretation: If the partial correlation remains strong, it suggests that stress independently impacts academic performance, regardless of sleep quality.

2. **Economics: Income and Health Outcomes**

Economists might examine the correlation between **income** (X) and **health outcomes** (Y), where **education level** (Z) could be a confounding variable. Higher education often leads to higher income and also better health awareness, which could influence health outcomes.

By controlling for education level, the partial correlation between income and health outcomes isolates the effect of income alone, providing insight into whether financial resources, independent of education, play a significant role in health.

Interpretation: A strong partial correlation here would imply that income, beyond education level, has a notable effect on health.

3. **Healthcare: Medication Adherence and Patient Recovery**

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

In a clinical study, researchers want to explore the relationship between **medication adherence** (X) and **recovery rate** (Y) in patients. **Severity of illness** (Z) could act as a confounding factor because patients with more severe conditions might adhere more closely to medication regimens and may also recover more slowly due to the severity itself.

Controlling for severity, a partial correlation analysis could reveal if medication adherence significantly affects recovery rates, independent of illness severity.

Interpretation: If the partial correlation is high, it would indicate that adherence to medication has a direct impact on recovery, beyond the influence of illness severity.

4. **Education: Technology Use and Test Scores**

An educational researcher might study the relationship between **technology use** (X) and **test scores** (Y) among students. **Parental involvement** (Z) may influence both, as involved parents might limit technology use and simultaneously encourage academic focus, which could improve scores.

By isolating the impact of parental involvement, the researcher can see whether technology use independently correlates with test scores, leading to insights about the role of technology in learning.

Interpretation: A near-zero partial correlation might suggest that the initial correlation between technology use and test scores was due to parental involvement rather than technology itself.

In each of these examples, the **partial correlation helps to clarify relationships by isolating the variable of interest** from others, giving a clearer understanding of causal or direct associations in complex data.

Part Correlation (or Semi partial Correlation)

Part correlation (also known as semi partial correlation) measures the relationship between two variables while controlling for the effect of one or more variables, but only on one of the variables being correlated. In other words, the effect of the control variable is removed only from one of the variables, rather than both.

Key Points of Part Correlation:

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

I. Partial Out the Effect: It only removes the influence of the control variable from one of the variables in the pair, rather than both, unlike partial correlation which adjusts for the control variable in both variables.

II. Range: The value of part correlation also ranges between -1 and 1.

- A value near 1 indicates a strong positive relationship after controlling for other variables on one variable.
- A value near -1 indicates a strong negative relationship.
- A value close to 0 means there's little to no linear relationship after accounting for the control variable in one variable.

Formula:

If we want to calculate the part correlation between X and Y, controlling for Z in X, the formula is:

The formula for **part correlation** (also known as semi-partial correlation) between two variables X and Y while controlling for a third variable Z is given by:

$$r_{X(Y \cdot Z)} = \frac{r_{XY} - r_{XZ} \cdot r_{YZ}}{\sqrt{1 - r_{XZ}^2}}$$

Where:

- r_{XY} is the Pearson correlation coefficient between X and Y.
- r_{XZ} is the Pearson correlation coefficient between X and Z.
- r_{YZ} is the Pearson correlation coefficient between Y and Z.

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

Explanation:

- **Numerator**: $r_{XY} - r_{XZ} \cdot r_{YZ}$ represents the difference between the raw correlation of X and Y and the product of their correlations with Z.
- **Denominator**: $\sqrt{(1 - r_{XZ}^2)(1 - r_{YZ}^2)}$ normalizes the result by adjusting for the strength of the correlation between X and Z, and Y and Z.

This formula helps determine the correlation between X and Y after removing the linear effects of Z on both variables.

Applications: Part correlation is typically used when researchers want to measure the unique contribution of a variable in explaining a relationship, especially in regression analysis. For example, it is used to determine the unique effect of one independent variable on the dependent variable while holding other variables constant.

Difference Between Partial and Part Correlation

Aspect	Partial Correlation	Part (Semipartial) Correlation
Definition	Measures the correlation between two variables while controlling for the effect of one or more additional variables on both.	Measures the correlation between two variables while controlling for the effect of one or more additional variables on only one variable.
Adjustment	Adjusts for the effect of control variables on both variables being correlated.	Adjusts for the effect of control variables on only one of the variables.
Use	To understand the unique relationship between two variables when the effect of other variables is removed from both.	To understand the unique effect of one variable on another while controlling for the effect of other variables on only one of them.
Complexity	More complex as it controls for confounders on both variables.	Simpler as it controls for confounders on just one variable.

Both partial and part correlation are valuable tools in understanding relationships between variables in the presence of confounders or control variables, with specific uses depending on the nature of the analysis.

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

11. Point-Biserial Correlation vs Phi Coefficient (Φ)

Point-Biserial Correlation

The point-biserial correlation is a special case of the Pearson correlation coefficient used when one variable is continuous (e.g., height, income) and the other is dichotomous (binary), meaning it has only two possible values (e.g., gender, pass/fail).

Key Characteristics of Point-Biserial Correlation:

i. Binary and Continuous Variables: It is used to measure the strength and direction of the association between a continuous variable and a dichotomous variable.
ii. Range: Like the Pearson correlation, the point-biserial correlation ranges between -1 and 1.

- A positive value indicates that higher values of the continuous variable tend to occur with one category of the binary variable.
- A negative value indicates that higher values of the continuous variable tend to occur with the other category.
- A value near 0 suggests little or no linear relationship between the two variables.

Formula:

The **point biserial correlation** (often denoted as r_{pb}) is used to measure the relationship between a continuous variable and a binary variable. The formula for point biserial correlation is:

$$r_{pb} = \frac{M_1 - M_0}{s} \cdot \sqrt{\frac{n_1 n_0}{n^2}}$$

Where:

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

- M_1 is the mean of the continuous variable for the group coded as 1 (the presence of the characteristic).
- M_0 is the mean of the continuous variable for the group coded as 0 (the absence of the characteristic).
- s is the standard deviation of the continuous variable.
- n_1 is the number of observations in the group coded as 1.
- n_0 is the number of observations in the group coded as 0.
- n is the total number of observations ($n = n_1 + n_0$).

Applications:

Education: To examine the relationship between exam scores (continuous variable) and pass/fail status (dichotomous).

Psychology: To study the relationship between an outcome variable like test anxiety (continuous) and group membership (e.g., treatment/no treatment).

Phi Coefficient (Φ)

The phi coefficient is a measure of association used when both variables are binary (dichotomous). It is a type of correlation coefficient used to assess the relationship between two dichotomous variables.

Key Characteristics of the Phi Coefficient:

I. Binary-Binary Correlation: It is used to determine the strength and direction of the relationship between two binary variables (e.g., gender and smoking status: male/female, smoker/non-smoker).

II. Range: The phi coefficient also ranges between -1 and 1.

- A value of 1 indicates a perfect positive association.
- A value of -1 indicates a perfect negative association.
- A value of 0 indicates no association between the variables.

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

Formula:

The **Phi coefficient** (often denoted as ϕ) is a measure of association for two binary variables. It is calculated as follows:

$$\phi = \frac{n_{11}n_{00} - n_{10}n_{01}}{\sqrt{(n_{10} + n_{11})(n_{01} + n_{00})(n_{10} + n_{01})(n_{11} + n_{00})}}$$

Where:

- n_{11}: Number of cases where both variables are 1 (True, True).
- n_{00}: Number of cases where both variables are 0 (False, False).
- n_{10}: Number of cases where the first variable is 1 and the second is 0 (True, False).
- n_{01}: Number of cases where the first variable is 0 and the second is 1 (False, True).

Applications:

Medical Research: To examine the relationship between a disease diagnosis (yes/no) and the presence of a particular risk factor (yes/no).

Marketing: To study the association between a customer buying a product (yes/no) and their response to a marketing campaign (yes/no).

Comparison: Point-Biserial vs. Phi Coefficient

The **Point-Biserial correlation** and the **Phi coefficient** are both measures used to assess the relationship between variables, specifically when dealing with **binary variables**. However, they are used in slightly different contexts and have unique interpretations:

1. Point-Biserial Correlation

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

Definition: The point-biserial correlation coefficient measures the strength and direction of the association between a continuous variable and a binary (dichotomous) variable.

Use Case: This is commonly used when you have one binary variable (e.g., **gender** with values 0 = female, 1 = male) and one continuous variable (e.g., **test scores**).

Calculation: It's similar to the Pearson correlation but adjusted for one variable being dichotomous. It calculates how much the mean of the continuous variable differs for the two groups defined by the binary variable.

Interpretation: A higher positive point-biserial correlation implies that one group tends to have higher values on the continuous variable than the other group. Values range from -1 to +1, with +1 indicating a strong positive association and -1 indicating a strong negative association.

Example: If you want to see the relationship between **exercise** (yes = 1, no = 0) and **blood pressure level** (a continuous measure), the point-biserial correlation can help you understand how blood pressure varies between people who exercise and those who do not.

2. Phi Coefficient

Definition: The phi coefficient measures the strength of association between two binary variables.

Use Case: This measure is used when both variables are binary. For example, if you have data on **smoking status** (yes = 1, no = 0) and **lung disease presence** (yes = 1, no = 0), the phi coefficient would be appropriate.

Calculation: The phi coefficient is essentially the Pearson correlation coefficient applied to two binary variables, and it uses a **2x2 contingency table** to compute the association.

Interpretation: Like other correlation coefficients, phi ranges from -1 to +1. A positive phi indicates that both variables are likely to have the same value (either both 1 or both 0), while a negative phi suggests they are inversely related.

Example: In a study of job applicants, you could use the phi coefficient to examine the association between **interview performance** (pass = 1, fail = 0) and **prior work experience** (yes = 1, no = 0).

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

Key Differences

Aspect	Point-Biserial Correlation	Phi Coefficient
Variables	One binary and one continuous	Two binary variables
Interpretation	Measures how a continuous variable differs across two binary groups	Measures association between two binary outcomes
Calculation	Similar to Pearson, adjusted for dichotomy	Pearson correlation on binary variables with 2x2 contingency table
Range of Values	-1 to +1	-1 to +1
Typical Use Case	Relationship between a category and continuous score	Association between two categorical events

Both the point-biserial correlation and the phi coefficient are valuable in different research contexts, depending on whether the variables involved are continuous, dichotomous, or both.

12. Point and Interval Estimation

Point and Interval Estimation are two approaches in statistics for estimating unknown population parameters (such as a population mean or proportion) based on sample data. These methods allow statisticians and researchers to make inferences about a larger population from a smaller sample.

1. Point Estimation

Definition: Point estimation involves using a single value, calculated from sample data, as an estimate of a population parameter. It provides a "best guess" or single estimate for the parameter.

Example: If we want to estimate the average height of all adults in a city, we might collect a sample and calculate the sample mean height. This sample mean serves as the **point estimate** of the population mean height.

Common Point Estimators:

- **Mean (M)**: The sample mean is a point estimate of the population mean.
- **Proportion ($\hat{p}$)**: The sample proportion is a point estimate of the population proportion.
- **Variance (s^2)**: The sample variance is a point estimate of the population variance.

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

Properties of a Good Point Estimator:

- **Unbiasedness**: The estimator's expected value should be equal to the true population parameter.
- **Consistency**: As the sample size increases, the estimator should converge to the true parameter value.
- **Efficiency**: Among all unbiased estimators, it has the smallest variance.
- **Sufficiency**: Uses all information in the sample to estimate the parameter.
- **Limitations**: Point estimates provide no information about the reliability or precision of the estimate, as they lack a margin of error or range of values.

2. Interval Estimation

Definition: Interval estimation provides a range (or interval) within which the population parameter is expected to lie, along with an associated confidence level. Unlike point estimation, interval estimation considers the variability inherent in sample data and offers a more realistic representation of uncertainty.

Confidence Interval: A confidence interval is a type of interval estimate that includes a lower and upper bound, indicating where the true population parameter is likely to be found with a specified level of confidence (usually 90%, 95%, or 99%).

Example: A 95% confidence interval for the mean height of adults in a city might be 65 to 70 inches, meaning we are 95% confident that the true average height lies within this range.

Formula for Confidence Interval for a Population Mean (when population standard deviation is known):

$$\bar{x} \pm Z\frac{\sigma}{\sqrt{n}}$$

- $\bar{x}$ = sample mean
- Z = Z-score corresponding to the desired confidence level
- σ = population standard deviation (or sample standard deviation if σ is unknown)
- n = sample size

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

If the population standard deviation is unknown and the sample size is small, the **t-distribution** is used instead of the Z-distribution.

Interpretation of Confidence Intervals: A 95% confidence interval does not mean there is a 95% chance that the interval contains the true parameter. Instead, it means that if we were to take many random samples and compute a confidence interval for each sample, 95% of those intervals would contain the true parameter.

Factors Affecting Confidence Interval Width:

- **Sample Size (n)**: Larger sample sizes result in narrower confidence intervals, as they provide more information about the population.
- **Confidence Level**: Higher confidence levels (e.g., 99% vs. 95%) result in wider intervals, as they require more certainty that the interval includes the population parameter.
- **Standard Deviation**: Greater variability in the data leads to wider intervals.

Key Differences between Point and Interval Estimation

Characteristic	Point Estimation	Interval Estimation
Output	Single value (point)	Range of values (interval)
Information Provided	No indication of accuracy	Indicates accuracy with a confidence level
Uncertainty	Lacks an error margin	Includes margin of error
Usefulness	Simpler and quicker	More informative and reliable
Example	Sample mean as an estimate of population mean	95% confidence interval for population mean

Advantages and Disadvantages

Point Estimation:

- **Advantages**: Simple, provides a quick estimate of the population parameter.
- **Disadvantages**: Lacks a measure of accuracy, potentially misleading if viewed without context.

Interval Estimation:

- **Advantages**: Provides a range of plausible values, incorporates sampling variability, and is more informative.

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

- **Disadvantages**: More complex to calculate and interpret, especially if the confidence level or sample size is misinterpreted.

Practical Applications

In real-world applications, interval estimation is preferred when making inferences because it gives a clearer picture of the uncertainty involved. For instance, in medical research, a confidence interval for the effectiveness of a new drug provides more information about its likely effect size and the reliability of the results, compared to a single-point estimate of its effectiveness.

In summary:

- **Point estimation** offers a single best guess of a population parameter.
- **Interval estimation** provides a range within which the parameter is expected to lie, with an associated confidence level, adding reliability and depth to statistical conclusions.

13. Type I and Type II Errors

Type I and Type II errors are central concepts in hypothesis testing that describe the possible mistakes researchers can make when evaluating data.

1. Type I Error:

Also called a **"false positive"**. This occurs when the test **incorrectly rejects a true null hypothesis**. In simpler terms, it's concluding there is an effect or difference when in reality, there isn't one.

The probability of making a Type I error is denoted by **alpha (α)**, often set at 0.05 or 5%, meaning there is a 5% risk of incorrectly finding a significant effect. *Example*: Concluding that a new drug works when it actually doesn't.

Context: In statistical testing, a Type I error means that you conclude there is a statistically significant effect or relationship when there is actually none.

Example: Suppose a company tests a new feature, expecting it to increase sales. If they conclude it does improve sales (by rejecting the null hypothesis that it has no effect) when, in fact, it doesn't, they have made a Type I error.

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

Implications: Type I errors are problematic because they lead to incorrect acceptance of new hypotheses. In critical areas like medicine, it could lead to adopting an ineffective treatment as a standard of care.

Control Mechanism: The significance level (alpha, α) is set to control the probability of a Type I error. Commonly, **α = 0.05** is used, which means there's a 5% chance of incorrectly rejecting the null hypothesis. However, this threshold can be made stricter (e.g., α = 0.01) in cases where a false positive could have serious implications, such as in clinical trials.

2. Type II Error:

Also called a **"false negative"**. This happens when the test **fails to reject a false null hypothesis**. In other words, it means concluding there is no effect or difference when there actually is one.

The probability of making a Type II error is represented by **beta (β)**, and **power (1 - β)** is the likelihood of correctly rejecting a false null hypothesis. *Example*: Concluding that a new drug has no effect when it actually does.

Context: A Type II error occurs when a test fails to identify a true effect, which means that you fail to reject the null hypothesis even though it is false.

Example: In a clinical setting, if a new drug truly improves health outcomes but the test fails to show statistical significance, leading to the drug's rejection, a Type II error has occurred.

Implications: Type II errors can lead to missed opportunities, as real effects go unrecognized. In business, it might mean rejecting a new process that could actually increase efficiency, or in public health, missing early signs of a disease outbreak.

Control Mechanism: The probability of a Type II error is denoted by **beta (β)**, and **power (1 - β)** represents the probability of correctly rejecting a false null hypothesis. Typically, researchers aim for a power of **80%** or higher, meaning there is an 80% chance of detecting a true effect if it exists. Larger sample sizes or more sensitive tests can help reduce Type II errors by increasing statistical power.

Key Differences and Trade-offs

There's a natural trade-off between Type I and Type II errors:

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

Reducing Type I Error: If you lower α to be very strict (say, 0.01), you're less likely to make a Type I error. However, this increases the likelihood of a Type II error because a smaller α makes it harder to reject the null hypothesis, potentially leading to missed true effects.

Reducing Type II Error: Increasing the power of a test (e.g., by increasing the sample size) reduces the chance of a Type II error, but may require more resources. In some fields, this is preferred because missed effects (Type II errors) are considered more harmful, as in clinical trials where a missed treatment effect could have severe consequences.

Examples in Practical Scenarios

I. **Medical Testing**:

- *Type I Error*: Concluding a new drug is effective when it isn't.
- *Type II Error*: Concluding a new drug is not effective when it actually is.

II. **Quality Control in Manufacturing**:

- *Type I Error*: Flagging a defect in a batch of products that is actually fine, leading to unnecessary re-inspections or recalls.
- *Type II Error*: Missing a defect in a batch that is actually flawed, leading to defective products reaching customers.

III. **A/B Testing in Marketing**:

- *Type I Error*: Concluding that a new ad campaign is better than the old one when it's not, leading to wasted advertising budgets.
- *Type II Error*: Concluding that the new ad campaign isn't effective when it actually is, missing out on potentially higher sales.

Minimizing Type I and Type II Errors

In research, a balanced approach is taken to minimize both types of errors:

I. **Set Appropriate Significance Levels**: Adjusting α based on the field's tolerance for false positives.

II. **Increase Sample Size**: Larger sample sizes reduce variability, increasing power and thus reducing the likelihood of Type II errors.

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

III. **Use One-Tailed vs. Two-Tailed Tests**: In cases where the direction of the effect is known, one-tailed tests can reduce the chance of Type II errors by focusing on a single direction of interest.

IV. **Replication and Cross-Validation**: Performing multiple tests or replications helps confirm results and reduce the likelihood of both error types.

14. Levels of Significance

In statistical analysis, the "level of significance" (often denoted by the Greek letter \alpha) is a threshold used to determine whether a hypothesis test's result is statistically significant. This level helps assess whether observed data deviate enough from the null hypothesis (the hypothesis of no effect) to reject it in favor of the alternative hypothesis.

Type I and Type II Errors in Context

Type I Error (False Positive): This occurs when a true null hypothesis is incorrectly rejected. The significance level \alpha directly controls the likelihood of making a Type I error. For instance, with \alpha = 0.05, there's a 5% chance of mistakenly concluding there's an effect when there is none.

Type II Error (False Negative): This error happens when the null hypothesis is false, but we fail to reject it. The probability of a Type II error is denoted by \beta, and 1 - \beta represents the **power** of a test, or its ability to detect an effect when one truly exists.

Common levels of significance are:

1. **0.05 Level** (5%): This is the most commonly used level. It implies a 5% risk of concluding that an effect exists when it actually doesn't. Results are considered statistically significant if the p-value is less than 0.05.

2. **0.01 Level** (1%): A more stringent level, often used in fields requiring greater certainty (like medical research). Here, there's only a 1% chance of rejecting the null hypothesis incorrectly.

3. **0.10 Level** (10%): Less stringent and occasionally used in exploratory research where preliminary insights are the goal. A 10% risk of a Type I error is accepted.


Interpreting Levels of Significance

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

A lower level of significance (like 0.01) reduces the likelihood of a Type I error (rejecting a true null hypothesis) but increases the chance of a Type II error (failing to reject a false null hypothesis). Researchers choose the level of significance based on factors such as the field of study, the potential consequences of errors, and the specific hypotheses being tested.

15. Skewness

Skewness is a measure of the asymmetry in a data distribution, indicating whether the data points are symmetrically distributed or skewed to one side. In a perfectly symmetrical distribution, the skewness is zero, and the data follow a bell-shaped curve (like the normal distribution). When a distribution is skewed, it means that the data are pulled more toward one tail, making one side of the distribution longer than the other.



Types of Skewness

1. **Positive Skewness (Right Skew)**

In a positively skewed distribution, the tail on the right side (higher values) is longer. This indicates that a few unusually high values are pulling the average to the right.

Characteristics:

- The mean is typically greater than the median.
- The majority of the data points lie to the left of the mean, creating a "tail" to the right.

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

Example: Income distribution in many societies is positively skewed, as most people earn lower to average incomes, but a few high-income individuals push the distribution to the right.

2. **Negative Skewness (Left Skew)**

In a negatively skewed distribution, the tail on the left side (lower values) is longer, suggesting that a few unusually low values are pulling the average to the left.

Characteristics:

- The mean is typically less than the median.
- Most data points are concentrated on the right, with a "tail" extending to the left.

Example: Scores on an easy test often show negative skewness, with most students scoring high and only a few scoring low.

3. **Zero Skewness (Symmetrical Distribution)**

When skewness is zero or close to zero, the distribution is symmetric, resembling a normal distribution (bell-shaped curve).

Characteristics:

- The mean and median are approximately equal.
- Data points are evenly distributed around the center.

Measuring Skewness

Formula for Skewness: One common formula for sample skewness (where n is the sample size, $\bar{X}$ is the sample mean, and s is the sample standard deviation) is:

Sample Skewness Formula:

$$\text{Skewness} = \frac{n}{(n-1)(n-2)} \sum_{i=1}^{n} \left(\frac{x_i - \bar{x}}{s} \right)^3$$

Where:

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

- n is the number of data points.
- x_i represents each data point.
- $\bar{x}$ is the mean of the data.
- s is the sample standard deviation.

Interpretation:

Skewness < 0: Negative skewness (left skew).

Skewness = 0: Symmetrical distribution.

Skewness > 0: Positive skewness (right skew).

Effects of Skewness on Data Analysis

I. **Influence on Central Tendency**: Skewness affects the relationship between the mean and median. In positively skewed data, the mean is usually higher than the median; in negatively skewed data, the mean is typically lower.
II. **Implications for Hypothesis Testing**: Many statistical tests assume a normal distribution. High skewness can violate these assumptions, making the test results less reliable.
III. **Outliers and Data Interpretation**: Skewness often indicates outliers or extreme values, which might be meaningful for the analysis or suggest the need for transformation to better fit statistical models.

Transforming Data to Address Skewness

In cases where skewness distorts analysis, data transformations like the **logarithmic**, **square root**, or **Box-Cox** transformation can help normalize the distribution. These methods adjust skewed data toward a more symmetrical shape, making it easier to apply statistical techniques that assume normality.

Practical Examples

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

Finance: Stock returns often show positive skewness, as occasional large gains create a "tail" on the right.

Environmental Science: Measurements of pollutants can exhibit positive skewness, as high pollution events are rare but can create a long right tail.

Skewness offers insights into the nature of the data, helping to identify potential anomalies and guiding decisions about which statistical methods are most appropriate.

16. Hypothesis Testing

Hypothesis testing is a statistical method used to make decisions about a population parameter based on sample data. It helps determine whether the observed data supports a specific hypothesis or whether the difference is due to chance.

The process involves the following steps:

1. **State the Null Hypothesis**: This is a statement of no effect or no difference. It assumes that any observed effect is due to random chance.

2. **State the Alternative Hypothesis**: This contradicts the null hypothesis and reflects the effect or difference we are testing for.

3. **Select a Significance Level**: This is the probability of rejecting the null hypothesis when it is true. Common values for are 0.05 or 0.01.

4. **Collect Data and Compute the Test Statistic**: A test statistic (like -score, -statistic, etc.) is calculated based on the sample data to compare against a critical value.

5. **Make a Decision**:

- **Reject** if the test statistic falls in the critical region or the -value is less than .
- **Fail to reject** if the test statistic is within the acceptable range or the -value is greater than .

Errors in Hypothesis Testing

In hypothesis testing, two types of errors can occur due to uncertainty and randomness in the sample data.

1. **Type I Error** :

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

Definition: Rejecting the null hypothesis when it is actually true.

Example: Concluding that a new drug is effective when it actually isn't.

If , there is a 5% chance of incorrectly rejecting the true null hypothesis.

2. **Type II Error** :

Definition: Failing to reject the null hypothesis when the alternative hypothesis () is true.

Example: Concluding that a new drug is not effective when it actually is.

The **power** of a test is , representing the probability of correctly rejecting a false null hypothesis.

Summary of Possible Outcomes in Hypothesis Testing

Actual Situation	Test Decision	Outcome	Description
Null Hypothesis is True	Reject Null Hypothesis	**Type I Error**	Incorrectly concluding there is an effect when there isn't (false positive). The probability of this error is the significance level, α.
Null Hypothesis is True	Fail to Reject Null	**Correct Decision**	Correctly concluding there is no effect (true negative).
Null Hypothesis is False	Reject Null Hypothesis	**Correct Decision**	Correctly concluding there is an effect (true positive). This is the desired outcome, representing the test's power (1 - β).
Null Hypothesis is False	Fail to Reject Null	**Type II Error**	Incorrectly concluding there is no effect when there is one (false negative). The probability of this error is denoted by β.

Hypothesis testing provides a structured way to make inferences, but the possibility of errors always exists, so significance levels and power calculations are crucial to minimizing incorrect decisions.

17. Standard Error

Standard Error (SE) is a measure of the variability or dispersion of a sample statistic (such as the sample mean) around the true population parameter. It essentially quantifies how much the sample statistic is expected to vary from one sample to another due to random

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

sampling. It's a way to quantify the uncertainty in the estimate of a population parameter. In the context of linear regression, standard errors are used for both the regression coefficients and residuals.

Definition: The standard error represents the standard deviation of the sampling distribution of a statistic. When we calculate a statistic (like a mean or proportion) from a sample, the standard error gives us an idea of how much that statistic would vary if we took multiple samples.

Formula for Standard Error of the Mean:

For a sample mean, the standard error is given by:

The **standard error of the mean (SEM)** is calculated using the following formula:

$$\mathrm{SEM} = \frac{s}{\sqrt{n}}$$

Where:

- s is the sample standard deviation.
- n is the number of observations in the sample.

Interpretation:

The standard error tells us how much the sample mean (or other statistic) is expected to deviate from the actual population mean.

A smaller standard error indicates that the sample mean is likely closer to the true population mean, which generally happens when the sample size is larger or when there is less variability in the data.

Standard Error vs. Standard Deviation:

Standard Deviation (SD): Measures the dispersion of individual data points around the mean of the sample or population.

Standard Error (SE): Measures the dispersion of sample means around the population mean. SE decreases as sample size n increases, while SD is unaffected by sample size.

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

Applications:

Confidence Intervals: The standard error is used to construct confidence intervals. This interval provides a range of values that is likely to contain the population mean.

Hypothesis Testing: Standard error plays a role in calculating test statistics (such as the t-statistic) to determine if observed differences are statistically significant.

Summary

The standard error is crucial in inferential statistics because it helps quantify the precision of sample estimates, guiding interpretations of how close a sample statistic (like a mean) might be to the actual population parameter.

18. Steps in Computation of Regression

The steps involved in the computation of regression, particularly linear regression, include:

1. Gather and Organize Data:

- Collect data for the dependent variable (Y) and the independent variable (X).
- Ensure the data is clean and appropriately formatted for analysis.

2. Calculate Key Statistics:

- Compute the **mean** of X ($\bar{X}$) and Y ($\bar{Y}$).
- Compute the **sum of squares** for X (SS_X) and the **sum of products** of X and Y (SP).

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

Formulas:

$$\mathrm{SS}_X = \sum (X_i - \bar{X})^2$$

$$\mathrm{SP} = \sum (X_i - \bar{X})(Y_i - \bar{Y})$$

3. Calculate the Regression Coefficients:

- Compute the **slope** (b_1) and **intercept** (b_0) of the regression line using:

$$b_1 = \frac{\mathrm{SP}}{\mathrm{SS}_X}$$

$$b_0 = \bar{Y} - b_1 \bar{X}$$

4. Form the Regression Equation:

- Combine the coefficients to write the equation of the line:

$$Y = b_0 + b_1 X$$

5. Evaluate the Fit of the Model:

- Compute the **coefficient of determination** (R^2) to assess how well the regression line fits the data:

$$R^2 = \frac{\mathrm{SSRegression}}{\mathrm{SSTotal}}$$

Where:

- $\mathrm{SS}_{\mathrm{Regression}}$ is the explained variation.
- $\mathrm{SS}_{\mathrm{Total}}$ is the total variation in Y.

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

6. Check Assumptions: Verify assumptions of linear regression such as linearity, independence, homoscedasticity (constant variance of errors), and normality of residuals.

7. Make Predictions: Use the regression equation to predict values of for given values of .

8. Interpret the Results: Analyze the coefficients to understand the relationship between and , and determine if the relationship is statistically significant using hypothesis tests (e.g., -test for).

These steps outline the process for simple linear regression. For multiple regression, similar steps are followed, but with calculations extended to handle multiple independent variables.

19. Interactional effect

"Interactional effect" refers to the phenomenon where the impact of one variable on an outcome depends on the level of another variable. In other words, the combined influence of two variables on an outcome is different than the effect of each variable individually. This concept is often used in research fields like psychology, sociology, and statistics, particularly in experiments and studies involving human behavior.

For example, suppose a study investigates how both stress levels and social support affect job satisfaction. The interactional effect would mean that the influence of stress on job satisfaction might change depending on whether someone has high or low social support. If social support is high, stress might have less of a negative impact on job satisfaction; if social support is low, stress could have a more significant negative impact.

Interactional effects are commonly analyzed using statistical methods like two-way ANOVA or regression analysis, which allow researchers to assess whether and how two or more variables interact to affect an outcome.

Types and Examples of Interactional Effects

I. **Moderation**: Here, one variable changes the strength or direction of the relationship between another independent variable and a dependent variable. For instance, in psychological research, a personality trait like resilience might moderate the effect of

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

life stress on mental health. Higher resilience could reduce the negative impact of stress on mental well-being, demonstrating an interactional effect.

II. **Synergistic or Amplifying Interactions**: This occurs when two variables together produce a greater effect than would be expected from their individual effects. For example, in health studies, smoking and high cholesterol levels may each independently increase the risk of heart disease, but together they might elevate the risk far beyond the sum of their separate effects.

III. **Antagonistic or Dampening Interactions**: In some cases, the combined effect of two variables is less than their individual impacts would predict. For example, in pharmaceutical studies, two drugs might have less of an effect together than when administered alone, which is called an antagonistic interaction.

IV. **Crossover Interactions**: These are cases where the direction of an effect changes based on the levels of the interacting variables. For example, consider a study on teaching methods where "interactive" versus "lecture-based" approaches are tested across different student personality types. Introverted students might perform better in lecture-based settings, while extroverted students perform better with interactive methods, showing a crossover interaction where the "best" teaching method depends on the personality of the students.

Why Interactional Effects Matter in Research

Understanding interactional effects allows researchers to:

I. **Develop targeted interventions**: For example, interventions in public health might be tailored to specific populations if certain risk factors only have significant effects under particular conditions.

II. **Create complex predictive models**: In fields like machine learning or econometrics, considering interactions among predictors helps refine models, making predictions more accurate and relevant to real-world contexts.

III. **Challenge oversimplified assumptions**: Without accounting for interactional effects, research might lead to misleading conclusions. For instance, if researchers assume that factors like stress and social support have additive effects on job satisfaction, they may miss out on the nuanced reality that social support might actually reduce the negative impact of stress, altering the overall outcome.

Statistical Methods for Interactional Effects

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

Detecting interactional effects requires statistical tests that allow for the examination of multiple variables at once, such as:

i. **Two-way ANOVA**: Useful for examining interaction effects when comparing groups based on two categorical independent variables.
ii. **Regression Analysis with Interaction Terms**: Interaction terms in a regression model allow for the investigation of how combinations of variables affect a dependent variable.
iii. **Multilevel Modeling**: Often used when data is nested (e.g., students within classrooms) and interactional effects might vary across levels of the hierarchy.

Practical Applications Across Fields

i. **Healthcare**: Understanding how lifestyle factors (like diet and exercise) interact with genetics can lead to personalized medicine.
ii. **Education**: Tailoring teaching methods based on an interaction of learning styles and content type to improve learning outcomes.
iii. **Marketing**: Marketers often look at how different customer segments respond to various advertising strategies, recognizing that the success of a strategy may depend on demographic factors.

Interactional effects add depth to data interpretation, offering insights that wouldn't be visible if variables were analyzed in isolation.

20. Correlation

Correlation is a statistical measure that describes the strength and direction of a relationship between two variables. It quantifies how changes in one variable are associated with changes in another variable. The correlation coefficient, typically represented by the letter r, ranges from -1 to +1, with values closer to ±1 indicating a stronger relationship.

Types of Correlation by Direction

1. **Positive Correlation**:

- In a positive correlation, as one variable increases, the other also increases. Similarly, as one variable decreases, the other decreases.
- **Example**: The relationship between height and weight in adults is often positively correlated: taller people tend to weigh more.

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

- **Correlation Coefficient**: $r > 0$.

2. **Negative Correlation**:

- In a negative correlation, as one variable increases, the other decreases.
- **Example**: There may be a negative correlation between the amount of exercise and body fat percentage; as exercise increases, body fat percentage often decreases.
- **Correlation Coefficient**: $r < 0$.

3. **Zero or No Correlation**:

- When there is no discernible relationship between the two variables, the correlation is zero or very close to zero.
- **Example**: The amount of time a person spends reading and their shoe size likely has no correlation.
- **Correlation Coefficient**: $r = 0$.

Types of Correlation by Strength

1. **Perfect Correlation**:

- A perfect correlation is an ideal situation where all points fall exactly on a straight line.
- **Perfect Positive Correlation**: $r = +1$; all points lie on a line with a positive slope.
- **Perfect Negative Correlation**: $r = -1$; all points lie on a line with a negative slope.

2. **Strong Correlation**:

- A strong correlation indicates that the points are closely clustered around a line, either positively or negatively sloped, though not perfectly aligned.
- **Example**: A correlation coefficient around $r = 0.8$ or $r = -0.8$ shows a strong relationship.

3. **Weak Correlation**:

- A weak correlation means that the points are more scattered around the line, indicating only a slight association.
- **Example**: Correlation coefficients around $r = 0.3$ or $r = -0.3$ suggest a weak relationship.

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

Types of Correlation Based on Nature

1. **Linear Correlation**:

- Linear correlation means that the relationship between the two variables can be represented by a straight line.
- **Example**: The relationship between study hours and test scores often follows a linear pattern; as study hours increase, test scores generally increase.
- **Measure**: Pearson's correlation coefficient (r) is typically used for linear relationships.

2. **Non-Linear (Curvilinear) Correlation**:

- In non-linear correlation, the relationship between variables is curved, meaning that a straight line cannot accurately describe the association.
- **Example**: The relationship between stress and performance often follows an inverted U-shape; performance may improve with stress up to a certain point but then declines as stress increases further.
- **Measure**: Spearman's rank correlation coefficient is used for non-linear associations.

Measures of Correlation

A. Pearson's Correlation Coefficient:

- Pearson's correlation measures the strength and direction of a **linear relationship** between two continuous variables.
- It assumes the data is normally distributed and is the most commonly used method.
- **Formula**:

$$r = \frac{\sum(X - \bar{X})(Y - \bar{Y})}{\sqrt{\sum(X - \bar{X})^2 \sum(Y - \bar{Y})^2}}$$

Interpretation:

- r = +1 : Perfect positive linear correlation.
- r = -1 : Perfect negative linear correlation.
- r = 0 : No linear correlation.

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

B. Spearman's Rank Correlation Coefficient:

- Spearman's rank correlation assesses the **monotonic relationship** between two variables and is used for ordinal, ranked, or non-normally distributed data.
- It does not require the relationship to be linear and is less sensitive to outliers.
- **Formula**:

$$r_s = 1 - \frac{6 \sum d_i^2}{n(n^2 - 1)}$$

Where:

- d_i = the difference between the ranks of corresponding values of X and Y,
- n = the number of paired observations.

Interpretation: Values close to +1 or -1 indicate strong relationships, while values close to 0 indicate weak or no relationship.

C. Kendall's Tau:

- Kendall's Tau is another non-parametric measure of correlation used when data does not meet the assumptions for Pearson's or Spearman's correlation.
- It compares the number of concordant and discordant pairs, providing an estimate of the **strength and direction** of the association.
- Often used with small sample sizes or ordinal data.

Importance of Correlation in Statistics

- **Predictive Modeling**: Correlation helps identify relationships that can be used to predict one variable based on another.
- **Data Analysis**: It helps in understanding relationships between variables, especially when exploring or visualizing data.
- **Multivariate Analysis**: Correlation is a key component in regression analysis, factor analysis, and principal component analysis.
- **Risk Management**: In finance, correlation is used to manage portfolio risks by diversifying investments across assets with low or negative correlations.

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

Limitations of Correlation

- **Correlation Does Not Imply Causation**: A high correlation does not mean that changes in one variable cause changes in the other. Other factors or variables may influence both.
- **Sensitive to Outliers**: Correlation coefficients, especially Pearson's, are sensitive to outliers, which can distort the strength and direction of the relationship.
- **Restricted to Linear Relationships**: Pearson's correlation only measures linear relationships, so it may not be suitable for non-linear associations.

In summary, correlation is a vital tool in statistics for identifying the relationship between variables, with various types and measures depending on the data and nature of the relationship. Understanding these relationships is essential in fields like psychology, finance, biology, and more, where identifying patterns and predicting outcomes are key.

21. Two-Way ANOVA

Two-way ANOVA (Analysis of Variance) is a statistical technique used to determine the effects of two independent variables on a single dependent variable and to assess whether there is an interaction between these independent variables. Unlike a one-way ANOVA, which tests the impact of a single independent variable, a two-way ANOVA allows researchers to evaluate the main effects of each independent variable and any interaction effects between them.

Key Concepts in Two-Way ANOVA

I. **Main Effects**: These are the individual impacts of each independent variable on the dependent variable. For example, if a researcher is testing the effects of both "teaching method" (lecture-based vs. interactive) and "student gender" (male vs. female) on "test scores," the main effects would be:
   - How the teaching method affects test scores, regardless of gender.
   - How gender affects test scores, regardless of the teaching method.

II. **Interaction Effects**: These occur if the effect of one independent variable on the dependent variable changes depending on the level of the other independent variable. In the example above, there would be an interaction effect if the teaching method impacts test scores differently for male and female students.

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

III. **Factorial Design**: Two-way ANOVA uses a factorial design, often denoted as 2 x 2, 3 x 3, etc., depending on the number of levels of each independent variable. A 2 x 2 design, for instance, would have two levels for each of the two variables (e.g., two teaching methods and two genders).

Steps in Conducting a Two-Way ANOVA

I. **Define Hypotheses**: Specify hypotheses for main effects (e.g., "teaching method has an effect on test scores") and interaction effects (e.g., "teaching method and gender interact to affect test scores").
II. **Collect Data**: Ensure random assignment and a representative sample, meeting the assumptions of ANOVA (normality, homogeneity of variances).
III. **Conduct the Analysis**: Use statistical software to calculate F-values for each main effect and interaction, comparing them to a critical value to assess significance.
IV. **Interpret Results**: Determine if each main effect and the interaction effect are statistically significant.

Merits of Two-Way ANOVA

I. **Efficient Analysis of Multiple Factors**: By analyzing two factors at once, two-way ANOVA provides a more comprehensive view of data than separate one-way ANOVAs, reducing the need for multiple tests and thereby minimizing the risk of Type I errors (false positives).
II. **Detection of Interaction Effects**: Two-way ANOVA can reveal if the relationship between the independent variables and the dependent variable changes under different conditions, providing insights into complex relationships.
III. **Greater Statistical Power**: Including two independent variables often increases the power of the test, as it can better explain variability in the dependent variable compared to a single-factor analysis.
IV. **Generalizability**: Analyzing multiple factors simultaneously allows researchers to assess the consistency of an effect across conditions, leading to results that are more generalizable.

Demerits of Two-Way ANOVA

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

I. **Complex Interpretation**: Interaction effects can be challenging to interpret, especially if the interaction is significant but the main effects are not. Researchers need to carefully examine and explain what the interaction means in the context of the study.
II. **Increased Sample Size Requirements**: To achieve reliable results, two-way ANOVA often requires a larger sample size, especially when both factors have multiple levels. Small samples may lead to unreliable or misleading conclusions.
III. **Assumptions of ANOVA**: Two-way ANOVA assumes normally distributed data, homogeneity of variances (equal variances across groups), and independent observations. Violations of these assumptions can lead to incorrect conclusions.
IV. **Limitations with Unequal Group Sizes**: When group sizes are unequal, two-way ANOVA may yield biased estimates. Special statistical adjustments are sometimes needed, which can complicate the analysis.
V. **Not Suitable for Non-Linear Relationships**: Two-way ANOVA is limited to testing linear relationships. If the relationship between variables is non-linear, other techniques like regression analysis with polynomial terms might be more appropriate.

Practical Applications of Two-Way ANOVA

Two-way ANOVA is commonly used in fields like psychology, education, and biology. For example:

I. **Education**: Comparing the impact of different teaching methods across various age groups on learning outcomes.
II. **Marketing**: Analyzing the effectiveness of advertising strategies across different demographics.
III. **Healthcare**: Examining how treatment type and dosage level affect recovery rates in patients.

Two-way ANOVA is a powerful tool for understanding how two factors, individually and interactively, affect a dependent variable. While it offers the advantage of a comprehensive analysis of complex data relationships, it also requires careful interpretation and attention to assumptions, especially when interaction effects are involved.

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

Achiever's Hive

This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963


Dear learners,

To understand Numerical Part of Statistics, Kindly visit the link of the playlist given :

https://youtube.com/playlist?list=PLXu-INR7XVTvPK_Ak4EAYE51U93_bGdR0&si=2aapRfVaB916dbL3

or You can also visit on our Channel : Achiever's Hive


Then go to the playlist section, here you will find a playlist with name MPC 006 Important Q/A Series (A picture as shown below is attached for your convenience). The video lessons look like this :




You will find all VIDEO LESSONS of the numerical part as mentioned in the index. It is done to make your learning easier.


This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963


1. Mann-Whitney U Test






This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963







This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963


2. Chi square






This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963







This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963


3. Regression Equation






This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963






This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

4. Kendall's Tau




	A	B	C	D	E	F
X	2	7	1	5	8	10
Y	4	5	6	8	10	9



	X	Y	R_x	R_y	CR_x	CR_y	C	D
A	2	4	2	1	1 - C	3 - C	3	2
B	7	5	4	2	2 - A	1 - A	4	0
C	1	6	1	3	3 - D	4 - D	2	1
D	5	8	3	4	4 - B	2 - B	2	0
E	8	10	5	6	5 - E	6 - E	0	1
F	10	9	6	5	6 - F	5 - F	0	0
							11	4


This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963




Achiever's


This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

5. Spearman's Rho






This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963




Achiever's Hive


This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

6. Pearson Product Moment Correlation






This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963




Achiever's


This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963


7. One Way Anova - 1







This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963







This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963






This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963






This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963




Achiever's Hive


This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963


8. One Way Anova – 2







This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963





This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963







This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963


Achiever's Hive


This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963







This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963






This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963







This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963

9. t- Test





This document is licensed to: Ms. J. Niranjana, Hyderabad, 91-9246336336
https://www.youtube.com/@AchieversHive - Contact us (WhatsApp): +91-8979564963



Achiever's Hive

Connect with us on:

YouTube: www.youtube.com/achievershive

Telegram: t.me/achievershive

Instagram: Achiever's Hive