


Heterogeneity of plasminogen activator activity produced by two murine tumour cell lines.

National Library of Canada - Clonal heterogeneity in plasminogen activator activity produced by two murine tumor cell lines



Description: -

-Heterogeneity of plasminogen activator activity produced by two murine tumour cell lines.

-

Canadian theses = -- Thèses canadiennes Heterogeneity of plasminogen activator activity produced by two murine tumour cell lines.

Notes: Thesis (M.Sc.) -- University of Toronto, 1995.

This edition was published in 1994



Filesize: 30.88 MB

Tags: #Articles #de #revues #: #« #Proteolytic #stability #»

Imperial Cancer Research Fund British Association for Cancer Research. Joint symposium held at the Royal College of Surgeons, Lincolns Inn Fields, London 30 November

GLYCOLIPIDS AND MALIGNANCY; A REVIEW OF SOME CURRENT IDEAS.

Invasion and Metastasis

Furthermore, while several inducible switches are currently available, none of them are regulated by FDA-approved molecules. The method may further comprise detecting an amplicon produced by the amplifying.

Clonal heterogeneity in plasminogen activator activity produced by two murine tumor cell lines

Parallel progression of primary tumours and metastases. Sepharose-lentil and Sepharose-ricin lectin columns were used to purify glycoMedicine, Cardiff. In addition, the ECM serves as a repository for secreted regulatory proteins and growth factors.

Articles de revues : « Proteolytic stability »

ALLEN, Department of Ultrastructure, Paterson Laboratories, Christie Hospital and Holt Radium Institute, Manchester. Natl of those mechanisms which normally regulate Sci. DURHAM, Departwith those immunologically of ment of Clinical Oncology, University of lymphocytes and fibroblasts wereerythrocytes, detected on Glasgow.

Heterogeneity of plasminogen activator expression in various moloney virus-induced tumor cell lines. Lack of correlation with tumor growth and cell phenotype, International Journal of Cancer

This in vitro selection of an aneuploid clone usually occurs within one year of serial passage.

Related Books

- [Da fisica da matéria à metafísica do espírito](#)
- [Nayōbāi kāngōen kānkhlang 2498-2502 \[1955-1959\] - phūnthān khwāmcharōen thāng sētthakit n](#)
- [Safe way to glory, in several exercises of general use.](#)
- [Cha teau Gaillard e tudes de castellologie me die vale, XIV - actes du colloque international tenu a](#)
- [Revolución teatral del Barroco](#)