Type-material of Central and South American clearwing moths (Lepidoptera, Sesiidae)

Smithsonian Institution Press - Biblioteca Virtual BIO



Description: -

Elections -- India.

Indian National Congress.

Clearwing moths -- Type specimens.

Clearwing moths -- South America -- Classification.

Clearwing moths -- Central America -- Classification.type-material of

Central and South American clearwing moths (Lepidoptera, Sesiidae)

-

no. 261.

Smithsonian contributions to zoology;

Smithsonian contributions to zoology; no. 261type-material of

Central and South American clearwing moths (Lepidoptera, Sesiidae)

Notes: Bibliography: p. 27-28. This edition was published in 1978



Filesize: 8.19 MB

Tags: #Cat #81: #Lepidoptera

Clearwing Moths (Lepidoptera: Sesiidae)

There are ten genera and 22 named species in North America; BC reports three genera and four species -- Hübner attacks apple and other rosaceous trees; L. Pagbubuod: Ang biyolohiya sa kamuraang gulang ng Zethera musides Semper na inalagaan sa laboratoryo ay bahagyang ipinakita sa pamamagitan ng iniitlog ng isang babaeng ispesimen na inobserbahan sa probinsiya ng Negros, Republika ng Pilipinas. Hind tibia with erect and elongate scales on upper surface; adults moving with characteristic scuttling run.

Butterflies and moths in the Sesiidae, Oecophoridae, and Lycaenidae Families

Gorbunov 1992: Revision of the types of the Sesiidae Lepidoptera, preserved in the collection of the Zoological Museum of Kiev State University.

Research Articles

Holcocera immaculella McDunnough attacks the cones and seeds of pines. Key words: biology, Catonephele, Ectima, egg, Eunica, Euphorbiaceae, Hamadryas, Heliconius, hostplant, immature stages, larva, morphology, Myscelia, Neotropical, pupa, South America, Venezuela. Al comparar el numero total de especies presentes por mes contra el diagrama ombrotermico, se encontro relacion entre la epoca humeda y alta riqueza especifica y entre la epoca seca y baja riqueza.

Butterflies and moths in the Sesiidae, Oecophoridae, and Lycaenidae Families

Dorow 1990: Neue Funde von Paranthrene tabaniformis ROTTEMBURG, 1775 Lepidoptera: Sesiidae. Certain genera and even subfamilies of longhorned beetles associate with or even require living hosts for larval development, and others associate with dead hosts 65. Révision des Aegeriidae de Barbarie.

Lepidoptera

Sesiidae: Bilan d'une saison de chasse aux phéromones Lepidoptera. How To Pronounce Mesolia oraculella How To Pronounce Mesolia pandavella How To Pronounce Mesolia pelopa How To Pronounce Mesolia plurimellus. Tropical Lepidoptera 2 1 : 43-51.

Erich Bauer, Antiquariat Goecke & Evers

The nymphs and adults have plaguelike aggregations when they feed. There are 10 species in 10 genera in BC. The bodies are striped lengthwise in alternating whitish green and darker green lines, with several dorsal maroon blotches.

lepidopteran

Cockerell 1888: Sesia myopaeformis in the Isle of Thanet.

Related Books

- Letterpress, printing & bookbinding plant and machinery.
 Sexe en mal damour de la révolution sexuelle à la régression érotique
 British and Foreign Unitarian Association. Address.
- Birtrānd Rāsil fikruhu wa-mawqi'uhu fi al-falsafah al-mu'āṣirah
- Nekotorye aktual'nye voprosy istorii i ekonomiki stran zarubezhnogo Vostoka sbornik stateĭ