

Birth to three - developmental learning and the handicapped child

Teaching Resources - Maternal depression and child development



Description: -

-
Notaries -- United States -- Encyclopedias.
Portuguese language -- Dictionaries -- English.
Commerce -- Dictionaries -- Portuguese.
Plant products industry -- Dictionaries -- Portuguese.
Singer, Paul, 1844-1911.
Bears -- Fiction.
Developmentally disabled children -- United States -- Family relationships.
Education, Preschool.
Developmentally disabled children -- Care -- United States.
Developmentally disabled children -- Education -- United States.
Birth to three - developmental learning and the handicapped child
-Birth to three - developmental learning and the handicapped child
Notes: Bibliography p. 287-295.
This edition was published in 1979



Filesize: 64.38 MB

Tags: #Maternal #depression #and #child #development

Identification and Evaluation of Mental Retardation

Children of depressed mothers appeared more passively noncompliant, with less mature expressions of age-appropriate autonomy. A simple, reliable infant development screening tool.

Identification and Evaluation of Mental Retardation

As they become adolescents and young adults, most of them will need professional intervention to help them become their own advocates in the health care system.

Identification and Evaluation of Mental Retardation

Families should be supported as they encourage independent functioning on the part of their adolescent or young-adult son or daughter. Evaluation of the child with idiopathic mental retardation. These disorders reflect an abnormality during the first 25 weeks of gestation in the early migration of the neurons into the normally six-layered cortex.

Maternal depression and child development

They are also poor models for negative mood regulation and problem solving. The boy could stand for a few seconds without support and was able to take a few steps. Monotherapy is preferable, and the medication used in pregnancy should be continued in the postnatal period.

Identification and Evaluation of Mental Retardation

In their 2002 statement, the United States Preventive Services Task Force recommended screening adults for depression in clinical practices that have systems in place to assure accurate diagnosis, effective treatment and follow-up. Class of recommendation: B, Level of evidence: I. The mother had subsequently given birth to a second child with the same syndrome. Family therapy. School-age children and adolescents from families with a depressed parent may benefit from a family-centered intervention, focusing on communication about the illness within the family and on the

development of resiliency in the child.

Maternal depression and child development

In school-age children and adolescents, the presence of difficulties in child adjustment and impaired functioning at home and in school should alert the physician to the possibility of maternal depression. Because postpartum depression can have long-term effects on mothers and children and its peak prevalence occurs at approximately three months, it has been suggested to screen for postpartum depression at the two-, six- and 12-month well-baby care visits.

Maternal depression and child development

Other risk factors for postpartum depression include a history of mood disorders, depression symptoms during the pregnancy and a family history of psychiatric disorders.

Related Books

- [Philosophy of Yei selfhelp project](#)
- [Communication and interpersonal relations - text and cases](#)
- [El estatuto de los ex presidentes y ex miembros de gobierno](#)
- [Hubert Harrison reader](#)
- [Companion to Greek and Roman political thought](#)