


Epilepsy and the oral manifestations of phenytoin therapy

S. Karger - DailyMed



Description: -

-
Physicians Assistants.
Allied Health Personnel.
Medical assistants.
Human rights advocacy -- History -- 20th century
Anticonvulsants -- Side effects
Phenytoin -- Side effects.
Epilepsy -- Chemotherapy. Epilepsy and the oral manifestations of phenytoin therapy
-
ABF working paper -- #2028.
Monographs in oral science -- vol. 9 Epilepsy and the oral manifestations of phenytoin therapy
Notes: Includes bibliographical references.
This edition was published in 1981



Filesize: 6.87 MB

Tags: #Phenytoin: #medicine #to #treat #epilepsy #and #trigeminal #neuralgia

Epilepsy and the oral manifestations of phenytoin therapy (eBook, 1981) [ne-x.uni.rf.gd]

A thorough understanding of the pathogenesis of this unwanted side effect is essential if we are to devise appropriate regimens for its prevention and treatment.

The effects of antiepileptic drugs on oral health

Reactions include nystagmus, ataxia, slurred speech, decreased coordination, somnolence, and mental confusion. If it's a problem, ask your pharmacist to recommend a painkiller. If that does not work, you may need to switch to a different medicine.

Epilepsy and the oral manifestations of phenytoin therapy (1981 edition)

Taking the amount of seizure medicine the doctor prescribes and taking it at the right time every single day are the most important steps in preventing seizures! Phenytoin slows down these electrical signals to stop seizures. One fairly common side effect in children is overgrowth of the gums.

Epilepsy and the oral manifestations of phenytoin therapy (eBook, 1981) [ne-x.uni.rf.gd]

TGF- β is stored within the cell as a homodimer, noncovalently bound to a protein called latency-associated protein LAP, which maintains TGF- β inactive. Thus, modulation of apoptosis could contribute to fibrosis in gingival tissues. Phenytoin oral suspension should be discontinued immediately if symptoms of angioedema, such as facial, perioral, or upper airway swelling occur.

Phenytoin: MedlinePlus Drug Information

There have been some reports of side effects in breastfed babies, but in most cases babies do not get any side effects.

Phenytoin

If you change your epilepsy medicine, your doctor will tell you whether you need to stop driving and for how long. What if I forget to take it? The patient may become comatose and hypotensive.

Phenytoin

Digital Library Federation, December 2002. Do not freeze the liquid. This means that it may be difficult to achieve a constant, appropriate level.

Related Books

- [Thalāth rasā'il fiqhīyah](#)
- [Munāza'āt al-awqāf wa-al-aḥkār - nizām al-waqf, al-istihqāq fi al-waqf...](#)
- [Secret corner - ...a little bit of heaven on Cornwall's rugged coast.](#)
- [Regved.](#)
- [Domain of inquiry](#)