


Ordovician trilobites from the Keele Range, northwestern Yukon Territory

Dept. of Energy, Mines and Resources - Trilobites from the Lower Member of the Gualcamayo Formation



Description: -

-Ordovician trilobites from the Keele Range, northwestern Yukon Territory

-Ordovician trilobites from the Keele Range, northwestern Yukon Territory

Notes: 11

This edition was published in -



Filesize: 17.55 MB

Tags: #Ordovician #trilobites #from #the #lower #part #of #the #Dawan #Formation #(Floian

Detrital geochronology of pre

Geological Survey of Canada, Bulletin 142; 76 pp.

Palaeontographica Canadiana Archives

The base of the Clarence River group is marked by the prominent Ordovician chert interval that is distributed throughout Yukon and Alaska. Geological Survey of Canada Bulletin 396; pp. Burgess Shale type fossils from a Lower Cambrian shallow shelf.

List of fossil sites

Geological Survey of Canada, Bulletin 125; 82 pp.

Ordovician trilobites from the lower part of the Dawan Formation (Floian

PRINGLE - On the fossil Xiphosura from the South Wales Coalfield with a note on the myriapod Euphoberia. Norsk Polarinstitutts Skrifter, 162: 1—125.

Related Books

- [Richard Raynal, solitary](#)
- [I‘lām al-siyāsī wa-al-Islām](#)
- [Castell de Castelldefels - arqueologia, història, art](#)
- [1848 - a szabadságharc története levelekben, ahogyan a kortársak látták](#)
- [Czechoslovakia = Tsche coslovaquie = Checoslovaquia.](#)