

Mechanisms of resistance to plant diseases

M. Nijhoff/W. Junk - Mechanisms of quantitative disease resistance in plants

Description: -

- Literary Criticism & Collections / Ancient & Classical
- Literary Collections / Ancient, Classical & Medieval
- Language Arts & Disciplines-General
- Greek literature--History and criticism
- General
- Ancient and Classical
- Language
- Ancient (Classical) Greek
- Literature - Classics / Criticism
- Language Arts & Disciplines
- Literary studies: classical, early & medieval
- Plants -- Disease and pest resistance.Mechanisms of resistance to plant diseases

- AAB 17

Advances in agricultural biotechnology ;Mechanisms of resistance to plant diseases

Notes: Includes bibliographies and index.

This edition was published in 1985



Filesize: 46.108 MB

Tags: #Plant #Life: #Resistance #to #Plant

#Diseases

Plant Life: Resistance to Plant Diseases

Apart from viruses, bacteria, fungi, and certain insect pests also tend to infect plants. Mechanisms of resistance in plants to infection by pathogens, n.J.

Mechanisms of quantitative disease resistance in plants

Some cultivars maintain resistance by limiting the production of certain chemicals that are essential nutrients for invading pathogens. It has a broad phenotype which has led to the successful evolution of these strains and hence an attack many plant species.

Mechanisms of Resistance to Plant Diseases

These viruses cause leaf mottling, stunted plant growth, and deformed inedible fruit and routinely reduce yields by 20—80%, depending on location and growing season. Natural products and plant disease resistance.

Mechanisms of plant resistance to viruses

Examples of well-characterized elicitors include PAMPs, avirulence effectors, Hrp proteins delivered by the type III secretion system, and fungal or oomycetal proteins, such as cryptogein and cerato-platanins Gurr and Rushton, 2005b; Table 21. Alternative splicing of transcripts encoding Toll-like plant resistance proteins — what's the functional relevance to innate immunity? In Arabidopsis plants, flagellin is recognized by the PRR FLS2 that upon flagellin perception activates signaling networks leading to the induction of immune responses Gomez-Gomez and Boller, 2002. Although there are not many studies on the CBL-CIPK signaling pathway in response to biological stress, it is not to say that it has a small effect.

Related Books

- [Flora centroamericana.](#)
- [Honor roll containing a pictorial record of the gallant and courageous men from Faribault County, Mi](#)
- [Immediate tasks of the Soviet government](#)
- [Risk and reason - safety, law, and the environment](#)
- [Practice of the British and French hospitals: viz. The Edinburgh, Military, and Naval Hospitals, LHo](#)