


Rhizomania - approaches to diagnosis and production of resistant cultivars

De Montfort University - RNAseq Analysis of Rhizomania



Description: -

-Rhizomania - approaches to diagnosis and production of resistant cultivars

-Rhizomania - approaches to diagnosis and production of resistant cultivars

Notes: Thesis (Ph.D) - De Montfort University, Leicester 1996.

This edition was published in 1996



Filesize: 48.87 MB

Tags: #Identification #of #rhizomania

Symptoms of Rhizomania, a new disease for Michigan red beets

Thus, the evidence of insertion seems to be coupled with the susceptible genotype.

In vitro propagation of three mosaic disease resistant cassava cultivars

Disease Cycle Beet necrotic yellow vein virus BNYVV is the causal agent of rhizomania. In a further aspect, the present invention also relates to a method for mediating or increasing a resistance to a pathogen, in particular BNYW, in a plant, preferably a plant of the genus Beta, which comprises a step of transforming a plant cell with the nucleic acid molecule according to the invention or the vector of the present invention. So sind bis jetzt weder hochauflösende Markerkarten noch verifizierte Kandidatengene für die genannten Rhizomania-Resistenzgene öffentlich verfügbar.

What is new with rhizomania and curly top management and effects of these viruses on storage

Breeding for resistance to rhizomania in sugar-beet *Beta vulgaris* L. Archives of Virology, 152, 59. Table VII-3 summarizes the more common diseases occurring on vegetables and general control recommendations.

Use of molecular markers in breeding for disease resistance in sugar beet (*Beta vulgaris* L.) : Rothamsted Research

Kocide are used as bactericides.

Identification of rhizomania

Generally, it was observed that 100% explants of all the cultivars evaluated sprouted within 2—3 days.

Chapter VII: Disease Management

Archives of Virology, 126, 329—335. The reactions were repeated at least twice, and only distinct, reproducible, polymorphic and well-resolved bands across all runs were considered for analysis.

Symptoms of Rhizomania, a new disease for Michigan red beets

Photo by Ben Werling, MSU Extension. Therefore, there is need to evaluate alternative propagation methods that are fast and tissue culture offers a feasible option.

Breeding for resistance to rhizomania in sugar beet: A review, Euphytica

Der Begriff umfasst auch ein synthetisch hergestelltes Nukleinsäuremolekül.

Related Books

- [Dynamical effects of dendritic structure on neural systems](#)
- [Last Liberal governments - the promised land, 1905-1910.](#)
- [How much risk? - a guide to understanding environmental health hazards](#)
- [Polski przemysł naftowy pod niemiecką okupacją w latach 1939-1945](#)
- [Tachymeter-Tafeln.](#)