

Study of staff and inpatients experiences of untoward incidents in the acute mental health setting.

Oxford Brookes University - Feeling safe or unsafe in psychiatric inpatient care, a hospital

Preventable adverse event	• Harm which caused to a patient due to a human or system error that could have been prevented.
Unpreventable adverse event	• Harm that cannot be prevented, since it occurs due to a complication that cannot be foreseen.
Potential adverse event	• Adverse event, such as hospital-acquired/nosocomial infections that could have been obviated.

Description: -

-study of staff and inpatients experiences of untoward incidents in the acute mental health setting.

-study of staff and inpatients experiences of untoward incidents in the acute mental health setting.

Notes: Thesis (M.Sc.) - Oxford Brookes University, Oxford, 2001.

This edition was published in 2001



Filesize: 30.83 MB

Tags: #Postincident #Review #of #Aggression #and #Violence #in #Mental #Health #Settings

NHS England » Improving care in mental health services

Patients are now being effectively supported and encouraged to engage effectively with community care teams and other care providers reducing their perceived dependency on inpatient services, which promotes the delivery of care in the least restrictive environment. The descriptions indicated that restrictive practices were more common than de-escalating practices when dealing with violent incidents. Nurses tended to be much angrier, frustrated, less tolerant and more affected by patient violence when it involved a patient with a personality disorder or substance misuse.

Postincident Review of Aggression and Violence in Mental Health Settings

A qualitative approach to postincident review was undertaken as part of the larger study.

A model to improve safety on acute inpatient mental health wards

The effect of clinician-patient alliance and communication on treatment adherence in mental health care: a systematic review. An evaluation by the Centre for Psychiatric Nursing at the University of Melbourne showed consistent use of the model in the first year, improvement in patient and staff safety, and a 36% reduction in seclusion use Fletcher et al, 2017. The medium security clinic had patients from surrounding catchment areas and also from other areas of Sweden.

The psychological impact of restraint in acute mental health settings: the experiences of staff and inpatients

Using this outcome definition, there were 69 unique outcomes reported across the studies. In summary, the studies reviewed suggest that CTI could be an effective method of reducing homelessness post-discharge and reducing early readmissions. We consider those results sustainable due to the large number of randomized excerpts.

Training healthcare assistants working in adult acute inpatient wards in Psychological First Aid: An implementation and evaluation study

The costs of replacing a nurse who leaves because of ill health are considerable, as are the costs of lost productivity due to presenteeism or absenteeism. This study explores the psychological impact of restraint for staff and patients who are involved in these procedures. Design and procedure This is an interview study focusing on the question of feeling safe or unsafe, with some optional open-ended questions.

Violence and aggression in psychiatric inpatient care in Sweden: a critical incident technique analysis of staff descriptions

The results from the pre-study showed significant differences between the two wards regarding documentation of staff-patient interactions in terms of categories of failed or successful attunement. During the time of the study, all clinics intended to provide single rooms for all patients.

Workplace Violence Toward Mental Healthcare Workers Employed in Psychiatric Wards

A nurse familiar with writing progress notes but who was not a part of the research team and the first author independently scored a sample of 90 excerpts using the SESPI.

Interventions to improve discharge from acute adult mental health inpatient care to the community: systematic review and narrative synthesis

Middelboe T, Schjødt T, Byrting K, Gjerris A. Furthermore, participants identified that organizations need to be open to the exploration of the phenomenon of violence and embrace an open, blame-free environment for nurses to feel comfortable reporting patient violence.

Related Books

- [Foot and mouth.](#)
- [Case hardened steels - microstructural and residual stress effects : proceedings of the symposium](#)
- [Northern Native Languages Project - final report](#)
- [Educación popular y formación docente de la Independencia al 23 de enero de 1958](#)
- [Woraus sind unsere Kleider? - ein ganzes Buch ist aus der Antwort geworden](#)