

Evaluation of a pharmacist documentation system for antibiotic appropriateness and patient outcome

National Library of Canada - A usability study to improve a clinical decision support system for the prescription of antibiotic drugs



Description: -

-evaluation of a pharmacist documentation system for antibiotic appropriateness and patient outcome

Canadian theses = Thèses canadiennes evaluation of a pharmacist documentation system for antibiotic appropriateness and patient outcome

Notes: Thesis (M.Sc.)--University of Toronto, 1993.

This edition was published in 1993



Filesize: 33.21 MB

Tags: #Summary #of #Core #Elements #for #Hospital #Antibiotic #Stewardship #Programs

Appropriateness of Surgical Antimicrobial Prophylaxis Practices in Australia

Prospective multicentre feasibility study of a quality of care indicator for intravenous to oral switch therapy with highly bioavailable antibiotics. In the US it has become customary to present a to the executive officers of the hospital administration.

Appropriateness of Surgical Antimicrobial Prophylaxis Practices in Australia

There is an assumption that if pharmacists provide evidence base advice on medications, doctors, in turn, will implement them. For example, a program will need to evaluate if policies and guidelines are being followed as expected processes and if interventions have improved patient outcomes and antibiotic use outcomes.

Antimicrobial stewardship

Though the primary goal of stewardship programs is to improve patient care, many studies have shown that they will pay for themselves, through savings in both antibiotic expenditures and indirect costs. Audit and feedback to reduce broad-spectrum antibiotic use among intensive care unit patients: a controlled interrupted time series analysis. These findings highlight the need for improvement in SAP prescribing and suggest potential targeted areas for action.

Outcome assessment of intervention on appropriateness of antibiotic use among geriatric patients: A prospective interventional study from a tertiary care referral hospital

Effect of a collaborative approach on the quality of prescribing for geriatric inpatients: a randomized, controlled trial. Number of patients who started with empirical systemic antibiotic therapy according to the local guideline Total number of patients who started with empirical systemic antibiotic therapy 6 Before starting systemic antibiotic therapy, at least 2 sets of blood cultures should be taken.

Appropriateness of Surgical Antimicrobial Prophylaxis Practices in Australia

We also found that time has been invested in the development of functionalities, which were not optimally used. Accountability The antibiotic stewardship program must have a designated leader or co-leaders who are accountable for program management and outcomes.

Related Books

- [A Month of Mystery - Book Two](#)
- [Crise allemande de la pensée française, 1870-1914.](#)
- [Bookcloth 1823-1980 - a study of early use and the rise of manufacture, Winterbottoms dominance of t](#)
- [6 \[i.e. Six\] février 1934](#)
- [Intukweensquaw of James Bay](#)