


# Heterogeneity of plasminogen activator activity produced by two murine tumour cell lines.

National Library of Canada - Heterogeneity of plasminogen activator expression in various moloney virus-induced tumor cell lines. Lack of correlation with tumor growth and cell phenotype, International Journal of Cancer



Description: -

-Heterogeneity of plasminogen activator activity produced by two murine tumour cell lines.

Canadian theses = -- Thèses canadiennes Heterogeneity of plasminogen activator activity produced by two murine tumour cell lines.

Notes: Thesis (M.Sc.) -- University of Toronto, 1995.

This edition was published in 1994



Filesize: 37.85 MB

Tags: #Articoli #di #riviste:

**Heterogeneity of plasminogen activator expression in various moloney virus-induced tumor cell lines. Lack of correlation with tumor growth and cell phenotype, International Journal of Cancer**

Lack of correlation with tumor growth and cell phenotype.

## Invasion and Metastasis

Reactivity with the same tumour extract in patients with preinvasive lesions of the THE ASSOCIATION OF HOST IM- cervix carcinoma in situ and dysplasia was MUNOGLUBULINS WITH SOLID less than that found in the benign control TUMOURS IN VIVO.

## Clonal heterogeneity in plasminogen activator activity produced by two murine tumor cell lines

Two sets of primers are used to flank the target nucleic acid. In 4 cases with microinvasive In order to ascertain w% Aether circulating lesions of the cervix, no reactivity was found. Tumor extension and cell morphology were investigated by histologic and cytomorphologic analysis.

## Articles de revues : « Proteolytic stability »

Micro-NMR for Rapid Molecular Analysis of Human Tumor Samples. Heussen, C and Dowdle, EB, 1980, Electrophoretic analysis of plasminogen activators in polyacrylamide gels containing sodium dodecyl sulfate and copolymerized substrates.

## Invasion and Metastasis

Recently, FOCMA was defined as a tumour-specific cell-surface antigen which is induced in cats by FeLV infection and may be coded for by the FeSV genome. Interstitial matrix provides the structure characteristic of connective tissues. The data illustrate the importance of consensus

cysteines in the secretion and proteolytic activity of ADAMTS13.

### **Clonal heterogeneity in plasminogen activator activity produced by two murine tumor cell lines**

CRITCHLEY, Department of Biochemistry, University of Leicester, Leicester. The urokinase plasminogen activator system: role in malignancy.

### **Articles de revues : « Proteolytic stability »**

After an antigen is bound to the external antigen recognition domain, CAR receptors cluster together and transmit an activation signal. The advent of major technological advances over the past decade in the field of sequencing and high-resolution analysis of disseminated cancer cells has made it possible to begin to address such complex biological questions. Alternatively or additionally, the subsequent results of a test performed by the device may be transmitted to another person or device.

### **Invasion and Metastasis**

The method of claim 1, wherein the sample collection unit is a slide. This provides a means for removing all affected tissue within a single surgery, which is especially beneficial when the risks of additional anesthesia or surgeries are confounded by comorbidities.

## Related Books

- [Pick for the IBM PC and compatibles](#)
- [Philosophical works of Frederick Pickles on cutting gentlemens dress](#)
- [Geography of the world](#)
- [Preliminary examination of halibut fishing grounds of the Pacific coast.](#)
- [Approaches to education for character - strategies for change in higher education](#)