

Category:Accipiter nisus

The proportion of nests used in the second or third breeding season after logging 55% did not differ significantly from that of 27 control nests 67%.

Translate nisus in French with contextual examples

The forest grass-snake *Elaphe longissima* is common amongst reptiles, and the birds, as well as the black woodpecker *Dryocopus martius*, the *Picus viridis* and the *Picus canis*, of the day-time birds of prey the sparrow-hawk *Accipiter nisus* is present, whilst a night-time one is the tawny owl *Otus scops*, a real rarity in the south. English 2000 examined the effect of the cyclodiene pesticide, dieldrin, on population growth rates in eurasian sparrowhawks *accipiter nisus* using life history data from two areas in Britain and were able to demonstrate that acute dieldrin poisoning increased the instantaneous mortality rate by 0.

Nest reuse by Goshawks after timber harvesting: Importance of distance to logging, remaining mature forest area and tree species composition

Spurvehøgen tager stort set ikke jagtbart vildt. Spurvehøgen lægger fra 4 til 7 æg afhængigt af hunnens alder og kondition.

Related Books

- [Nilai-nilai luhur budaya spiritual bangsa dalam ungkapan bahasa Jawa dan Batak - penyusun, Ratnawati](#)
- [Proceso de beatificación del maestro Juan de Ávila](#)
- [Deyimler, atasözleri, beyitler ve anlamdaş kelimeler](#)
- [Ateşten gömlek](#)
- [Groundwater Probability - Virden \(West Half\) West of Principal Meridian; Saskatchewan-Manitoba.](#)