

# Trauma rehabilitation

Lippincott Williams & Wilkins - Center of Excellence for Trauma Rehabilitation and Research

| Box No 1                 |                            |                                    |
|--------------------------|----------------------------|------------------------------------|
| Injury severity category | Initial Glasgow Coma Scale | Duration of post-traumatic amnesia |
| Mild                     | 12-15                      | Less than 24 hours                 |
| Moderate                 | 9-11                       | 1-7 days                           |
| Severe                   | 3-8                        | 1-4 weeks                          |

Description: -

-  
Wounds and Injuries -- rehabilitation  
Wounds and injuries -- Patients -- Rehabilitation  
Wounds and injuries -- TreatmentTrauma rehabilitation  
-Trauma rehabilitation

Notes: Includes bibliographical references and index.

This edition was published in 2006



Filesize: 4.56 MB

Tags: #Acute #Rehabilitation #after #Trauma: #Does #it #Really #Matter?

## Trauma Rehab

The Rehabilitation of People with Traumatic Brain Injury is a portable reference for resident physicians, physiatrists and other professionals interested in the treatment of patients who have sustained a traumatic brain injury. Propensity score matching was used to identify a control group of patients who were not discharged to an IRF. Be sure to discuss any concerns with your healthcare provider before rehab.

### Acute Rehabilitation after Trauma: Does it Really Matter?

All of the mental health professionals at Deer Hollow are experts in their field and used evidence-based treatment to assist patients in the road to recovery.

## Trauma Rehab

How do I get ready for rehab after traumatic brain injury? Having access to specialized resources and equipment is critical for severely injured patients. Total functional independence measure scores improved from 63.

### Acute Rehabilitation after Trauma: Does it Really Matter?

Every person's needs and abilities after TBI are different. We all need help, and here at Deer Hollow, we have the resources and care to help you learn to cope with your trauma-related stress. Traumatic brain injury TBI occurs when a sudden injury causes damage to your brain.

## Rehabilitation and after care

Goal: This center was created to explore the differences in rehabilitation diagnosis, therapy prescription, and interventions needed for each subtype of trauma patient e.

---

## Related Books

- [Europa-America Latina - nuove forme de cooperazione](#)
- [Notes on the butterflies of the genus \*Velamysta\* \(Ithomiidae\) with a key to the species by W.T.M. For](#)
- [Justice on the grass - a story of genocide and redemption](#)
- [Hostiensis summa avrea - ad vetustissimos codices summa fide diligentiaque nuno primum collata ...](#)
- [Socialist campaign book](#)