

Psychiatric disorders in obstetrics

F.A. Davis Co. - Obstetric and gynaecological conditions associated with psychiatric disorder

What's New in Obstetric Medicine?
9 Key Questions (and Answers) about
Substance Use and Psychiatric Disorders
in Pregnancy

Learning Objectives:

- Review the epidemiology of behavioral and mental disorders in pregnancy and their impact on perinatal outcomes
- Become familiar with recommendations for screening, intervention and treatment of substance use and psychiatric disease

Description: -

-

Childbirth -- Psychological aspects.

Pregnancy -- Complications. Psychiatric disorders in obstetrics

-Psychiatric disorders in obstetrics

Notes: Bibliography: p. 5-6.

This edition was published in 1967



Filesize: 37.92 MB

Tags: #Psychiatric #Disorders

Psychiatric Disorders

Many institutions use the PHQ-2, the first two questions, as the initial screen. In summary, we found a strong and consistent relationship between mental illness and obstetric complications. Depression is more common in female adolescents than in male adolescents.

Premenstrual Syndrome and Premenstrual Dysphoric Disorder

Perinatal Depression: Prevalence, Screening Accuracy, and Screening Outcomes. Studies show that women diagnosed with a premenstrual disorder do not have higher levels of estrogen or progesterone than the general population, and accepted explanations as to why some women could be more sensitive to fluctuations in these sex hormones are lacking.

Psychiatric Disorders

The prevalence of PMS is not associated with age, educational achievement, or employment status. All mesh complications requiring further surgical intervention should be recorded on the British Society of Urogynaecology BSUG national database. Some disorders or their treatments will affect the hypothalamic—pituitary—gonadal axis, causing anovulatory cycles and various menstrual disturbances such as secondary amenorrhea or abnormal uterine bleeding.

Obstetrics and Gynecology Patients: Menstrual Cycle, Pregnancy, and Postpartum

One or more of the following symptoms must be present: 1. Mothers should not reduce or discontinue medications on their own. Screening for depression and suicide also should include screening for nonsuicidal self-injury.

Premenstrual Syndrome and Premenstrual Dysphoric Disorder

Oral contraceptives with and without drospirenone seem to be effective at relieving abdominal bloating, mastalgia, headache, weight gain, and swelling of extremities.

Mental Health Disorders in Adolescents

In children and adolescents, the mood may be irritable rather than sad.

Related Books

- [Our criminal law = - \[Notre droit pénal\]](#)
- [Kinship in Europe - approaches to long-term developments \(1300-1900\)](#)
- [Rescued from Barry](#)
- [Pre-historic America](#)
- [From rumour to removal.](#)