


Tumors arising from the blood-vessels of the brain - angiomatic malformations and hemangioblastomas

Charles C. Thomas - 84841592



Description: -

-

Brain Neoplasms. Tumors arising from the blood-vessels of the brain - angiomatic malformations and hemangioblastomas

- Tumors arising from the blood-vessels of the brain - angiomatic malformations and hemangioblastomas

Notes: Includes bibliographies.

This edition was published in 1928



Filesize: 41.33 MB

Tags: #Cushing, #Harvey #1869

Tumors Arising from the Blood Vessels of the Brain: Angiomatic Malformations and Hemangioblastomas.

Epidural injections of steroids are considered safer, but cases of aseptic meningitis have been reported following this procedure as well. He had a temperature of 38. Some idea of the frequency of occurrence of drug-induced aseptic meningitis can be obtained from cases reported with drugs in specific therapeutic areas.

Drug

MRI may show diffuse supratentorial white matter abnormalities that clear up after recovery from aseptic meningitis. Prevention involves the avoidance of drugs suspected to produce aseptic meningitis in susceptible individuals.

Methods of Circulatory Arrest

Mean follow-up was 21 months. Does not prevent regrowth following subtotal excision. Ibuprofen is the most frequently implicated drug in drug-induced aseptic meningitis.

Methods of Circulatory Arrest

Ibuprofen should not be prescribed for patients with systemic lupus erythematosus because of the frequent occurrence of ibuprofen-induced aseptic meningitis with the background of this disease. Two further cases of trimethoprim-sulfamethoxazole-induced aseptic meningitis have been reported, and complete recovery occurred in both after discontinuation of the drug 23.

Related Books

- [Myologie, angéiologie, névrologie, topographie](#)
- [Temps sociaux](#)
- [La-dwags kyi yul glu sñan pa'i pi wan](#)
- [\[Kniga dlia vnuchek - puteshestvie na rodinu](#)
- [Guidelines for vaccine-preventable disease education](#)