


Programme to Prevent Malnutrition

The Ministry - Cost effectiveness of a community based prevention and treatment of acute malnutrition programme in Mumbai slums, India



Description: -
-Programme to Prevent Malnutrition
-Programme to Prevent Malnutrition
Notes: Caption title.
This edition was published in 1992



Filesize: 42.68 MB

Tags: #Community

Factors affecting malnutrition in children and the uptake of interventions to prevent the condition

While it is possible that the magnitude of these effects has decreased since the time of the GAO analysis, the most recent comprehensive literature review, published by USDA in October 2004, found that more recent studies have continued to find that WIC participation during pregnancy reduces the incidence of low birthweight and has a positive impact on other birth outcomes.

Food and Nutrition Programs: Reducing Hunger, Bolstering Nutrition

Washington, DC: United States Agency for International Development. These were classified as controls but were not matched by age or sex to the cases. D'Agostino RB 1998 Propensity score methods for bias reduction in the comparison of a treatment to a non-randomized control group.

Preventing Acute Malnutrition among Young Children in Crises: A Prospective Intervention Study in Niger

Disinfect soiled water with this free and easy technique using solar radiation.

Factors affecting malnutrition in children and the uptake of interventions to prevent the condition

Interventions to prevent malnutrition must target these underlying causes.

Cost effectiveness of a community based prevention and treatment of acute malnutrition programme in Mumbai slums, India

They received intensive training and were monitored and supervised closely by field supervisors based on real time performance tracking. Similarly, children with an episode of diarrhoea within the last 6 months had a higher odds of malnutrition Adjusted OR, 2.

Malnutrition

Uncertainty analysis was undertaken to develop scenarios with Dirichlet probabilities assigned to each possible outcome. Groups receiving cash

transfer were given nutritional information on purchasing food for the targeted child, which included education on diet diversity e.

Malnutrition

The Food Stamp Program reduces poverty. A clinical trial in Malawi showed non-inferiority of SC+ for treating MAM in children aged 6—59 months, compared with two LNSs.

Related Books

- [British portraits - winter exhibition, 1956-57.](#)
- [Handbook of reporting methods](#)
- [Mental game](#)
- [Public interest profiles, 1992-1993](#)
- [Competition laws of NAFTA, Canada, Mexico, and the United States - a practitioners guide](#)