


# Ifosfamide in cancer therapy - a comparison with cyclophosphamide

Universitätsverlag - Uroprotection in patients receiving cyclophosphamide and ifosfamide



Description: -

-  
 Art, Medieval -- Exhibitions  
 Art, Dutch -- Exhibitions  
 Unión Alimentaria Sanders S.A.  
 Personnel management -- Spain -- Alicante (Province)  
 Industrial management -- Spain -- Alicante (Province)  
 Human capital -- Spain -- Alicante (Province)  
 Cyclophosphamide.  
 Cancer -- Chemotherapy.  
 Antineoplastic agents. Ifosfamide in cancer therapy - a comparison with cyclophosphamide  
 -Ifosfamide in cancer therapy - a comparison with cyclophosphamide  
 Notes: Includes bibliographical references.  
 This edition was published in 1993



Filesize: 38.63 MB

Tags: #Ifosfamide #versus #cyclophosphamide #in #combination #drug #therapy #for #metastatic #breast #cancer

## Ifosfamide vs cyclophosphamide in cancer therapy

Ifosfamide and its metabolites are known to be substantially excreted by the kidney, and the risk of toxic reactions to this drug may be greater in patients with impaired renal function. Repeat every 3 weeks for 12 cycles. Baxter and Vialflex are trademarks of Baxter International Inc.

## Uroprotection in patients receiving cyclophosphamide and ifosfamide

Includes cases reported as phlebitis and irritation of the venous walls.

## Bone Cancer Treatment Regimens

Response rates were 85% 95% CI 77—93% in the CIP arm and 90% 95% CI 84—96% in the CEP arm; complete response rates were 48 and 52%.

## A phase II randomised clinical trial comparing cisplatin, paclitaxel and ifosfamide with cisplatin, paclitaxel and epirubicin in newly diagnosed advanced epithelial ovarian cancer: long

Beyer-Boom ME, de Voogt HJ, Schaberg A 1978 The effect of cyclophosphamide treatment on the epithelium and stroma of the urinary bladder.  
 Allen LM, Creaven PJ 1972 In vitro activation of ifosfamide NSC-109724, a new oxazaphosphorine, by rat liver microsomes. Safety assessment of intensive induction with vincristine, ifosfamide, doxorubicin, and etoposide VIDE in the treatment of Ewing tumors in the EURO-E.

## Bone Cancer Treatment Regimens

After second-line chemotherapy laparoscopy was repeated, and the treatment was continued until pathological evidence of complete response. Injections are prepared for parenteral use by adding Sterile Water for Injection, USP or Bacteriostatic Water for Injection, USP benzyl alcohol or parabens preserved, to the vial and shaking to dissolve.



---

## Related Books

- [Vale un Perù](#)
- [Mémoires dun imbécile](#)
- [Celebration of literature and response - children, books, and teachers in K-8 classrooms](#)
- [Messerschmitt Bf 109 - Luftwaffe fighter](#)
- [Hanguksarül pinnaen myǿng nonsǒl](#)