

Northeastern forest survey board-foot volume equations

Dept. of Agriculture, Forest Service, Northeastern Forest Experiment Station - Development of monitoring and assessment of forest biomass and carbon storage in China



Description: -

-

Volume (Cubic content)

EquationsNortheastern forest survey board-foot volume equations

-

Communication arts books

Research note NE -- 271.

Research note NE -- 271.Northeastern forest survey board-foot volume equations

Notes: Includes bibliographical references (p. 3).

This edition was published in 1979



Filesize: 54.101 MB

Tags: #Forest #survey #cubic

[PDF] Northeastern Forest Survey Revised Cubic Foot Volume Equations D

Distribution of the equations with respect to different forest types Forest and vegetation classification of Indonesia has been developed by van Steenis , adapted by Whitmore and further refined by Kartawinata et al. The survey compiled 184 wood density values that included 113 species from 15 studies.

[PDF] Northeastern Forest Survey Revised Cubic Foot Volume Equations D

The last chapter tells how to obtain copies of the EWDB for a State. FIA samples land to estimate forest attributes, such as tree species composition, volumes of trees, and growth, removals, and mortality of trees, along with other variables, such as type of ownership.

A review of forest and tree plantation biomass equations in Indonesia

This field should have an entry if DBHCUR is 1. In contrast to the existing CRM branch CRs, which approached 100 per cent of AGB in small trees, branch biomass under the new formulation did not exceed 50 per cent of AGB in any of the eastern conifers observed here Figure F.

Big BAF sampling in mixed species forest structures of northeastern North America: influence of count and measure BAF under cost constraints

The majority of the equations were from plantations crops and agroforestry systems.

A review of forest and tree plantation biomass equations in Indonesia

In 1979, he transferred to the Forest Service and has been working with the Forest Inventory and Analysis Unit at the Northeastern Station since. X X X 401 water hickory *Carya aquatica* 14 4 X X X.

Improved accuracy of aboveground biomass and carbon estimates for live trees in forests of the eastern United States

While the NB FDS provides detailed overstory stratification, only four broad mature forest types were used in the analyses presented here: tolerant hardwood mixtures THMIX ; intolerant hardwood mixtures IHMIX ; softwood mixtures SWMIX ; and spruce — balsam fir SPBF. As long as mBAF was greater than about double cBAF, optimal sampling error varied very little, only as mBAF got close to cBAF did optimal sampling errors increase Figure A and C. Timberland Forest land that is producing or capable of producing crops of industrial wood.

Related Books

- [Ritorno del legame sociale](#)
- [Asyl - Anspruch und Wirklichkeit](#)
- [Kandinsky, Franz Marc, August Macke - drawings and watercolours.](#)
- [Ishkāfiyat al-fikr al-Zaydi fi al-Yaman al-mu‘āṣir - qirā’ah fi al-qirā’āt al-sab‘ li-tura](#)
- [Rosses - and other poems](#)