

Relationship between family size and maternal and child health.

- - **Maternal profiles and social determinants of malnutrition and the MDGs: What have we learnt?**

Variables	Total
(N=384) (%)	
Had mothers ever heard any MCH information?	
Yes	341 88.8
No	43 11.2
Sources of MCH information	
1. Media (TV, radio, and/or smart phone)	174 45.3
2. Health personnel at health facilities (Doctor, nurse, midwife, and/or other health workers)	306 79.7
3. Health personnel at community level (Doctor, nurse, midwife, and/or other health workers together with village health volunteer and local authority)	222 57.8
4. Social network (family, relatives, friends, and/or neighbors)	186 48.4
(a) Data shown as number (%) unless indicated.	
(b) Multiple answers from mothers who had ever heard any MCH information from each source.	

Description: -

-

Pakistan -- Economic conditions.

Pakistan -- Commerce.

Exports -- Pakistan.

Pressure groups -- Pakistan.

Infants -- Health and hygiene.

Women -- Health and hygiene.

Family size.relationship between family size and maternal and child health.

-relationship between family size and maternal and child health.

Notes: Bibliography: p. 25-[27]

This edition was published in 1970



Filesize: 57.38 MB

Tags: #Family #Planning's #Role #in #Improving #Maternal #and #Child #Health_Patrici...

[Impact of family planning on maternal

Immediate and continuous postpartum contact with the baby promotes bonding by facilitating sensitive maternal responsiveness to infant signals, thus facilitating maternal behavior which helps the infant become securely attached. Looking ahead to a post-2015 agenda for Sustainable Development Goals these results indicate that progress on the health of mothers and children will depend on multiple underlying determinants improving at the same time. A line of best fit curve is added showing a similar slope for both years, as well as a shift rightward and downward in 2010 compared to 1990.

Title V Maternal and Child Health Services Block Grant Program

In all three, trained staff can help parents get ongoing care for their children. The World Health Organization recommends that mothers breastfeed their children for the first two years of life, whereas the American Academy of Pediatrics and the American Academy of Family Physicians recommend that mothers do so for at least the first six months, and continue as long as is mutually desired. A conference or teaching room where parents can meet with staff members for discussion periods should be available.

Determinants of patterns of maternal and child health service utilization in a rural community in south eastern Nigeria

Center for Disease Control and Prevention: Office of Smoking and Health. A specialized wooden device that is, an infantometer was used to take the measurement in a supine position. Competing interests The authors declare that they have no competing interests.

Frontiers

Métodos de medición de las desigualdades. .

Related Books

- [New kid](#)
- [Georg Trakl - eine Biographie mit Bildern, Texten und Dokumenten](#)
- [Imperial Lady - A Fantasy of Han China](#)
- [Saddest summer of Samuel S.](#)
- [Piémont-Ligurie roman](#)