


# Mechanism of molecular interactions at metal filament surfaces - a study of the interaction of hydrocarbons with tungsten and rhenium filament surfaces using mass spectrometric and magnetron techniques.

The author - ECS Transactions, Volume 25, Number 8, September 2009, 2009



Description: -

-mechanism of molecular interactions at metal filament surfaces - a study of the interaction of hydrocarbons with tungsten and rhenium filament surfaces using mass spectrometric and magnetron techniques.

-

Thesesmechanism of molecular interactions at metal filament surfaces - a study of the interaction of hydrocarbons with tungsten and rhenium filament surfaces using mass spectrometric and magnetron techniques.

Notes: Ph.D. thesis. Typescript.

This edition was published in 1975



Filesize: 25.19 MB

Tags: #CVD #Diamond #Growth

**Review: mass spectrometry in Russia.**

The volatility and thermal stability of a series of hafnium complexes with guanidines and amidinate ligand systems have been investigated. The design method involves an evolutionary algorithm based on CFD simulations. The approach is based on the generation of a universal scale for the chromatography data using a multiple point normalization method.

## Publications of the CVD Diamond Group

The process energy especially during the flame treatment serves additional as a heating source for the substrate. In the instrument the gas ion source with the reduced discrimination and adjustable width of an output collimating split and the ion detector with a wide dynamic range of measured ion currents were used. An electrical surface resistance value of 8.

ECS Transactions, Volume 25, Number 8, September 2009, 2009

The IST films grown below 225oC exhibit an amorphous structure and the films grown at 300oC include various crystalline phases of IST, In-Sb, and In-Te.

## CVD Diamond Growth

As was noted in the Introduction, the first example of ESI volatile compounds. Poly-silicon deposition rate profiles on silicon wafers in several temperatures around 873 K along axial direction in commercial low-pressure chemical vapor deposition LPCVD reactor were investigated by both experiments and numerical simulations considering coupled computational fluid dynamics CFD and chemical reactions.

## **Publications of the CVD Diamond Group**

The dielectric properties have been evaluated at microscopic as well as at nanoscopic scale by scanning impedance microscopy.

## Related Books

- [Treasure-trove - the law and practice of antiquity.](#)
- [Mrs Fix-It easy home repair](#)
- [Communication from the Commission to the Council on a European Communities programme concerning the](#)
- [Acceso de la Argentina a la sociedad del conocimiento](#)
- [American art at home in Britain - the last four decades](#)