


Stratigraphy, igneous petrology and evolution of the Sinclair Group in southern south west Africa.

University of Cape Town - Stratigraphy, igneous petrology and evolution of the Sinclair Group in southern south west Africa. (edition)



Description: -

-
Geology - South Africa - Sinclair Group
Stratigraphy, igneous petrology and evolution of the Sinclair Group in southern south west Africa.

-
Cape Town University. Bulletin 16
Stratigraphy, igneous petrology and evolution of the Sinclair Group in southern south west Africa.

Notes: 11

This edition was published in -



Filesize: 30.27 MB

Tags: #Petrogenesis #of #the #felsic #rock #units #of #the #late

First evidence of the Cambrian basement in Upper Peninsula of Thailand and its implication for crustal and tectonic evolution of the Sibumasu terrane

Journal of Asian Earth Sciences 36, 168—182.

Isotopic evolution of the Middle to Late Proterozoic Awasib Mountain terrain in southern Namibia

Radiometric ages reveal a distinct younging along the rift from the southern toward the northeastern branch.

The intracratonic branch of the Damara Orogen in South West Africa II. Discussion of relationships with the Pan

Rb—Sr, Sm—Nd, and U—Pb geochronology of the rocks within the Khlong Marui shear zone, southern Thailand. On discute la genèse des unités felsitiques et on considère que les magmas se sont formés par fusion partielle de la croûte sialique; la source thermique ayant provoqué la fusion peut être recherchée dans des injections répétées de magma basique pendant l'évolution de la série. Both provinces appear to have evolved from earlier Archaean terrains, i.

First evidence of the Cambrian basement in Upper Peninsula of Thailand and its implication for crustal and tectonic evolution of the Sibumasu terrane

Es werden durchschnittliche Haupt- und Spurenelementanalysen vorgelegt, die in Ca-reiche und Ca-arme Gruppen gegliedert werden können, jede mit unterschiedlichen Ca-Wert-Bereichen und Spurenelementverteilungs-Muster. The geophysical signature of the rift is dominated by positive Bouguer gravity anomalies, flanked by gravity lows which follow a linear trend. Basic rocks with ophiolitic affinities are confined to the highest thrust sheet.

The Koras—Sinclair—Ghanzi rift in southern Africa. Volcanism, sedimentation, age relationships and geophysical signature of a late

middle proterozoic rift system

The role of strike-slip faults in the displacement of the Palaeotethys suture zone in Southeast Thailand. The join between the two can therefore not be regarded as a triple junction in the plate tectonic sense. That includes the Tengchong—Baoshan terrane, the Greater Himalayan sequence GHS, Lesser Himalayan sequence LHS, the Lhasa terrane, the Namche Barwa Complex, Iran, and Turkey Fig.

Petrogenesis of the felsic rock units of the late

Amalgamating eastern Gondwana: the evolution of the Circum-Indian Orogens. The Hf model age 1244 Ma to 1827 Ma and wide range of ϵ_{HfT} values between +4.

The Koras—Sinclair—Ghanzi rift in southern Africa. Volcanism, sedimentation, age relationships and geophysical signature of a late middle proterozoic rift system

Geochronology and geochemistry of granite magmatism in Thailand in relation to a plate tectonic model. A geodynamic model is proposed for the evolution of grabens, aulacogens, ensialic geosynclines and regions with tectono-thermal overprinting interpreting these structures as different reactions of the heterogeneous crust to mantle diapirs of different sizes.

Related Books

- [Goblin market](#)
- [Truss and his island](#)
- [Aja Taiheiyō Sensō](#)
- [Special situations in financial planning](#)
- [Taking Root \(Rookie Read-About Science\)](#)