

Gibbon and siamang - a series of volumes on the lesser apes.

S. Karger - Alternative splicing in lecithin:cholesterol acyltransferase mRNA: An evolutionary paradigm in humans and great apes — Johns Hopkins University



Description: -

-

China -- Relations -- United States

United States -- Relations -- China

Siamang

Gibbons Gibbon and siamang - a series of volumes on the lesser apes.

-Gibbon and siamang - a series of volumes on the lesser apes.

Notes: Includes bibliography.

This edition was published in 1972



Filesize: 38.29 MB

Tags: #Muscle #moment #arms #and #function #of #the #siamang #forelimb #during #brachiation

[PDF] Activity Budgets of Two Captive White

A climatic change occurs such that areas of wet lowland rainforest gradually became both cooler and drier. The frequency and duration ranges of all types and classes of song vocalisations are summarised in Table 3.

Geissmann, 2000: Duet songs of the siamang (*Hylobates syndactylus*)

Evolution of communication in gibbons Hylobatidae. In another siamang group in Sumatra, the subadult male was twice observed producing solo songs within the territory of his family group before his dispersal Palombit 1992:319. The first attempts of great call sequences during a song bout usually consist of a few long barks of the female.

Pair Bonds in Monogamous Apes: a Comparison of the Siamang *Hylobates Syndactylus* and the White

This has prevented testing of the hypothesis that pair bonds differ in the siamang *Hylobates syndactylus* and the white-handed gibbon *Hylobates lar* Chivers, 1972. So even though our hypothetical ancestor is, for the time being, still the same size and eating the same quality diet, the energetic costs associated with the procurement of this diet are increasing. Acknowledgements For permission to study the animals in their care and for useful information about the animals' previous history I wish to thank Dr.

Evolution of Gibbons and Siamang

By 5 y of age, this figure has decreased to 44% and by adulthood, it is 23%. In other instances, one animal was observed to grunt alone for some time before it was joined by its partner. Structure and organisation Thomas Geissmann This section was published as: Geissmann, T.

[PDF] Activity Budgets of Two Captive White

Apparently, most songs are produced either without any recognisable external stimulus or in response to songs of neighboring groups.

Orangutan odyssey

Universals of human music are difficult to identify but probably include a steady, accentuated beat see Arom, Nettle, and Mâche, this volume. Gut kinetics refers to the pattern of movement of ingesta, both particulate and liquid, through the digestive tract ,.

Gibbon take: DNA study reveals king of the swingers

In Studen, the original constellation consisted of one solitary female Vr , one pair Bb+Ra and one family group Ko+Cr +Li+Al. Figure 6 Sonagrams of female great calls of two gibbon species, *Hylobates lar* a and H. Feeding and Ranging Behavior of the White-Handed Gibbon *Hylobates lar* in Khao Yai National Park, Thailand, Unpublished PhD Dissertation, Washington University, St.

Related Books

- [Zavode - antropogeografska istraživanja](#)
- [Principles of radiation protection - a textbook of health physics.](#)
- [Politik und Kultur in der russischen Provinz - Nowgorod - Woronesh - Saratow - Jekaterinburg](#)
- [Tà chān pè hūà lǒn thā](#)
- [Continuemos!](#)