


Relations among quality of life, social support and long-term clinical outcome for adults living with HIV in the era of highly active antiretroviral therapy

National Library of Canada - Factors Related to Quality of Life in Treatment



Description: -

-Relations among quality of life, social support and long-term clinical outcome for adults living with HIV in the era of highly active antiretroviral therapy

-

Canadian theses = -- Th`eses canadiennes Relations among quality of life, social support and long-term clinical outcome for adults living with HIV in the era of highly active antiretroviral therapy

Notes: Thesis (M.Sc.) -- University of Toronto, 2002.

This edition was published in 2002



Filesize: 42.42 MB

Tags: #Social #support #and #quality #of #life #over #time #among #adults #living #with #HIV #in #the #HAART #era

Quality of life and sexual health among transgender people and people living with HIV in Sweden

Longitudinal analysis of clinical markers following antiretroviral therapy initiated during acute or early HIV Type I infection. Wig N, Lekshmi R, Pal H, Ahuja V, Mittal CM, Agarwal SK.

Quality of life of People living with HIV and AIDS attending the Antiretroviral Clinic, University College Hospital, Nigeria

Emotional control, pain, sleep and depression among HIV-positive persons. The Higher CD4 levels, the better ART adherence, the higher QOL of patient's.

Quality of life and sexual health among transgender people and people living with HIV in Sweden

The statistics R^2 and adjusted R^2 were used to verify the percentage of variance related to the increase in QOL explained by the model. Respondents took between 30 and 45 minutes to fill out the questionnaire.

SciELO

Quality of life, psychosocial health, and antiretroviral therapy among HIV-positive women in Zimbabwe. In conclusion, ART improved HRQoL of HIV-infected patients after one year of treatment.

Factors Related to Quality of Life in Treatment

It allows us to gain a full appreciation of the different ways individuals adapt to a situation they perceive to be stressful.

Perceptions of quality of life among Ugandan patients living with HIV: a qualitative study

The mean age of male participants was higher than that of their female counterparts, whereas the mean CD4 T cell count of female participants was higher than that of their male counterparts. This instrument assesses three dimensions of primary appraisal: threat, challenge and centrality.

Quality of life in HIV/AIDS :Basavaraj K H, Navya M A, Rashmi R, Indian J Sex Transm Dis

With the current prevalence of 4.

Related Books

- [Ring for Jeeves.](#)
- [Expert system applied to pollution control.](#)
- [Nishida and Western philosophy](#)
- [Socialism affluence - choice and the Welfare state.](#)
- [Education and practice - the critical interface](#)