


Study of adenosine triphosphate-sensitive potassium channels in rat hearts

De Montfort University - Effect of adenosine triphosphate



Description: -

-study of adenosine triphosphate-sensitive potassium channels in rat hearts

-study of adenosine triphosphate-sensitive potassium channels in rat hearts

Notes: Thesis (Ph.D) - De Montfort University, Leicester 1996.

This edition was published in 1996



Filesize: 68.72 MB

Tags: #Activation #of #SUR2B/Kir6.1 #subtype #of #adenosine #triphosphate

Inhibition of vascular adenosine triphosphate

Coronary arterial P O₂ was increased during halothane, although this effect was statistically significant only in the presence of glibenclamide. This approach avoided the systemic hemodynamic effects of the drugs, which simplified interpretation of the findings. This blockade was reversible, and the channel activities were restored by washing out glibenclamide.

Ischemic preconditioning with opening of mitochondrial adenosine triphosphate

THORNTON, in, 2005 K ATP Channel Hyperinsulinism K ATP channel HI is the most common form of HI.

Ischemic preconditioning with opening of mitochondrial adenosine triphosphate

In addition, it has been reported that K ATP channels play an important role in ischemic preconditioning : A brief period of ischemia can make the heart more resistant to subsequent, more severe episodes of ischemia. Furthermore, it has been shown that various drugs, such as cromakalim, may modulate the opening of the K ATP channels. Statistical Analysis Student's t test for paired samples was used to assess effects of the volatile anesthetics, cromakalim, sodium nitroprusside, acetylcholine, adenosine, and glibenclamide relative to the predrug control value.

Activation of the adenosine triphosphate sensitive mitochondrial potassium channel is involved in the cardioprotective effect of isoflurane

Finally, we also found that systolic lengthening persisted when CBF was increased maximally with adenosine during intracoronary enflurane administration. Coronary Effects of Volatile Anesthetics The increases in CBF caused by the volatile anesthetics were accompanied by decreases in SS and MVO₂ reflecting a direct negative inotropic effect, and thus the values for O₂ extraction decreased markedly. However, some studies used experimental approaches that minimized these complicating factors.

Related Books

- [Boston book trade, 1790-1799.](#)
- [Reactive voting in Danish general elections 1971-1979 - a revisionist interpretation](#)
- [Minguo zhen xi duan kan duan kan. Chongqing juan. \(9 vols.\)](#)
- [Textos escolares de lengua materna en la enseñanza media argentina, 1900-1950](#)
- [Women in Islam - tradition and transition in the Middle East](#)