

Factors affecting the detection of psychiatric morbidity by general practitioners

University of Birmingham - Detection of child mental health disorders by general practitioners



Description: -

-Factors affecting the detection of psychiatric morbidity by general practitioners

-Factors affecting the detection of psychiatric morbidity by general practitioners

Notes: Thesis (M.Med.Sc.) - University of Birmingham, Department of Public Health and Epidemiology.

This edition was published in 1995



Filesize: 21.101 MB

Tags: #PSYCHIATRIC #MORBIDITY #AND #REFERRAL #IN #GENERAL #PRACTICE—A #SURVEY #OF #GENERAL #PRACTITIONERS #IN #BANGALORE #CITY

Factors contributing to the recognition of anxiety and depression in general practice

While GPs in the current study reported working slightly fewer direct patient hours per week than GPs in the Bettering the Evaluation and Care of Health BEACH study an ongoing program that continuously collects information about clinical activities in general practice clinics within Australia , other GP characteristics were similar. Psychiatric morbidity in an Italian general practice. A greater effort to increase awareness of the problem among primary care and mental health care providers.

GP detection of health risk factors among general practice patients at risk of primary and secondary stroke

In addition, rostering of patients and salary-based or hourly-fee compensation should be more promoted. Participants were identified as having this risk factor if they reported consuming more than two standard drinks on a typical day or more than four standard drinks on any drinking occasion.

Physical morbidity and mortality in people with mental illness

Subjects who completed the Health Monitor in 2012 and who scored high on the K10 were invited to participate. This would allow psychiatrists to develop better liaison with their fellow professionals and share responsibility with them, which undoubtedly would encourage good medical practice.

General practitioners' management of mental disorders: A rewarding practice with considerable obstacles

Política de saúde mental: revisão dos dois últimos anos. Kringos DS, Boerma WGW, Hutchinson A, van der Zee J, Groenewegen PP: The breadth of primary care: a systematic literature review of its core dimensions.

Detection of child mental health disorders by general practitioners

Patient factors associated with guideline-concordant treatment of anxiety and depression in primary care. Only 47% of the new patients who met Bedford College diagnostic criteria for anxiety, depression, or illdefined disorder had their psychiatric disorder recognized by their GP.

Detection of psychiatric morbidity in the primary medical care setting in Brazil

Although detection and management of modifiable lifestyle risk factors is important for all patients, it is likely to be particularly important for those with a history of stroke or heart disease.

Related Books

- [Christian videos for children and young people](#)
- [On the take - how Americas complicity with big business can endanger your health](#)
- [Droit du travail](#)
- [McLean scores again](#)
- [Metamathematik](#)