


Characterization of a topoisomerase II α gene rearrangement in adriamycin-resistant P388 leukemia

National Library of Canada - Introduction



Description: -

-Characterization of a topoisomerase II α gene rearrangement in adriamycin-resistant P388 leukemia

-

Denkmaltopographie Bundesrepublik Deutschland -- Bd. 12.1

News stand library pocket edition -- 21A

Canadian theses = Thèses canadiennes
Characterization of a topoisomerase II α gene rearrangement in adriamycin-resistant P388 leukemia

Notes: Thesis (M.Sc.)--University of Toronto, 1993.

This edition was published in 1993



Filesize: 50.39 MB

Tags: #Natural #and #Acquired #Resistance #to #Cancer #Therapies

Frontiers

Br J Cancer 1997 76 486 93 10.

DNA topoisomerase II mutations and resistance to anti-tumor drugs, BioEssays

A similar tendency towards decreased transgene expression levels was also observed following administration of vincristine. Although these drugs block the formation of glucosylceramide from ceramide , , , , direct interaction with GCS has not been demonstrated. ERCC1 and RRM1 gene expressions but not EGFR are predictive of shorter survival in advanced non-small-cell lung cancer treated with cisplatin and gemcitabine.

WikiGenes

In the past, the main effort was aimed at developing highly specific inhibitors acting on single RTKs. Other gene products that interact with topoisomerase II for chromosome segregation Focusing exclusively on topoisomerase II as the enzyme responsible for chromosome segregation at mitosis is too simplistic, given the high complexity of molecular interactions during cell division. Alterations in the levels or affinity of these enzymes in the cellular system develop the drug resistance.

DGIdb

Much progress has been made in identifying agents and developing strategies for enhancing cellular accumulation of topo II inhibitors in tumors with the MDR phenotype.

Characterization of camptothecin

Jacoby DR, Fraefel C and Breakefield XO.

Targeting Ceramide Metabolism—a Strategy for Overcoming Drug Resistance

Additional research or modeling is also needed to identify what combination of targets can be expected to optimize therapy for particular cancer types. Breast cancer stem cells: Obstacles to therapy. Structure activity relationship study suggested that phenol moiety at 4-position of the central pyridine regardless of chlorophenyl moiety at 2-position of the central pyridine has an important role in dual topoisomerase inhibitory activity as well as antiproliferative activity.

Related Books

- [Summary report on water pollution, Western Great Lakes drainage basin](#)
- [Economics of public finance - an economic analysis of government expenditure and revenue in the Unit](#)
- [Erziehungsidee und Schulwirklichkeit - zur pädagogischen Evaluation des Unterrichts](#)
- [Soil survey \(reconnaissance\), the Red River Valley area, Minnesota](#)
- [Sequencing of instruction in relation to hierarchies of competence](#)