


Pharmacology of pseudoaconitine and japaconitine considered in relation to that of aconitine

Published for the Royal Society by Dulau and Co - The Pharmacology of Pseudoaconitine and Japaconitine Considered in Relation to That of Aconitine. [Abstract] on JSTOR



Description: -

-
Pharmacology. pharmacology of pseudoaconitine and japaconitine considered in relation to that of aconitine

- pharmacology of pseudoaconitine and japaconitine considered in relation to that of aconitine

Notes: From the Philosophical transactions of the Royal Society of London, Series B. Vol 195.

This edition was published in 1902



Filesize: 57.58 MB

Tags: #Meridian #Institute

The Pharmacology of Pseudoaconitine and Japaconitine Considered in Relation to That of Aconitine. [Abstract] on JSTOR

Curacoa Aloes is obtained from A.

Aconitum (U. S. P.)—Aconite.

Outside calyx as seen from beneath. Aloin is slow in action, requiring from 10 to 12 hours. Poisoning: Have irritation of intestinal canal, causing pain, vomiting, and purging, cold sweats, prostration, sometimes convulsions, collapse.

Primitive luke

Uses Preparations of aconite and its principal alkaloid, aconitine, when applied to the skin, produce tingling followed by numbness; they are used in certain forms of neuralgia and rheumatism.

Meridian Institute

Pseudoaconitine, also known as nepaline C 36H 51NO 12 , is an extremely toxic found in high quantities in the of , also known as Indian Monkshood, which belongs to the family.

The Pharmacology of Pseudoaconitine and Japaconitine Considered in Relation to That of Aconitine : Cash, J. : Free Download, Borrow, and Streaming : Internet Archive

Aconitum Napellus, common monkshood, is a doubtful native of Britain, and is of therapeutic and toxicological importance. As in poisoning, the patient is conscious and clear-minded to the last.

Pseudaconitina

In aphthous conditions, with fever, associate it with phytolacca. It tapers gradually, and is either dark grey and nearly smooth daughter root or brownish, and marked with not very prominent, paler, longitudinal ridges parent root.

Aconitum

There are 50 Pinaceae species that yield this resin. Added to cimicifuga, it greatly increases the curative influence of this agent in acute rheumatism, and more especially where there is a tendency to muscular spasm. The shape of the flower is specially designed to attract and utilize bee visitors, especially the humble bee.

Aconitum

Hlavně vietnamské a čínské náplaste sú pomerne lacné. The underground portion of the plants are dug up after the stem has died down, and the smaller of the 'daughter' roots that have developed at the side of the old roots are selected for replanting in December or January to form new stock, the young roots being planted about a foot apart each way.

Related Books

- [Workplace crime - proceedings and resources of Internal Business Theft Conference of the National Co](#)
- [Taste of France](#)
- [Senshō haiku dan](#)
- [Adab al-nisāʾī al-Lubnānī al-muʿāṣir - dirāsāt, siyar, bibliyūghrāfiyā](#)
- [Smalltalk](#)