

Sub-clinical lead poisoning

Academic Press - Lead

Description: -

- California -- Biography.
- California -- Social conditions -- 19th century -- Sources.
- California -- Gold discoveries -- Sources.
- Gold miners -- California -- Biography.
- Pioneers -- California -- Biography.
- Frontier and pioneer life -- California -- Sources.
- Politics and government
- European Union
- 1974-
- International Relations - General
- History / Europe / General
- International Relations - Diplomacy
- International
- EU (European Union)
- Politics/International Relations
- Portugal
- Politics / Current Events
- International - Economics
- Political Science
- EU & European institutions
- Pennsylvania -- Politics and government -- 1951-
- Leader, George M., 1918-
- Fiction - General
- Fiction
- Fiction / General
- General
- Modern fiction
- Literature of special Lesbian interest
- Rembrandt Harmenszoon van Rijn, 1606-1669 -- Sources -- Indexes.
- Lead poisoning
- Sub-clinical lead poisoning
- Sub-clinical lead poisoning
- Notes: Bibliography: p. [168]-198.
- This edition was published in 1974



Tags: #Lead #poisoning #in #cattle

Subclinical lead poisoning: A preventable disease

At other times more than 20 cows have been lost in a single poisoning incident. The population reference laboratory normal ranges for thyroxine are set wide compared to the normal individual range, so that a fall in thyroxine levels at the lower end of the range may elevate the TSH above normal.

MEDICAL: CONDITIONS: LEAD POISONING : DATABASE SEARCH

RESULTS : BOOKS : INTERNET SOURCES: A Collection of Lead Poisoning Books, Database Search Results and Internet Sources

Lead poisoning can affect any cattle operation.

SUB-CLINICAL LEAD POISONING IN CHILDREN, The Journal of Child Psychology and Psychiatry

It is often difficult to distinguish lead poisoning from other diseases that affect the nervous system of cattle. The response to the ingestion of lead-containing materials is relatively slow so that a child with pica ingesting flakes of lead-based paint containing 1,000 µg of lead per day is unlikely to present with clinical symptoms for 4 - 5 months Chisolm and Harrison, 1956; Bartrop and Killala, 1967. Lead used to be in paint; older houses may still have lead paint.

Subclinical lead poisoning: A preventable disease

Boiled linseed oil, which contains lead, may poison livestock when it is used as a laxative.



Filesize: 65.12 MB

Related Books

- [RACER # 3605347](#)
- [Models for Writers 9e & Easy Writer 3e](#)
- [Demise of the reasonable man - a cross-cultural study of a legal concept](#)
- [Erlebt, gedacht, vollbracht - Erinnerungen an ein Jahrhundert](#)
- [Proverbes et dictions français](#)