


Netherlands health interview survey 1981-1991

SDU/CBS-Publications - The World Mental Health Survey Initiative



Description: -

- Netherlands -- Statistics, Medical.
Health surveys -- Netherlands. Netherlands health interview survey 1981-1991

-Netherlands health interview survey 1981-1991

Notes: Includes bibliographical references.

This edition was published in 1992



Filesize: 17.65 MB

Tags: #The #World #Mental #Health #Survey #Initiative

SSRS Full

In 2018, this ranged from 9. Determinants of suicidal ideation differed from determinants of suicide attempts. Table 4: Share of the population reporting that they had chronic depression, 2014 % Source: Eurostat In 15 of the EU Member States, self-perceived chronic depression peaked within the age group covering people aged 75 years and over.

Riots explode across Netherlands over covid restrictions

Dual Smoker For respondents who smoke BOTH cigarettes and bidis. A full-scale parliamentary enquiry is to be held into how thousands of innocent parents could have been accused of child benefit fraud.

Studies comparing vaccinated to unvaccinated populations

Plans are being drawn up to take coronavirus into account, and this may include allowing pupils to fail one more exam but still pass their certificate overall. By contrast, a higher share of the total number of deaths among men rather than women was attributed to mental and behavioural disorders in Slovenia, Poland, Estonia and Romania; there was no difference between the shares for men and women in Bulgaria. Comparability - over time Between EHIS wave 1 and 2 implementation, modifications in the model questions were limited in order to safeguard the comparability of the results over time.

Face

Also, with this type of interview, the interviewer can gain a deeper insight to specific answers by treating the questionnaire like a meaningful discussion and deducing the validity of each response.

Health Interview Survey

Emphasise your role and describe how appealing your present job is but do not overdo it; you are about to leave. The 17 participating EU Member States strived towards comparability via a standard questionnaire, guidelines and translation recommendations. Urinary incontinence, problems in controlling the bladder N.

Regarding the national implementation of EHIS wave 2, an overview of the practices adopted is presented below.

Surveys

Table 5: In-patient average length of stay for mental and behavioural disorders and Alzheimer's disease, 2013 and 2018 days Source: Eurostat , and Among 10 of the 25 EU Member States for which data are available no comparison available for Greece or Portugal , the average length of a hospital stay for people treated for mental and behavioural disorders fell between 2013 and 2018; the largest reductions were recorded in Finland and Malta 2013-2017 , down 14. The pattern in Iceland was almost the reverse of the general situation witnessed for the whole of the EU-27 insofar as the highest proportion of the population reporting chronic depression was recorded among those aged 15-24 years 21. The data of each person are ordered according to the value of the total equivalised disposable income.

Related Books

- [Primary health care in Nigeria - female circumcision / edited by Bernard E. Owumi.](#)
- [Immigration dans le roman francophone contemporain](#)
- [The political economy of rural livelihoods in transition economies - land, peasants and rural poverty](#)
- [United Nations microfiche standard.](#)
- [Carotid endarterectomy - principles and technique](#)