


Ecological study of some root aphids in Egypt (Hemiptera-Hemoptera: Aphididae)

General Organisation for Government Printing Offices - Cereal Aphids and their Biological Control Agents in Egypt



Description: -

-
Brangwyn, Frank.
Governmental investigations
Electric power-plants
Accounting
Special education -- Québec (Province)
Special education -- France.
Judgment (Logic)
Hemiptera -- Egypt.
Aphididae. ecological study of some root aphids in Egypt (Hemiptera-Hemoptera: Aphididae)
- ecological study of some root aphids in Egypt (Hemiptera-Hemoptera: Aphididae)
Notes: At head of title: United Arab Republic. Ministry of Agricultural Extension Dept., Editing and Publication [sic] Administration.
This edition was published in 1963



Filesize: 70.22 MB

Tags: #Hemiptera

Isolation and efficacy of the endophytic fungus, *Beauveria bassiana* (Bals.) Vuillemin on grapevine aphid, *Aphis illinoisensis* Shimer (Hemiptera: Aphididae) under laboratory conditions

The highest peak was by mid-February. Biology of the camphor scale and a method for predicting the time of appearance of stages in the field. Wingless immature forms are rather long-bodied and paler than adults, with a dark spinal stripe and a light dusting of whitish-grey wax.

Studies on the biology of some root aphids in Egypt (Hemiptera

The first appeared in the.

Effectiveness of ladybirds as biological control agents: Patterns and processes

Consequently, not all injury causes damage, and damage threshold, or boundary, defines the level of injury where damage occurs Bardner and Fletcher 1974 and Pedigo et al. Parasite populations can be monitored by assessing the proportion of aphid mummies relative to unparasitized aphids.

Isolation and efficacy of the endophytic fungus, *Beauveria bassiana* (Bals.) Vuillemin on grapevine aphid, *Aphis illinoisensis* Shimer (Hemiptera: Aphididae) under laboratory conditions

Parasites of potato-infesting aphids in north-eastern Maine. No adverse effects of transgenic- Bt tomato plants expressing the toxin Cry3Bb against Coleoptera were found on the biology of M.

Isolation and efficacy of the endophytic fungus, *Beauveria bassiana* (Bals.) Vuillemin on grapevine aphid, *Aphis illinoisensis* Shimer (Hemiptera: Aphididae) under laboratory conditions

Studies on the aphid fauna of Sinai Governorates.

Hemiptera

Journal of Chemical Ecology, 30 4 :819-834. The obtained 5 isolates were DNA sequenced for ITS region.

Hemiptera

In some groups of Hemiptera, there are variations of winged, short-winged, and wingless forms within a single species. Molecular Fingerprinting of Certain Cereal Aphids in Egypt Hemiptera: Sternorrhyncha: Aphididae Using RAPD and ISSRs Markers. Characteristics of the open empty mummies of the cabbage aphid *Brevicoryne brassicae* L.

Related Books

- [Kept in the dark](#)
- [Education in cities](#)
- [Colors from nature - growing, collecting, and using natural dyes](#)
- [Autoregulation of blood flow - proceedings of an international symposium held June 10, 11, 12, 1963,](#)
- [Kuttā te ādamī te hora kahānām](#)