

# Obstetric analgesia and anesthesia - a manual for medical students, physicians in training, midwives, nurses, and other health personnel

## World Federation of Societies of Anaesthesiologists - OBSTETRIC ANALGESIA AND ANESTHESIA IN GENERAL PRACTICE

European Union Standards for Training and Minimum Instruction for Anaesthetists (1980)

<ul style="list-style-type: none"> <li>Basic sociology and socio-medical questions</li> <li>Basic pharmacology</li> <li>Psychology</li> <li>Principles and methods of teaching</li> <li>Health and social legislation and health organization</li> <li>Professional ethics and professional legislation</li> <li>Sex education and family planning</li> <li>Legal protection of mother and infant</li> </ul>	<ul style="list-style-type: none"> <li>Analgesia, anesthesia and resuscitation</li> <li>Physiology and pathology of the newborn infant</li> <li>Care and supervision of the newborn infant</li> <li>Psychological and social factors</li> </ul>
--	---

**B. Practical and clinical training**  
This training is to be dispersed under appropriate supervision:

- Advising of pregnant women, involving at least 100 pre-natal examinations.
- Supervision and care of at least 40 pregnant women.
- Conduct by the student of at least 40 deliveries, where this number cannot be reached owing to the lack of available women in labour, it may be reduced to a minimum of 30, provided that the student assists with 50 further deliveries.
- Active participation with breech deliveries. Where this is not possible because of lack of breech deliveries, practice may be in a simulated situation.
- Performance of episiotomy and initiation into suturing. Initiation shall include theoretical instruction and clinical practice. The practice of suturing includes suturing of the wound following an episiotomy and a simple perineal laceration. This may be in a simulated situation if absolutely necessary.
- Supervision and care of 40 women at risk in pregnancy, or labour or post-natal period.
- Supervision and care (including examination) of at least 100 post-natal women and healthy new-born infants.
- Observation and care of the new-born requiring special care, including those born pre-term, post-term, underweight or ill.
- Care of women with pathological conditions in the fields of gynaecology and obstetrics.
- Initiation into care in the field of medicine and surgery. Initiation shall include theoretical instruction and clinical practice.

The theoretical and technical training (Part A of the training programme) shall be balanced and coordinated with the clinical training (Part B of the same programme) in such a way that the knowledge and experience listed in this Annex may be acquired in an adequate manner.

Clinical instruction shall take the form of supervised in-service training in hospital departments or other health services approved by the competent authorities or bodies. As part of this training, student midwives shall participate in the activities of the departments concerned in so far as those activities contribute to their training. They shall be taught the responsibilities involved in the activities of midwives.

Description: -

-

Cake.

Analgesia.

Anesthesia in obstetrics. Obstetric analgesia and anesthesia - a manual for medical students, physicians in training, midwives, nurses, and other health personnel

-Obstetric analgesia and anesthesia - a manual for medical students, physicians in training, midwives, nurses, and other health personnel

Notes: Bibliography: p. 191-192.

This edition was published in 1980



Filesize: 20.410 MB

Tags: #Safe #Prevention #of#the #Primary #Cesarean #Delivery

## OBSTETRIC ANALGESIA AND ANESTHESIA IN GENERAL PRACTICE

However, this risk increase may not be entirely related to the duration of the second stage per se, but rather to health care provider actions and interventions in response to it eg, operative delivery and the associated risks of perineal trauma. Epidural anesthesia is commonly used by Canadian women to relieve pain—even though it can result in prolonged labour and increase the need for forceps and vacuum birth.

### Obstetric Analgesia and Anesthesia

Footnote 74 Skoll A, St Louis P, Amiri N, Delisle M, Lalji S. Supporting Healthy And Normal Physiologic Childbirth: A Consensus Statement By ACNM, MANA, and NACPM. It is also important to recognize that recovery is a process and some families may not be ready at a particular time; offering the option of discussing their experience in the future is recommended.

### Obstetrics and Gynecology Clinical Practice Guidelines Summaries

As new cohorts of AA graduates become available to play a role in the delivery of anesthetic care, the CAS will evaluate, with the safety of patients and quality of care as the primary principles in mind, the pertinence of revisiting its guidelines.

### ARC 3708C

More recent studies also show that outcomes are also improved in preterm infants born before 24 weeks, and also for the late preterm 34 +0 to 37 +6 weeks.

## Related Books

- [Occidente, 1932-1935](#)
- [Shin-plasters.](#)
- [Gramatika bosanskoga jezika](#)
- [Profsoiuzy Kalmykii - stranitsy istorii](#)
- [Isaac Newton and physics for kids - his life and ideas with 21 activities](#)