

Thermodynamics of Rare-Earth Compounds.

s.n - Thermodynamic study of the rare earth vapour complexes: ScAl₂Cl₉ and YAl₂Cl₉

RE	$\Delta H_{\text{aq}}(700^\circ\text{C})$ kJ mol ⁻¹	$\Delta H^\circ_{\text{f,ox}}$ kJ mol ⁻¹	$\Delta H^\circ_{\text{f,el}}$ kJ mol ⁻¹
RHABDOPHANES			
LaPO ₄ · 0.804 H ₂ O	203.53 ± 1.95 (9)	-342.92 ± 4.29	-2220.9 ± 4.5
CePO ₄ · 0.732 H ₂ O	179.56 ± 0.93 (9)	-325.23 ± 7.21	-2189.7 ± 9.4
PrPO ₄ · 0.709 H ₂ O	199.67 ± 0.98 (9)	-321.54 ± 6.05*	-2181.7 ± 6.4*
	210.67 ± 2.32 (5)*		
NdPO ₄ · 0.746 H ₂ O	197.10 ± 1.22 (8)	-309.80 ± 4.57	-2178.7 ± 5.1
SmPO ₄ · 0.636 H ₂ O	195.88 ± 0.48 (8)	-311.13 ± 4.22	-2156.8 ± 5.1
EuPO ₄ · 0.555 H ₂ O	191.51 ± 0.91 (10)	-300.15 ± 4.07	-2042.4 ± 4.9
GdPO ₄ · 0.533 H ₂ O	186.76 ± 0.92 (8)	-304.45 ± 3.96	-2119.1 ± 7.2
MONAZITES			
LaPO ₄	151.26 ± 0.82 (11)	-346.11 ± 3.37	-1994.4 ± 4.3
CePO ₄	112.28 ± 0.84 (10)	-316.25 ± 6.46	-1963.8 ± 9.4
PrPO ₄	147.57 ± 0.96 (8)	-326.11 ± 8.13*	-1983.5 ± 6.3*
	163.31 ± 1.89 (7)*		
NdPO ₄	144.80 ± 1.33 (9)	-308.78 ± 3.77	-1964.7 ± 5.1
SmPO ₄	142.66 ± 1.54 (8)	-301.77 ± 3.36	-1965.7 ± 5.3
EuPO ₄	139.98 ± 1.15 (11)	-286.90 ± 2.56	-1870.6 ± 4.9
GdPO ₄	139.33 ± 0.96 (8)	-293.80 ± 1.86	-1956.1 ± 7.2

All errors in the table are propagated as two standard deviations of the means; numbers in brackets are numbers of individual measurements.
*drop solution enthalpy in lead borate at 800°C. Enthalpies of formation of Pr-containing rhabdophane and monazite from oxides are calculated using thermodynamic cycle from high temperature drop solution experiments with lead borate at 800°C.

Description: -

-Thermodynamics of Rare-Earth Compounds.

-

Report of investigations (United States. Bureau of Mines) --

5468Thermodynamics of Rare-Earth Compounds.

Notes: 1

This edition was published in 1959



Filesize: 38.81 MB

Tags: #Heat #capacity #and #thermodynamic #functions #of #some #rare

Thermodynamics of rare earth elements and uranium in gallium based quaternary metallic alloys

In addition, 15 mol L —1 H₃PO₄ 85% Normapur was used as the source of the phosphate anions. Willit JL, Miller WE, Battles JE 1992
Electrorefining of uranium and plutonium—a literature review.

Thermodynamic properties of rare-earth alloys by electrochemical emf measurements

They also contain significant amounts of uranium;. Johnson MB, James DD, Bourque A, Dabkowska HA, Gaulin BD, White MA. Your datasets will also be searchable on Mendeley Data Search, which includes nearly 11 million indexed datasets.

Thermodynamics of rare earth metal interaction with copper

Effects of BaO sintering additive on dielectric properties of Y₂Ti₂O₇ microwave dielectric ceramics.

Thermodynamic properties of rare-earth alloys by electrochemical emf measurements

They are thermodynamically metastable with respect to the corresponding monazites plus water at all temperatures under ambient pressure conditions. JCT Style guide Please consult the for further details on the requirements for submitting your paper to The Journal of Chemical Thermodynamics.

Thermodynamics of rare earth elements and uranium in gallium based quaternary metallic alloys

Electrooptic and piezoelectric properties of La₂Ti₂O₇ single crystal. Conclusion The enthalpies of formation of a series of rhabdophanes were measured by high temperature drop solution calorimetry and the energetics of dehydration and transformation to monazite plus water were determined by DSC.

Related Books

- [Història de la literatura catalana a Mallorca a l'Edat Mitjana](#)
- [200 tips for growing vegetables in the Pacific Northwest](#)
- [Schulpädagogik und Fachdidaktik - zur Aktualität des Werkes von Johannes Kühnel \(1869-1928\)](#)
- [Ring as a support to, and element of teaching methodology with special reference to the early years](#)
- [Report of Childrens Programmes Workshop](#)