

# Reinstatement of the turrid genus *Bellaspira* Conrad, 1868 (Mollusca: Gastropoda) with a review of the known species

Los Angeles County Museum of Natural History - ⓘ Gastropods described in 1868 .. Info

Description: -

-

Bereavement -- Religious aspects -- Christianity

Grief -- Religious aspects -- Christianity

Alvesta kommun (Sweden) -- Bibliography.

Canary Islands -- Imprints

Printers -- Canary Islands

Ninemile Creek (Missoula County, Mont.).

Groundwater -- Quality -- Montana -- Creeks.

Water quality biological assessment -- Montana -- Creeks.

Algae -- Montana -- Creeks.

English language -- Composition and exercises.

English language -- Grammar.

Mollusks -- Classification

Turridae, Fossil -- Classification

Turridae -- Classification  
Reinstatement of the turrid genus *Bellaspira* Conrad, 1868 (Mollusca: Gastropoda) with a review of the known species

-

Figli in cielo -- 2

6

Smålands bibliografi,

no. 189

Los Angeles County Museum. Contributions in science, Reinstatement of the turrid genus *Bellaspira* Conrad, 1868 (Mollusca: Gastropoda) with a review of the known species

Notes: Includes bibliographical references.

This edition was published in 1970

Tags: #*Bellaspira* #*clarionensis*

## List of smaller malacological Works

Rectification of the generic placement of *Sclewdoris tanyin* Marcus, 1971. Distally the pelvic process is narrow and straight and connected with its antimer by



Filesize: 62.11 MB

ligamentous tissue. The toothed surface is narrow, bearing for most of its length two rows of sharp pointed teeth.

## Full text of of the Natural History Museum

Traces of a cardinal process have not been observed. There are no ventral frontal laminae. *Adelatecrinus* is erected to accommodate *Atelecrinus sulcatus* AH Clark and *Adelatecrinus vallatus* new species.

## Research Publications

To show the equatorial layer separated into two parts by a median layer of shell material in the peripheral zone; 9, X 40 ; 12, X 20 ; in specimens previously identified as L.

## Research Publications

The olfactory tracts are well separated and each tract is short and thick comprising at least twelve separate nerves each of which branches to feed the individual laminae of the nasal rosette. The posterior border of the bone is rounded and partially overlapped by a relatively small, ovoid intercalar to which is attached the inferior limb of the posttemporal Figs 4B, 5A,B, 6D. The ossified anterior wall of the ethmoid forms most of the nasal cavity and a thin, vertical septum of ethmoid cartilage separates the cavities medially.

### **List of smaller malacological Works**

In the Bregmacerotidae the third and subsequent vertebrae bear parapophyses to which are attached cartilage-formed ribs Fig. Bathymetrically, the species range in depth from 36 to 2034 m, although they occur most frequently between 200 and 1000 m.

### **Bellaspira**

The longitudinal ribs are oblique. However, a wide size-range of anal maculae, their size negatively correlated with their number, occurs in the monotypic serranochromine genus *Pharyngochromis*, some individuals of which have spots as large as those in *Chetia brevis*. Proximal ray morphology varies substantially with size in *P.*

## Related Books

- [Proceedings of the Hague Peace conferences - translation of the original texts](#)
- [The 2007-2012 World Outlook for Dry Purchased Noodle Products of All Shapes, Sizes, and Types with L](#)
- [Practical forms of proceedings, under the laws of New Jersey.](#)
- [Who gets to narrate the world? - contending for the Christian story in an age of rivals](#)
- [Defence - Exchange of Notes Between Canada and the United Republic of Tanzania : Ottawa, December 19](#)