

Parasites of domestic and feral cats from the London area.

- - TWS Issue Statement: Feral and Free

Helminth infection	Domestic, 45 (24%)		Stray, 273 (76%)	
	Infected cats	Prevalence %	Infected cats	Prevalence %
Unique infections				
Toxocara cati	0	0	6	22
Toxocara leonae	0	0	1	34
Toxocara canis	1	12	8	29
Physalium proglutidum	4	47	9	33
Opisthorchis catenatus	25	284	79	29
Angiostrongylus vasorum	2	24	3	11
Trichostrongylus axei	0	0	0	0
Total unique infection	32		116	
Mixed infections				
Opisthorchis catenatus/Toxocara canis	0	0	6	22
Opisthorchis catenatus/Toxocara cati	0	0	4	15
Opisthorchis catenatus/Physalium proglutidum	3	33	8	3
Opisthorchis catenatus/Angiostrongylus vasorum	0	0	3	11
Opisthorchis catenatus/Toxocara canis/Toxocara leonae	1	12	0	0
Opisthorchis catenatus/Physalium proglutidum/Trichostrongylus axei	0	0	1	34
Angiostrongylus vasorum/Toxocara cati	1	12	0	0
Total mixed infections	5		22	

Description: -

-Parasites of domestic and feral cats from the London area.

-Parasites of domestic and feral cats from the London area.

Notes: Thesis (M.Phil.) - North East London Polytechnic, 1982.

This edition was published in 1982



Filesize: 35.43 MB

Tags: #Top #PDF #Parasites #of #domestic #owned #cats #in #Europe: #co #infestations #and #risk #factors

[Download PDF] Parasites of domestic and feral cats from the London area by Susan Nichol Ebook

They have vibrissae on the muzzle, eyebrows, and elbows which function as haptic receptors. Soon the problem becomes out of control.

PRIME PubMed

Estrus females call loudly to potential mates, while continually rolling on the ground. Cat fleas are found on a much wider phylogenetic diversity of wild mammals than dog fleas.

Prevalence of intestinal parasites in domestic cats from the London area

Toxoplasma gondii seroprevalence was 22. Show more 9 Giemsa stain was used in carrying out quantification of malaria parasites. Males tend to make territories more often than females.

Prevalence of zoonotic parasites in feral cats of Central Virginia, USA

Noise, destructive behaviours, excrement, biting and overpopulation of domestic dogs and cats are some of the common problems that can result in stressors and difficulties in owners, neighbours, the community and pets Cutt et al. One day later, after treatment, all animals were returned to their home environment SHE.

Related Books

- [Anekdoty, aforizmy i melochishki o damakh](#)
- [Social psychology of HIV infection](#)
- [Biofeedback - Making O](#)
- [China shakes the world](#)
- [Yugoslavia in crisis](#)