


Fetal well-being - physiological basis and methods of clinical assessment

CRC Press - Assessment of Fetal Well being



Description: -

-
Prenatal Diagnosis -- methods.
Pregnancy Complications -- prevention & control.
Fetal Diseases -- prevention & control.
Fetal monitoring.
Prenatal diagnosis. Fetal well-being - physiological basis and methods of clinical assessment
-Fetal well-being - physiological basis and methods of clinical assessment
Notes: Includes bibliographical references and index.
This edition was published in 1990



Filesize: 33.36 MB

Tags: #NCLEX

The Quality of Clinical Maternal and Neonatal Healthcare

Most clinical applications have included assessment of other fetal biophysical indices, such as heart rate. Amniocentesis results are available sooner than CVS results are.

Antepartum and intrapartum foetal monitoring

Tan P, Steinbach M, Kumar V. Concealed: Blood is retained in uterine cavity.

Well Being For Women

Assessment can help figure this out.

Antepartum and intrapartum foetal monitoring

These decelerations vary in size, shape, and relationship to uterine contractions.

Physiological Measurement

Impact of fetal blood sampling on vaginal delivery and neonatal outcome in deliveries complicated by pathologic fetal heart rate: a population based cohort study. Contraction stress test CST may be used as a confirmatory test when the NST is nonreactive or inadequate. They concluded that one acceleration was just as reliable as two in predicting healthy fetal status.

Antepartum and intrapartum foetal monitoring

Several planes including the abdominal view, four-chamber, five-chamber, short-axis and three-vessel views have to be assessed in order to get

spatial information on different cardiac chambers and vessels, as well as their connections to each other. The relatively low sub-sample of cases for the risk monitoring function is due to the fact that only 58 out of 82 cases 71% were actively monitored by partograph during stage one labor, and only 76 out of 82 cases 93% were kept in the maternity unit for direct immediate postpartum observation during the first two hours after delivery. Any diagnosis should have clinical utility, meaning it aids the mental health professional in determining prognosis, the treatment plan, and possible outcomes of treatment APA, 2013.

Chapter 19 Assessment of Fetal Well

Williams Obstetrics, 24th edition, McGraw-Hill Education. NLN Competencies: Relationship-Centered Care: Communicate information effectively; listen openly and cooperatively.

Well Being For Women

These chest wall movements differed from those following birth in that they were discontinuous. The new algorithm was compared to other known methods basing on the collected research material and defined performance measures. ST analysis for intrapartum fetal monitoring.

Related Books

- [Divinas mutantes - carta de relación del itinerario de la poesía femenina en México](#)
- [Kaisōshū ni yoru kindai bungaku](#)
- [Fatigue industrielle - colloque](#)
- [Haynes automotive heating & air conditioning systems manual - the Haynes repair manual for automotiv](#)
- [Landor - a biographical anthology. Compiled and edited by Herbert Van Thal. With an introd. by Mal](#)