


Measuring clinical outcome in stroke (acute care)

Clinical Effectiveness & Evaluation Unit, Royal College of Physicians - Performance measures for in



Description: -

- Natural history.

Buñuel, Luis, -- 1900-1983.

Critical care medicine.

Outcome assessment (Medical care)

Cerebrovascular disease -- Treatment -- Evaluation.

Cerebrovascular disease -- Patients -- Care. Measuring clinical

outcome in stroke (acute care)

-Measuring clinical outcome in stroke (acute care)

Notes: Includes bibliographical references.

This edition was published in 2000



Filesize: 64.12 MB

Tags: #In

Outcome Measurement in Rehabilitation

Symmetry in external work SEW : a novel method of quantifying gait differences between prosthetic feet.

Performance measures for in

Results underscore the importance of optimizing acute stroke care for all patients.

Performance measures for in

A history of hypertension was found in 77. A significant difference in death, length of stay and unplanned readmission occurred among units treating stroke. Using recombinant tissue plasminogen activator to treat acute ischemic stroke in China: analysis of the results from the Chinese National Stroke Registry CNSR Stroke.

Outcome Measurement in Rehabilitation

However, no difference was found by demographic characteristics and 30-day fatality between patients included in the sample and those not included. Health at a Glance 2001.

Stroke

We excluded 124 patients because their medical charts could not be found after five searches on different dates or because the charts contained insufficient information for diagnosis of an ischemic stroke Figure. Age disparities in stroke quality of care and delivery of health services. RC contributed to the study design, data collection and its monitoring and interpretation of the results and collaborated in the drafting of the manuscript.

Outcome Measurement in Rehabilitation

Proportions and 95% confidence intervals were calculated for the population characteristics and for the study variables. To check the quality of data collection, an external audit was carried out by blinded and independent researchers. Most patients 89% were independent in daily life before the current stroke onset.

In

Mean age was 70 years SD 12 years and 146 56% were male. Outcome measures for individuals with stroke: process and recommendations from the American Physical Therapy Association neurology section task force.

Related Books

- [Restauro e museografia - centralità della storia](#)
- [Dramatic works of Samuel Foote, Esq. - to which is prefixed a life of the author.](#)
- [The Arrandel Motto V3 - A Novel](#)
- [Treasury of Jewish humor](#)
- [A Moments Enchantment \(Candlelight Ecstasy Romance, No 404\)](#)