

Variations in family practitioners, rheumatologists and orthopaedic surgeons perceptions of the outcomes, alternatives and indications for knee replacement

National Library of Canada - Full text of the Medical Association of the State of



Description: -

-Variations in family practitioners, rheumatologists and orthopaedic surgeons perceptions of the outcomes, alternatives and indications for knee replacement

- Canadian theses = -- Thèses canadiennes Variations in family practitioners, rheumatologists and orthopaedic surgeons perceptions of the outcomes, alternatives and indications for knee replacement
Notes: Thesis (M.Sc.) -- University of Toronto, 1996.
This edition was published in 1996



Filesize: 5.82 MB

Tags: #The #Use #of #Guidelines #for #Managing #and #Treating #Osteoarthritis

Difference Between Rheumatologist & Orthopedic Doctor

For consulting a medical specialist, a yearly compulsory deductible is levied. Arthritis Care Res 1997; 10: 343—51.

Publications Facilitated by ACREU

The numeric pain scale is an 11-point scale for adults and children over 10 years old.

Does the implementation of a care pathway for patients with hip or knee osteoarthritis lead to fewer diagnostic imaging and referrals by general practitioners? A pre

More than 20,000 teenagers use the State Health Department system for preventive contraceptive services each year.

Pain Relief: Putting Pain Management in Your Hands

JULY 1981 JOURNAL OF THE MEDICAL ASSOCIATION OF THE STATE OF ALABAMA 23 Mothers, partly because of their greater contact with their children, often generate considerable hostility even if unconscious of this towards them. Pre and post surgical management of the hip and knee. However, there was no significant difference in the same parameters between the two groups at the second week 0.

Health Content A

All co-authors contributed to the interpretation of results and provided feedback on drafts of the manuscript. The secondary outcomes will be pain intensity; disability and function; kinesiophobia and global perceived effect at 3, 6, and 12 months after randomization; and kinesiophobia and global perceived effect at completion of treatment.

Full text of of the Medical Association of the State of

Soever and MacKay concluded that comparison of outpatient care to other rehabilitation approaches warranted further study to determine the optimal processes of care following TJA 80. Data on file, Wyeth Laboratories.

Related Books

- [Sheng biao mian bo yuan li ji qi zai dian zi ji shu zhong di ying yong](#)
- [Samotechnyĭ transport na khlebopriemnykh i zemoperarabatyvaiushchikh predpriatiakh](#)
- [\[Ĉhotmai čhāk Mŭ'ang Thai\] = - Letters from Thailand](#)
- [Canada](#)
- [Gideon Lincecum, 1793-1874 - a biography.](#)